

Boundary layers in a Riemann-Liouville fractional derivative two-point boundary value problem

José Luis Gracia and Martin Stynes

Abstract A two-point boundary value problem whose highest-order term is a Riemann-Liouville fractional derivative of order $\delta \in (1, 2)$ is considered on the interval $[0, 1]$. It is shown that the solution u of the problem lies in $C[0, 1]$ but not in $C^1[0, 1]$ because $u'(x)$ blows up as $x \rightarrow 0$ for each fixed value of δ . Furthermore, $u'(1)$ blows up as $\delta \rightarrow 1^+$ if and only if the constant convection coefficient b satisfies $b \geq 1$.

1 Introduction

Let $\delta \in (1, 2)$. Let $g \in C^1(0, 1]$ with $g' \in L_1[0, 1]$. The Riemann-Liouville fractional derivative D_{RL}^δ of order δ associated with the point $x = 0$ is defined by

$$D_{RL}^\delta g(x) = \frac{d^2}{dx^2} \left[\frac{1}{\Gamma(2-\delta)} \int_{t=0}^x (x-t)^{1-\delta} g(t) dt \right] \quad \text{for } 0 < x \leq 1;$$

see [6].

In this paper we shall consider the two-point boundary value problem

$$-D_{RL}^\delta u(x) + bu'(x) = f \quad \text{for } x \in (0, 1), \quad (1a)$$

$$u(0) = 0, \quad u(1) + \alpha_1 u'(1) = \gamma_1, \quad (1b)$$

José Luis Gracia

Institute of Mathematics and Applications, University of Zaragoza, Spain and Department of Applied Mathematics, University of Zaragoza, Spain, e-mail: jlgracia@unizar.es

Martin Stynes

Applied Mathematics Division, Beijing Computational Science Research Center, China and Department of Mathematics, National University of Ireland, Cork, e-mail: m.stynes@ucc.ie

where b, f, α_1, γ_1 are given constants. We assume that $\alpha_1 \geq 0$, as in 2nd-order elliptic problems. Remark 1 below explains the necessity of imposing the Dirichlet condition $u(0) = 0$ at $x = 0$.

Existence and uniqueness of solution to (1) is discussed in [4]. Assume that f and γ_1 are not both zero as otherwise the solution to (1) is $u \equiv 0$ in the set of functions $u \in C[0, 1]$, with u' and $D_{RL}^{\delta-1}u$ absolutely continuous on $[0, 1]$ (see [4, Theorem 2.8]).

Problem (1) is used to model anomalous diffusion processes; for example, we refer to [3] for a motivation of this model.

In [8] we considered a related problem where the Riemann-Liouville derivative of (1a) is replaced by a Caputo fractional derivative, and discussed under what circumstances one would observe a boundary layer in its solution at $x = 1$ as $\delta \rightarrow 1^+$ (with the other data of the problem fixed). Our main aim in the present paper is similar: to determine when $u'(1)$ blows up as $\delta \rightarrow 1^+$.

In Section 2 we solve (1) exactly using Laplace transforms. We shall see easily that in general $|u'(x)| \rightarrow \infty$ as $x \rightarrow 0$ for each fixed value of $\delta \in (1, 2)$, so $u \notin C^1[0, 1]$. A more demanding investigation in Section 3 exploits properties of Mittag-Leffler functions to show that $u'(1)$ blows up as $\delta \rightarrow 1^+$ when $b \geq 1$ but no such singular behaviour is present when $b < 1$.

Notation. We use the ‘‘big O’’ notation in its sharp form. Thus when we write for example $g = O(1/(\delta - 1))$ as $\delta \rightarrow 1^+$, we mean that $\lim_{\delta \rightarrow 1^+} [(\delta - 1)g]$ exists and is non-zero. Throughout the paper C denotes a generic constant that is independent of δ but may depend on b, f, α_1 and γ_1 . Set $\|u\|_\infty = \max_{x \in [0, 1]} |u(x)|$.

2 Solution via Laplace transform

We compute the solution of the problem (1) by using the Laplace transform. Our analysis makes heavy use of the two-parameter Mittag-Leffler function (see, for example, [1, 6])

$$E_{\alpha, \beta}(z) = \sum_{k=0}^{\infty} \frac{z^k}{\Gamma(\alpha k + \beta)} \quad \text{for } \alpha, \beta, z \in \mathbb{R} \text{ with } \alpha > 0, \quad (2)$$

which is an entire function if, furthermore, $\beta > 0$.

The Laplace transform \mathcal{L} of the Riemann-Liouville fractional derivative is [6, (2.248)]

$$\mathcal{L}\{D_{RL}^\delta u\} = s^\delta \mathcal{L}\{u\} - C_1 - sC_2,$$

with $C_1 = [D_{RL}^{\delta-1}u](0)$ and $C_2 = [D_{RL}^{\delta-2}u](0)$. Thus, taking the Laplace transform of (1a), one obtains

$$\frac{f}{s} = -[s^\delta \mathcal{L}\{u\} - C_1 - sC_2] + b[s\mathcal{L}\{u\} - u(0)] = -[s^\delta \mathcal{L}\{u\} - C_1 - sC_2] + bs\mathcal{L}\{u\}$$

and therefore

$$\mathcal{L}\{u\} = \frac{C_1}{s(s^{\delta-1}-b)} + \frac{C_2}{s^{\delta-1}-b} - \frac{f}{s^2(s^{\delta-1}-b)}. \quad (3)$$

Now the Laplace transform of the Mittag-Leffler function is [6, (1.80)]

$$\mathcal{L}\left\{x^{\beta-1}E_{\alpha,\beta}(\pm\lambda x^\alpha)\right\} = \frac{s^{\alpha-\beta}}{s^\alpha \mp \lambda}.$$

Hence one can deduce from (3) that the solution of (1) is

$$u(x) = C_1 x^{\delta-1} E_{\delta-1,\delta}(bx^{\delta-1}) + C_2 x^{\delta-2} E_{\delta-1,\delta-1}(bx^{\delta-1}) - f x^\delta E_{\delta-1,\delta+1}(bx^{\delta-1}). \quad (4)$$

In this formula the constants C_1 and C_2 must be chosen to satisfy the boundary conditions (1b). The boundary condition $u(0) = 0$ forces $C_2 = 0$.

Remark 1. Recall that $1 < \delta < 2$. From (4) one sees that to obtain a solution u that lies in $C[0,1]$, the formulation of the problem (1) must include the homogenous Dirichet boundary condition $u(0) = 0$ in order to eliminate the singular component $x^{\delta-2} E_{\delta-1,\delta-1}(bx^{\delta-1})$.

The value of C_1 in (4) will be deduced from the boundary condition (1b) at $x = 1$. By [6, (1.82)] one has

$$D_{RL}^\gamma(x^{\beta-1}E_{\alpha,\beta}(\lambda x^\alpha)) = x^{\beta-\gamma-1}E_{\alpha,\beta-\gamma}(\lambda x^\alpha), \quad (5)$$

for constant α, β, γ and λ . When $\gamma = 1$ one has $D_{RL}^\gamma = d/dx$ [1, p.27]; hence (4) yields

$$u'(x) = C_1 x^{\delta-2} E_{\delta-1,\delta-1}(bx^{\delta-1}) - f x^{\delta-1} E_{\delta-1,\delta}(bx^{\delta-1}). \quad (6)$$

Thus, by (1b) one has

$$\begin{aligned} \gamma_1 &= u(1) + \alpha_1 u'(1) \\ &= C_1 [E_{\delta-1,\delta}(b) + \alpha_1 E_{\delta-1,\delta-1}(b)] - f [E_{\delta-1,\delta+1}(b) + \alpha_1 E_{\delta-1,\delta}(b)] \end{aligned}$$

and consequently

$$C_1 = \frac{\gamma_1 + f [E_{\delta-1,\delta+1}(b) + \alpha_1 E_{\delta-1,\delta}(b)]}{E_{\delta-1,\delta}(b) + \alpha_1 E_{\delta-1,\delta-1}(b)}. \quad (7)$$

Substituting (7) into (4) and (6) yields closed-form representations of the solution

$$\begin{aligned} u(x) &= \gamma_1 x^{\delta-1} \frac{E_{\delta-1,\delta}(bx^{\delta-1})}{E_{\delta-1,\delta}(b) + \alpha_1 E_{\delta-1,\delta-1}(b)} \\ &+ f \left[x^{\delta-1} \frac{E_{\delta-1,\delta+1}(b) + \alpha_1 E_{\delta-1,\delta}(b)}{E_{\delta-1,\delta}(b) + \alpha_1 E_{\delta-1,\delta-1}(b)} E_{\delta-1,\delta}(bx^{\delta-1}) - x^\delta E_{\delta-1,\delta+1}(bx^{\delta-1}) \right] \end{aligned} \quad (8)$$

and its first-order derivative

$$\begin{aligned}
u'(x) &= \gamma_1 x^{\delta-2} \frac{E_{\delta-1,\delta-1}(bx^{\delta-1})}{E_{\delta-1,\delta}(b) + \alpha_1 E_{\delta-1,\delta-1}(b)} \\
&+ f \left[x^{\delta-2} \frac{E_{\delta-1,\delta+1}(b) + \alpha_1 E_{\delta-1,\delta}(b)}{E_{\delta-1,\delta}(b) + \alpha_1 E_{\delta-1,\delta-1}(b)} E_{\delta-1,\delta-1}(bx^{\delta-1}) - x^{\delta-1} E_{\delta-1,\delta}(bx^{\delta-1}) \right].
\end{aligned} \tag{9}$$

Using the elementary identity

$$E_{\delta-1,i}(z) = z E_{\delta-1,\delta-1+i}(z) + \frac{1}{\Gamma(i)} \quad \text{for } i = 0, 1, 2 \tag{10}$$

in (8) and (9), we get

$$\begin{aligned}
u(x) &= \gamma_1 \frac{E_{\delta-1,1}(bx^{\delta-1}) - 1}{E_{\delta-1,1}(b) - 1 + \alpha_1 E_{\delta-1,0}(b)} \\
&+ \frac{f}{b} \left\{ x - \frac{E_{\delta-1,2}(b) - 1 + \alpha_1 [E_{\delta-1,1}(b) - 1]}{E_{\delta-1,1}(b) - 1 + \alpha_1 E_{\delta-1,0}(b)} \right\} \\
&+ \frac{f}{b} \left\{ \frac{E_{\delta-1,2}(b) - 1 + \alpha_1 [E_{\delta-1,1}(b) - 1]}{E_{\delta-1,1}(b) - 1 + \alpha_1 E_{\delta-1,0}(b)} E_{\delta-1,1}(bx^{\delta-1}) - x E_{\delta-1,2}(bx^{\delta-1}) \right\}
\end{aligned} \tag{11}$$

and

$$\begin{aligned}
u'(x) &= \gamma_1 x^{-1} \frac{E_{\delta-1,0}(bx^{\delta-1})}{E_{\delta-1,1}(b) - 1 + \alpha_1 E_{\delta-1,0}(b)} + \frac{f}{b} \\
&+ \frac{f}{b} x^{-1} \left\{ \frac{E_{\delta-1,2}(b) - 1 + \alpha_1 [E_{\delta-1,1}(b) - 1]}{E_{\delta-1,1}(b) - 1 + \alpha_1 E_{\delta-1,0}(b)} E_{\delta-1,0}(bx^{\delta-1}) - x E_{\delta-1,1}(bx^{\delta-1}) \right\}.
\end{aligned} \tag{12}$$

Lemma 1. For $j = 1, 2$ the function

$$\phi(x) = x^{\delta-j} E_{\delta-1,\delta-j+1}(bx^{\delta-1}), \quad \text{with } x > 0, \tag{13}$$

is a solution of

$$-D_{RL}^{\delta} \phi + b \phi' = 0.$$

Proof. It follows from [6, (1.82)] that

$$\begin{aligned}
-D_{RL}^{\delta}\phi + b\phi' &= -x^{-j}E_{\delta-1,-j+1}(bx^{\delta-1}) + bx^{\delta-j-1}E_{\delta-1,\delta-j}(bx^{\delta-1}) \\
&= x^{-j} \left[-\sum_{k=0}^{\infty} \frac{(bx^{\delta-1})^k}{\Gamma((\delta-1)k-j+1)} + \sum_{k=0}^{\infty} \frac{(bx^{\delta-1})^{k+1}}{\Gamma((\delta-1)k+\delta-j)} \right] \\
&= x^{-j} \left[-\sum_{k=1}^{\infty} \frac{(bx^{\delta-1})^k}{\Gamma((\delta-1)k-j+1)} + \sum_{k=1}^{\infty} \frac{(bx^{\delta-1})^k}{\Gamma((\delta-1)k-j+1)} \right] \\
&= 0,
\end{aligned}$$

where in the first series we have used $\Gamma(-1) = \Gamma(0) = \infty$.

Lemma 2. *The function*

$$\psi(x) = x^{\delta}E_{\delta-1,\delta+1}(bx^{\delta-1}), \quad \text{with } x > 0, \quad (14)$$

is a solution of

$$-D_{RL}^{\delta}\psi + b\psi' = -1.$$

Proof. It follows from [6, (1.82)] that

$$\begin{aligned}
-D_{RL}^{\delta}\psi + b\psi' &= -E_{\delta-1,1}(bx^{\delta-1}) + bx^{\delta-1}E_{\delta-1,\delta}(bx^{\delta-1}) \\
&= -\sum_{k=0}^{\infty} \frac{(bx^{\delta-1})^k}{\Gamma((\delta-1)k+1)} + \sum_{k=0}^{\infty} \frac{(bx^{\delta-1})^{k+1}}{\Gamma((\delta-1)k+\delta)} \\
&= -\sum_{k=0}^{\infty} \frac{(bx^{\delta-1})^k}{\Gamma((\delta-1)k+1)} + \sum_{k=1}^{\infty} \frac{(bx^{\delta-1})^k}{\Gamma((\delta-1)k+1)} \\
&= -1.
\end{aligned}$$

Observe that the functions $\phi(x)$ and $\psi(x)$ in Lemmas 13 and 14 are infinitely differentiable for $x > 0$.

Using Lemmas 1 and 2 it is straightforward to verify that the function u defined in (8) satisfies (1). In addition, from (9) we have $|u'(x)| \rightarrow \infty$ as $x \rightarrow 0^+$ for each fixed value of $\delta \in (1, 2)$.

3 Boundary layers in the solution

We now discuss the behaviour of $\|u\|_{\infty}$ and $u'(1)$ when $\delta \rightarrow 1^+$.

Note immediately that (9) and the hypothesis $1 < \delta < 2$ imply that $u'(x)$ blows up as $x \rightarrow 0^+$. This is a singularity in u , not a boundary layer (in the typical usage of this terminology in singularly perturbed differential equations), and we do not discuss it further.

Thus we investigate the other endpoint $x = 1$. This will involve different cases depending on the value of the convective term b ; cf. [8].

Let $\beta > \alpha > 0$ and $y \in \mathbb{R}$. We begin with the useful Mittag-Leffler identity

$$E_{\alpha,\beta}(y) = \frac{1}{\alpha\Gamma(\beta-\alpha)} \int_{t=0}^1 (1-t^{1/\alpha})^{\beta-\alpha-1} E_{\alpha,\alpha}(ty) dt \quad (15)$$

of [5, Lemma 2], which is easily proved by expanding $E_{\alpha,\alpha}(ty)$ as an infinite series in powers of ty and then integrating term by term. In [5] this identity is used to prove that $E_{\alpha,\beta}(-y)$ is completely monotonic for $0 < \alpha \leq 1$, $\beta \geq \alpha$ and $y \geq 0$. Hence in particular

$$E_{\alpha,\beta}(y) \geq 0 \text{ and } (d/dy)E_{\alpha,\beta}(y) \geq 0 \text{ for } 0 < \alpha \leq 1, \beta \geq \alpha, y \leq 0. \quad (16)$$

Of course one has trivially $E_{\alpha,\beta}(y) > 0$ for $y \geq 0$ (and any $\alpha \geq 0$, $\beta > 0$) from the definition (2). One can sharpen (16) to

$$E_{\alpha,\beta}(y) > 0 \text{ for } 0 < \alpha \leq 1, \beta > \alpha, y \leq 0 \quad (17)$$

because in (15) the integrand is continuous and non-negative with $E_{\alpha,\alpha}(0) = 1/\Gamma(\alpha) > 0$.

Furthermore, the identity (15) and the properties $E_{\alpha,\alpha}(0) = 1/\Gamma(\alpha)$ and $E_{\alpha,\alpha}(s) \geq 0$ for all $s \in \mathbb{R}$ imply that for $i = 0, 1$ one has

$$0 < E_{\delta-1,\delta+1+i}(y) < E_{\delta-1,\delta+i}(y) \text{ for all } y \in \mathbb{R}. \quad (18)$$

Thus for the quotients appearing in (8) and (9) it follows that

$$0 < \frac{E_{\delta-1,\delta+1}(b) + \alpha_1 E_{\delta-1,\delta}(b)}{E_{\delta-1,\delta}(b) + \alpha_1 E_{\delta-1,\delta-1}(b)} \leq \frac{E_{\delta-1,\delta+1}(b)}{E_{\delta-1,\delta}(b)} + \alpha_1 < 1 + \alpha_1. \quad (19)$$

3.1 Case $b \leq 0$

In this subsection assume that $b \leq 0$. By (16) and (18), for $0 \leq x \leq 1$ and $i = 0, 1$ one has

$$0 < E_{\delta-1,\delta+i}(bx^{\delta-1}) \leq E_{\delta-1,\delta+i}(0) \leq 1/\theta,$$

where $\theta := \min\{\Gamma(x) : 1 \leq x \leq 2\} \approx 0.8856$. Invoking this inequality and (19) in (8) yields $\|u\|_\infty \leq C$ (for some constant C) for $1 < \delta < 2$.

Similarly, (16) implies that $E_{\delta-1,\delta-1}(bx^{\delta-1}) \leq E_{\delta-1,\delta-1}(0) \leq 1$; combining this inequality and (19) with (9) yields $|u'(1)| \leq C$ (for some constant C) for $1 < \delta < 2$, so there is no boundary layer at $x = 1$ as $\delta \rightarrow 1^+$ when $b \leq 0$.

3.2 Case $0 < b < 1$

In this subsection assume that $0 < b < 1$.

The definition (2) yields $0 \leq E_{\delta-1,i}(bx^{\delta-1}) \leq E_{\delta-1,i}(b)$ for $i = 1, 2$ and $0 \leq x \leq 1$. The analysis in [8, Subsection 2.2.3] shows that

$$\frac{1}{4(1-b)} \leq E_{\delta-1,2}(b) \leq \frac{1}{1-b}, \quad (20)$$

$$\frac{1-b^{1+[1/(\delta-1)]}}{1-b} \leq E_{\delta-1,1}(b) \leq \frac{1}{\theta(1-b)}, \quad (21)$$

where $\theta \approx 0.8856$ was defined earlier and $[n]$ denotes the greatest integer satisfying $[n] \leq n$. Similarly one has

$$0 \leq E_{\delta-1,0}(bx^{\delta-1}) \leq E_{\delta-1,0}(b) = \sum_{k=1}^{\infty} \frac{b^k}{\Gamma(k(\delta-1))} \leq \frac{1}{\theta} \sum_{k=1}^{\infty} b^k = \frac{b}{\theta(1-b)}. \quad (22)$$

It follows from (11) and (20)–(22) that

$$\|u\|_{\infty} \leq C$$

for some constant C whose value depends on b but is independent of δ .

By (12) and (20)–(22) we get $|u'(x)| \leq C$ (where C depends on b but not on δ) for $x > c > 0$ where $c \in (0, 1)$ is any fixed constant. It follows that u does not have a boundary layer at $x = 1$ as $\delta \rightarrow 1^+$ when $0 < b < 1$.

3.3 Case $b = 1$

In this subsection assume that $b = 1$.

For any constant $r \geq 0$ we have

$$\begin{aligned} \int_{x=r}^{\infty} \frac{dx}{\Gamma(x)} &= \int_{x=0}^{\infty} \frac{dx}{\Gamma(x+r)} = \sum_{k=0}^{\infty} \int_{x=k(\delta-1)}^{(k+1)(\delta-1)} \frac{dx}{\Gamma(x+r)} \\ &= \lim_{\delta \rightarrow 1^+} \sum_{k=0}^{\infty} \frac{\delta-1}{\Gamma(k(\delta-1)+r)} \\ &= \lim_{\delta \rightarrow 1^+} (\delta-1)E_{\delta-1,r}(1), \end{aligned}$$

where the penultimate equality holds true by the theory of Riemann sums in integration. Now Table VI of [2] gives the numerical values

$$\int_{x=0}^{\infty} \frac{dx}{\Gamma(x)} \approx 2.808, \quad \int_{x=0}^1 \frac{dx}{\Gamma(x)} \approx 0.541, \quad \int_{x=1}^2 \frac{dx}{\Gamma(x)} \approx 1.085,$$

so

$$\int_{x=1}^{\infty} \frac{dx}{\Gamma(x)} \approx 2.267, \quad \int_{x=2}^{\infty} \frac{dx}{\Gamma(x)} \approx 1.182.$$

Thus

$$\lim_{\delta \rightarrow 1^+} (\delta - 1)E_{\delta-1,i}(1) \approx \begin{cases} 2.808 & \text{if } i = 0, \\ 2.267 & \text{if } i = 1, \\ 1.181 & \text{if } i = 2. \end{cases} \quad (23)$$

We first deduce bounds for $\|u\|_\infty$. Observe that the first two terms in (11), viz.,

$$\gamma_1 \frac{E_{\delta-1,1}(bx^{\delta-1}) - 1}{E_{\delta-1,1}(b) - 1 + \alpha_1 E_{\delta-1,0}(b)}, \quad \frac{f}{b} \left\{ x - \frac{E_{\delta-1,2}(b) - 1 + \alpha_1 [E_{\delta-1,1}(b) - 1]}{E_{\delta-1,1}(b) - 1 + \alpha_1 E_{\delta-1,0}(b)} \right\}$$

are bounded by some constant C , so we analyse the third term $(f/b)\{\dots\}$ with $b = 1$. Invoking (23) we obtain

$$\lim_{\delta \rightarrow 1^+} \frac{E_{\delta-1,2}(1) - 1 + \alpha_1 [E_{\delta-1,1}(1) - 1]}{E_{\delta-1,1}(1) - 1 + \alpha_1 E_{\delta-1,0}(1)} \approx \frac{1.181 + 2.267\alpha_1}{2.267 + 2.808\alpha_1} \geq \frac{1.181}{2.808} > 0.42. \quad (24)$$

If δ is sufficiently close to 1 and $0 < x < 0.42$, by (24) and the trivial inequality $E_{\delta-1,1}(x^{\delta-1}) \geq E_{\delta-1,2}(x^{\delta-1})$ we have

$$\begin{aligned} & \frac{E_{\delta-1,2}(1) - 1 + \alpha_1 [E_{\delta-1,1}(1) - 1]}{E_{\delta-1,1}(1) - 1 + \alpha_1 E_{\delta-1,0}(1)} E_{\delta-1,1}(x^{\delta-1}) - x E_{\delta-1,2}(x^{\delta-1}) \\ & \geq (0.42 - x) E_{\delta-1,2}(x^{\delta-1}) \\ & \geq (0.42 - x) \sum_{k=0}^{\lfloor 1/(\delta-1) \rfloor} \frac{(x^{\delta-1})^k}{\Gamma(3)} \\ & = (0.42 - x) \frac{1 - (x^{\delta-1})^{1+\lfloor 1/(\delta-1) \rfloor}}{2(1 - x^{\delta-1})} \\ & > (0.42 - x) \frac{1 - x}{2(1 - x^{\delta-1})} \end{aligned}$$

because $(x^{\delta-1})^{1+\lfloor 1/(\delta-1) \rfloor} < (x^{\delta-1})^{1/(\delta-1)} = x$. But $(1-x)/(1-x^{\delta-1}) \rightarrow \infty$ as $\delta \rightarrow 1^+$ since $x > 0$. Consequently $\lim_{\delta \rightarrow 1^+} \|u\|_\infty = \infty$.

We now show that $u'(1)$ blows up as $\delta \rightarrow 1^+$. Set $x = 1$ in (12), multiply by $\delta - 1$ then take the limit as $\delta \rightarrow 1^+$, and appeal to (23): this yields

$$\lim_{\delta \rightarrow 1^+} (\delta - 1)u'(1) \approx \left[\frac{2.808(1.181 + 2.267\alpha_1)}{2.267 + 2.808\alpha_1} - 2.267 \right] f.$$

That is, $|u'(1)| = O(1/(\delta - 1))$ as $\delta \rightarrow 1^+$. Thus, the derivative of u at $x = 1$ blows up as δ tends to 1^+ when $b = 1$ and $f \neq 0$.

Figure 1 displays the exact solution for two values of δ when $b = 1$ and δ equals 1.01 and 1.0001. Note that the scales on the vertical axes in the two plots are different and a typical boundary layer is not observed at $x = 1$ although $u'(1)$ is large. In [7] a related problem (where the Riemann-Liouville derivative is replaced by a Caputo derivative) is analysed in detail.




Fig. 1 Exact solution of (1) for $b = 1, f = 1, \alpha_1 = 0, \gamma_1 = 0$ and $\delta = 1.01$ (left figure) and $\delta = 1.0001$ (right figure).

3.4 Case $b > 1$

In this subsection assume that $b > 1$. We begin with a technical lemma. Recall the asymptotic relation

$$E_{\delta-1,n}(b) = \frac{1}{\delta-1} b^{(1-n)/(\delta-1)} \exp(b^{1/(\delta-1)}) + O\left(\frac{1}{(\delta-1)^2}\right) \quad \text{as } \delta \rightarrow 1^+ \quad (25)$$

of [8, (2.19)]; in this formula the index n must be fixed independently of δ .

Lemma 3.

$$\begin{aligned} & E_{\delta-1,\delta+1}(b)E_{\delta-1,\delta-1}(b) - [E_{\delta-1,\delta}(b)]^2 \\ &= O\left(\frac{1}{(\delta-1)^3} b^{1/(\delta-1)} \exp(b^{1/(\delta-1)})\right) \quad \text{as } \delta \rightarrow 1^+. \end{aligned} \quad (26)$$

Proof. By (10) we have

$$\begin{aligned} & E_{\delta-1,\delta+1}(b)E_{\delta-1,\delta-1}(b) - [E_{\delta-1,\delta}(b)]^2 \\ &= \frac{1}{b^2} [E_{\delta-1,2}(b) - 1] E_{\delta-1,0}(b) - \frac{1}{b^2} [E_{\delta-1,1}(b) - 1]^2 \\ &= \frac{1}{b^2} \left\{ E_{\delta-1,2}(b)E_{\delta-1,0}(b) - [E_{\delta-1,1}(b)]^2 - E_{\delta-1,0}(b) + 2E_{\delta-1,1}(b) - 1 \right\} \\ &= O\left(\frac{1}{(\delta-1)^3} b^{1/(\delta-1)} \exp(b^{1/(\delta-1)})\right) \quad \text{as } \delta \rightarrow 1^+, \end{aligned} \quad (27)$$

on invoking (25), because the highest-order terms cancel and the expression in (27) is then the dominant term among those remaining.

We use Lemma 3 to analyse the behaviour of u near $x = 1$. From (9) one has

$$\begin{aligned}
u'(1) &= \gamma_1 \frac{E_{\delta-1,\delta-1}(b)}{E_{\delta-1,\delta}(b) + \alpha_1 E_{\delta-1,\delta-1}(b)} \\
&\quad + f \left[\frac{E_{\delta-1,\delta+1}(b) + \alpha_1 E_{\delta-1,\delta}(b)}{E_{\delta-1,\delta}(b) + \alpha_1 E_{\delta-1,\delta-1}(b)} E_{\delta-1,\delta-1}(b) - E_{\delta-1,\delta}(b) \right] \\
&= \frac{1}{E_{\delta-1,\delta}(b) + \alpha_1 E_{\delta-1,\delta-1}(b)} \times \\
&\quad \left\{ \gamma_1 E_{\delta-1,\delta-1}(b) + f E_{\delta-1,\delta+1}(b) E_{\delta-1,\delta-1}(b) - f [E_{\delta-1,\delta}(b)]^2 \right\}.
\end{aligned}$$

Here

$$\left\{ \dots \right\} = O\left(\frac{1}{(\delta-1)^3} b^{1/(\delta-1)} \exp(b^{1/(\delta-1)})\right)$$

by Lemma 3 and (25). From (25) we also get


$$E_{\delta-1,\delta}(b) + \alpha_1 E_{\delta-1,\delta-1}(b) = \frac{1 + \alpha_1 b^{1/(\delta-1)}}{b(\delta-1)} \exp(b^{1/(\delta-1)}) + O\left(\frac{1}{(\delta-1)^2}\right).$$

Consequently

$$|u'(1)| = O\left(\frac{b^{1/(\delta-1)}}{(\delta-1)^2} \left(1 + \alpha_1 b^{-1/(\delta-1)}\right)\right) \quad \text{as } \delta \rightarrow 1^+. \quad (28)$$

Thus $u'(1)$ blows up when $\delta \rightarrow 1^+$ and $b > 1$, and this behaviour is more extreme if $\alpha_1 = 0$, i.e., if there is a Dirichlet boundary condition at the endpoint $x = 1$. Figure 2 indicates that a layer appears at $x = 1$ when δ is near 1. One can prove analytically that this layer is present, but this derivation is too long to include here.

Fig. 2 Exact solution of (1) for $b = 1.1$, $f = 1$, $\alpha_1 = 0$, $\gamma_1 = 1.7$ and $\delta = 1.03$.



In closing, we mention that when $b > 1$, $\|u\|_\infty$ is unbounded as $\delta \rightarrow 1^+$. The analysis needed to show this resembles the analysis given above for $u'(1)$.

Acknowledgements This paper was written while the second author was visiting the University of Zaragoza, supported by the Institute of Mathematics and Applications (IUMA). The re-

search was also partly supported by the projects MTM2013-40842-P, UZCUD2014-CIE-09 and the Diputación General de Aragón.

References

1. Kai Diethelm. *The analysis of fractional differential equations*, volume 2004 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2010. An application-oriented exposition using differential operators of Caputo type.
2. Arne Fransén and Staffan Wrigge. High-precision values of the gamma function and of some related coefficients. *Math. Comp.*, 34(150):553–566, 1980.
3. Bangti Jin, Raytcho Lazarov, Joseph Pasciak, and William Rundell. Variational formulation of problems involving fractional order differential operators. *Math. Comp.* (To appear).
4. Natalia Kopteva and Martin Stynes. Analysis and numerical solution of a Riemann-Liouville fractional derivative two-point boundary value problem. (Submitted for publication).
5. Kenneth S. Miller and Stefan G. Samko. A note on the complete monotonicity of the generalized Mittag-Leffler function. *Real Anal. Exchange*, 23(2):753–755, 1997/98.
6. Igor Podlubny. *Fractional differential equations*, volume 198 of *Mathematics in Science and Engineering*. Academic Press Inc., San Diego, CA, 1999. An introduction to fractional derivatives, fractional differential equations, to methods of their solution and some of their applications.
7. Martin Stynes and José Luis Gracia. Blow-up of solutions and interior layers in a Caputo two-point boundary value problem. (Submitted to BAIL 2014 proceedings).
8. Martin Stynes and José Luis Gracia. Boundary layers in a two-point boundary value problem with a Caputo fractional derivative. *Comput. Methods Appl. Math.*, (15):79–95, 2015.