

Recent developments of universal modules

H. Melis Tekin Akcin

September 4-8, 2017

Contact: hmtekin@hacettepe.edu.tr, Hacettepe University

Abstract: For a commutative algebra R , its de Rham cohomology is an important invariant of R . In this talk, an infinite chain of de Rham-like complexes is introduced where the first member of the chain is the de Rham complex. The complexes are called *approximations of the de Rham complex*. Their cohomologies are found for polynomial rings.

References

- [1] Brown, W. C. The algebra of differentials of infinite rank. *Can. J. Math.* **XXV** (1) (1973), 141–155.
- [2] Erdogan, A. Homological dimensions of the universal modules for hypersurfaces, *Comm. Algebra*, **24** (5) (1996) 1565–1573.
- [3] Hart, R. Glued algebras and differential operators, *Bull. LMS*, **23** (1991) 351–355.
- [4] Hart, R. Higher derivations and universal differential operators, *J. Algebra*, **184** (1996) 175–181.
- [5] Heyneman R. G. and Sweedler, M. E. Affine Hopf algebras I, *J. Algebra*, **13** (1969) 192–241.
- [6] Nakai, Y. High order derivations 1, *Osaka Journal of Mathematics*, **7** (1970) 1–27.
- [7] Osborn, H. Modules of differentials 1, *Mathematische Annalen*, **170** (1967) 221–244.