

# ERROR ESTIMATES FOR HIGHER-ORDER FINITE VOLUME SCHEMES FOR CONVECTION DIFFUSION PROBLEMS

DIETMAR KRÖNER\* AND MIRKO ROKYTA†

**Abstract.** It is still an open problem to prove a priori error estimates for finite volume schemes of higher order MUSCL type, including limiters, on unstructured meshes, which show some improvement compared to first order schemes. In this paper we use these higher order schemes for the discretization of convection dominated elliptic problems in a convex bounded domain  $\Omega$  in  $\mathbb{R}^2$  and we can prove such kind of an a priori error estimate. In the part of the estimate, which refers to the discretization of the convective term, we gain  $h^{1/2}$ . Although the original problem is linear, the numerical problem becomes nonlinear, due to MUSCL type reconstruction/limiter technique.

**Key words.** linear convection dominated diffusion equation in 2D, upwind finite volume scheme, first and higher order finite volume schemes, a priori error estimates, MUSCL type reconstruction/limiter.

**AMS subject classifications.** 65N15, 35J25, 76M25

**1. Introduction.** There are many Finite Volume and Discontinuous Galerkin schemes for solving elliptic convection dominated problems and nonlinear conservation laws on unstructured grids in multi dimensions, as

$$\begin{aligned} \partial_t v + \operatorname{div} f(v) &= 0 && \text{in } \mathbb{R}^n \times \mathbb{R}^+ \\ v(x, 0) &= u_0(x) && \text{on } \mathbb{R}^n. \end{aligned} \tag{1.1}$$

While for strongly elliptic problems like

$$\begin{aligned} -\Delta v &= f && \text{in } \Omega \\ v(x) &= 0 && \text{on } \partial\Omega \end{aligned} \tag{1.2}$$

with dominating diffusion, no stabilization is necessary for numerical schemes, we need some upwinding [21] or, for higher order schemes, a suitable stabilization [22], [32], [45] for convection dominated problems like

$$\begin{aligned} -\varepsilon \Delta v + \operatorname{div}(bv) + cv &= f && \text{in } \Omega \\ v(x) &= 0 && \text{on } \partial\Omega. \end{aligned} \tag{1.3}$$

The same statement holds also for nonlinear conservation laws as in (1.1). In this case the stabilization is obtained by reconstruction technique with so called limiters. They make the scheme nonlinear, even in cases where the underlying partial differential equation (1.1) is linear. For finite volume schemes, the reconstruction with limiters can be realized in a very easy way even on unstructured grids, e.g. by MUSCL type discretizations. However, the theoretical background for these schemes, in particular when applied to conservation laws, is not yet satisfactorily developed. Concerning the convergence of both first and higher order schemes, there are results in the case of nonlinear scalar conservation laws [13], [17], [34], [33], [15], and in the case of weakly coupled systems of conservation laws [40]. For conservation laws as in (1.1), a priori error estimates of the form

$$\|v - u_h\|_{L^\infty(L^1)} \leq ch^{\frac{1}{4}} + \text{approximation error of data} \tag{1.4}$$

are available [3], [12], [42], [13], [2], [7]. Here,  $v$  denotes the exact solution of the underlying partial differential equation and  $u_h$  the approximative numerical solution obtained by a first order finite volume scheme in multi dimensions on unstructured grids. From numerical experiments one would expect  $h^{\frac{1}{2}}$  in (1.4), but the proof for this on unstructured grids is an open question.

---

\*Universität Freiburg, Institut für Angewandte Mathematik, Hermann–Herder–Str. 10, 79104 Freiburg, Germany (dietmar@mathematik.uni-freiburg.de).

†Charles University, Department of Mathematical Analysis, Sokolovská 83, 186 75 Praha, Czech Republic (rokyta@karlin.mff.cuni.cz). M. Rokyta was partially supported by Prvoug P47.

For smooth solutions of the linear transport equations

$$\partial_t v + (a \cdot \nabla)v = 0 \quad \text{in } \mathbb{R}^n \times \mathbb{R}^+ \quad (1.5)$$

one gets [2]

$$\|v - u_h\|_p \leq ch. \quad (1.6)$$

There are also no error estimates of the type  $\|v - u_h\| \leq ch^\beta$  for higher order finite volume schemes for conservation laws in multi dimensions on unstructured grids including limiters, with  $\beta > \frac{1}{4}$ . To get results in this direction, concerning nonlinear hyperbolic conservation laws, seems to be very difficult. Theoretically justified error analysis for upwind finite volume schemes of higher order, which would also indicate the higher order convergence rate, remains an open problem. See for example [15], [42], [13], [12], [14].

Therefore in this paper we apply the higher order finite volume schemes with limiters to a linear convection dominated stationary diffusion equations like (1.3) in multi dimensions on partially unstructured, locally irregular grids (see Assumption 2.2 and Remark 2.3 (c), as well as left part of the Fig. 2.1). In this case the numerical scheme becomes nonlinear, because of the limiter. We can show that we gain  $h^{\frac{1}{2}}$  in the error estimate for the term, which refers to the discretization of the convective term (i.e.  $\frac{h^3}{\varepsilon}$ ), compared to first order schemes. For the higher order scheme we get (see Theorem 4.1)

$$\|z_h\|_\varepsilon^2 \leq c\left(\varepsilon h^2 + h^{4-2\delta} + \frac{h^3}{\varepsilon}\right)\|v\|_{2,2}^2 + c\frac{h^4}{\varepsilon} \sum_j R_j^2 |T_j| \quad (1.7)$$

and if  $v \in W^{3,2}(\Omega)$ , in particular for smooth solutions, we have

$$\|z_h\|_\varepsilon^2 \leq c\left(\varepsilon h^3 + h^3 + \frac{h^3}{\varepsilon}\right)\|v\|_{3,2}^2 + c\frac{h^4}{\varepsilon} \sum_j R_j^2 |T_j|.$$

Later we will show that the term  $\sum_j R_j^2 |T_j|$  (see (4.8)) is of the same order with respect to  $\varepsilon$  as  $\|v\|_{2,2}^2$ . In the case of the first order scheme we get (for comparison) the following result

$$\|z_h\|_\varepsilon^2 \leq c\left(\varepsilon h^2 + h^3 + \frac{h^2}{\varepsilon}\right)\|v\|_{2,2}^2. \quad (1.8)$$

Let us briefly mention some related results. In [28], [23] a convection diffusion equation like (1.3) with  $\varepsilon = 1$  and general elliptic part is considered. They prove error estimates of the form

$$\|v - u_h\|_{L^2} \leq ch$$

for finite volume schemes of first order. Further results for elliptic and parabolic equations for finite volume schemes are obtained in [4], [24] and the results of an interesting benchmark problem are published in [25]. Lube considered in [36] discretizations of (1.3) but with  $B(v)\nabla v$  instead of  $\text{div}(bv)$  and proved

$$\|v - u_h\| \leq c_\varepsilon h^k (\varepsilon^{\frac{1}{2}} + h^{\frac{1}{2}})$$

for the streamline diffusion method. Here  $k$  is the degree of the local polynomials and

$$\|v\|^2 = \varepsilon |v|_{1,2}^2 + \sum_i \delta_i \|B(u_h)\nabla v\|_{0,2,T_i}^2.$$

A-priori error estimates of the type (1.4), e.g. with the  $\varepsilon \|\cdot\|_{H^{1,2}} + \|\cdot\|_{L^2}$ -norm, are also known for the streamline diffusion shock capturing method applied to the linear transport equation with  $h^{\frac{3}{2}}$ , see [31].

For dominating diffusion problems there are error estimates for first order schemes (cf. [28], [23] for stationary case), which show  $\|v - u_h\|_{L^2} \leq ch$ . In [26], [37] convergence for a first order combined finite volume–finite element method in the non-stationary case was proved.

For second order TVD Rung-Kutta Discontinuous Galerkin methods with piecewise polynomials of order  $k$  in space a priori error estimates of the form

$$\|u - u_h\|_{L^2} \leq ch^{k+\frac{1}{2}}$$

for smooth solutions  $u$  of (1.1) have been proved in [47]. More advanced results for (also hybridized) Discontinuous Galerkin methods can be found e.g. in [43], [16], [39]. For finite element approximations of convection diffusion problems we refer to [5], [41] and for systems to [29].

The following papers deal with finite volume schemes for elliptic problems which are not convection dominated. In [9], [10] and [8] general frameworks for the construction and analysis of higher-order finite volume methods are developed and optimal error estimates are derived. In particular this has also been done for higher order finite volume schemes on rectangular partitions in [6]. Multiscale discontinuous Galerkin schemes for second order elliptic equations with rough coefficients are considered in [44]. Detailed error estimates are derived and confirmed by numerical experiments. Limiters are not included.

Finite volume schemes for elliptic problems without any error estimate are considered in the following papers. Development of higher order schemes are studied in [30]. Oevermann and Klein consider in [38] finite volume schemes of second order for two-dimensional elliptic equations with variable, discontinuous coefficients. Different limiting procedure for diffusion problems are discussed in [35]. In [18] a review of some of the most successful higher-order numerical schemes for the compressible Navier-Stokes equations on unstructured grids are presented. They use Moving-Least-Squares approximations for the construction of higher order finite volume schemes on unstructured grids. They can be used for direct reconstruction of the fluxes at cell edges for hyperbolic and elliptic problems.

**2. The problem.** Consider the following boundary value problem

$$Lv := -\varepsilon \Delta v + \operatorname{div}(bv) + cv = f \quad \text{in } \Omega, \quad (2.1)$$

$$v = 0 \quad \text{on } \partial\Omega \quad (2.2)$$

where  $\Omega$  is a convex polygonal domain in  $\mathbb{R}^2$  and  $b(x)$ ,  $c(x)$ ,  $f(x)$  are functions which are sufficiently smooth on  $\bar{\Omega}$  and such that  $0 < c_0 \leq c(x) \leq c_1$ ,  $\operatorname{div} b = 0$  in  $\bar{\Omega}$ . Moreover we suppose that the diffusion parameter  $\varepsilon$  is a positive constant,  $0 < \varepsilon \leq 1$ .

We consider that  $\bar{\Omega} = \bigcup_j \bar{T}_j$ , where  $T_j \in \mathcal{T}_h$  are open triangles,  $h := \sup_j \operatorname{diam}(T_j)$ ,  $0 < h < h_0$ . Furthermore, all boundary triangles are mirrored by the boundary of  $\Omega$  to get a corresponding ghost triangles (see Fig. 2.1 on the right). The set of all ghost triangles will be denoted by  $\mathcal{T}_G$ ,  $\mathcal{T}_G \cap \mathcal{T}_h = \emptyset$ .

NOTATION 2.1. We denote by

- (i)  $|T_j|$ : the volume of triangle  $T_j$ ;  $T_j \in \mathcal{T}_h \cup \mathcal{T}_G$
- (ii)  $x_j$ : the center of gravity of  $T_j$  (i.e.,  $x_j$  is the center of the inscribed circle to the triangle  $T_j$ )
- (iii)  $\bar{x}_j$ : intersection of the perpendicular bisectors of  $T_j$  (i.e.,  $\bar{x}_j$  is the center of the circumscribed circle to the triangle  $T_j$ )
- (iv)  $v_j := v(\bar{x}_j)$
- (v)  $N_j$ : the set of the numbers of the neighboring triangles to  $T_j$ ,  $T_j \in \mathcal{T}_h$
- (vi)  $T_{j\ell} := T_j \cup T_\ell$
- (vii)  $S_{j\ell}$ ,  $\ell \in N_j$ : the joint edge of  $T_j$  and  $T_\ell$  with length  $|S_{j\ell}|$ , where  $T_j \in \mathcal{T}_h$ ,  $T_\ell \in \mathcal{T}_h \cup \mathcal{T}_G$
- (viii)  $x_{j\ell}$ : the midpoint of  $S_{j\ell}$
- (ix)  $d_{j\ell} := |x_\ell - x_j|$
- (x)  $\bar{d}_{j\ell} := |\bar{x}_\ell - \bar{x}_j|$

(xi)  $\gamma_{j\ell} := \frac{|S_{j\ell}|}{d_{j\ell}}$ ;  $\gamma := \min \gamma_{j\ell}$ ;

(xii)  $n_{j\ell}$ : the outward unit normal to  $T_j \in \mathcal{T}_h$  in the direction of  $T_\ell$ ,  $\ell \in N_j$ .

We assume that there exists an  $\eta > 0$  such that all angles of all triangles  $T_j \in \mathcal{T}_h$  are less than  $\frac{\pi}{2} - \eta$ . Therefore, both  $x_j$  and  $\bar{x}_j$  lie strictly inside of  $T_j$  for all  $j$  and there is a constant  $c_\eta > 0$  independent of  $h$  such that  $\gamma > c_\eta$ .

Moreover we assume that  $\mathcal{T}_h = \mathcal{T}_R \cup \mathcal{T}_S$ , such that  $\mathcal{T}_R \cap \mathcal{T}_S = \emptyset$ , where

$$\mathcal{T}_R = \{T_j \in \mathcal{T}_h; T_j \text{ is equilateral and } T_\ell \text{ is equilateral } \forall \ell \in N_j\}, \quad (2.3)$$

and

$$\mathcal{T}_S = \{T_j \in \mathcal{T}_h; T_j \text{ is not equilateral or } \exists \ell \in N_j \text{ s.t. } T_\ell \text{ is not equilateral}\}. \quad (2.4)$$

The triangles in  $\mathcal{T}_R$  and  $\mathcal{T}_S$  are called regular and singular, respectively.

**ASSUMPTION 2.2.** *We assume that the triangulation is locally irregular in the sense of Heinrich (cf. [27, par. 2.2.2, p. 27]), i.e. that the set  $\mathcal{T}_S$  consists of the finite number of strips of triangles, each being of the width of  $O(h)$ . See the left of Fig. 2.1 for the triangles of  $\mathcal{T}_R$  (white color) and the triangles of  $\mathcal{T}_G$  (darker color).<sup>1</sup>*

**REMARK 2.3.** (a) If we have  $c = 0$  everywhere then we have to change the norm in (4.6): the  $c_0$ -term has to be cancelled. In the estimates from below and above in Sections 7 and 8 the term  $\psi_{N_j}$  is equal to zero and will not appear. Then all the arguments can be repeated. If  $c = 0$  only in a subset of  $\Omega$  then there will be some problems. We can change the definition of the norm in (4.6) by  $\sum_{T_j} c_j |z_j|^2 |T_j|$  but then, e.g. the  $c_0$ -term in the last estimate in the proof of Lemma 7.1 (and correspondingly in the proof of Lemma 7.2) cannot be controlled by the left-hand side. Therefore degeneration of  $c$  in part of  $\Omega$  is not allowed in this framework.

(b) The condition  $\operatorname{div} b = 0$ , mentioned in the beginning of Section 2, ensures that (3.10) and (3.13) are valid, and that we can obtain the estimate (8.16). This means that a local conservation property holds. This is essential to get the final estimates. Therefore, also a condition of the type  $-\frac{1}{2} \operatorname{div} b + c \geq 0$  would not be sufficient.

(c) The analysis is performed essentially only on uniform grids consisting of identical equilateral triangles. The MUSCL procedure is used only on these elements and furthermore, the set of non-equilateral triangles is assumed to be very small (strips of width  $O(h)$ ). Even then the non-equilateral triangles are assumed to have all angles less than  $\frac{\pi}{2} - \eta$ . These conditions are rather restrictive concerning domains which can be triangulated by the locally irregular triangulation. It can be shown, however, that every quadrilateral and pentagon can be triangulated by a reasonable finite amount of acute triangles (see [46]), so we can handle the acute triangulation of local quadrilaterals and pentagons occurring in the finite number of strips of irregularity.

**ASSUMPTION 2.4.** *For the solution  $v$  of (2.1) and (2.2) we assume  $v \in W^{2,2}(\Omega)$  and that  $v$  can be extended onto a small strip  $\omega_d$  of the width of  $O(h)$  outside of  $\Omega$  such that we have  $v(\bar{x}_\ell) = -v(\bar{x}_j)$  if  $T_\ell \subset \omega_d$  is the mirrored ghost triangle to  $T_j$ . For the continuation  $v_d$  of  $v$  we assume*

$$\|v_d\|_{W^{2,2}(\Omega_d)} \leq c \|v\|_{W^{2,2}(\Omega)},$$

where the constant  $c$  is independent of  $v$  and  $\Omega_d := \Omega \cup \omega_d$ .

In the context of the locally irregular grid we will also use the following result (cf. [27, p. 189] and the references there):

**THEOREM 2.5.** *Let  $\Omega$  be a convex polygonal domain in  $\mathbb{R}^2$  and  $\omega_h \subset \Omega$  be the strip of the width of  $O(h)$ ,  $0 < h < h_0$ . Then there is a constant  $c > 0$  independent of  $h$  such that for all  $v \in W^{j+1,2}(\Omega)$ ,  $j = 0, 1, 2$ , we have*

$$\|v\|_{W^{j,2}(\omega_h)} \leq ch^{\frac{1}{2}} \|v\|_{W^{j+1,2}(\Omega)}, \quad j = 0, 1, 2. \quad (2.5)$$

<sup>1</sup>For drawing the pictures we are grateful to Mr. Johannes Schäfer.




FIG. 2.1. A locally irregular mesh and ghosts cells.

We split the set  $\mathcal{E}$  of the edges  $S_{j\ell} = \overline{T_j} \cap \overline{T_\ell}$ , with  $T_j, T_\ell \in \mathcal{T}_h$ , into three parts,  $\mathcal{E} = \mathcal{E}_R \cup \mathcal{E}_S \cup \mathcal{E}_M$ , where

$$\begin{aligned} \mathcal{E}_R &:= \{S_{j\ell}; S_{j\ell} \not\subset \partial\Omega; \text{ both } T_j \text{ and } T_\ell \text{ are regular}\}, \\ \mathcal{E}_S &:= \{S_{j\ell}; S_{j\ell} \not\subset \partial\Omega; \text{ both } T_j \text{ and } T_\ell \text{ are singular}\}, \\ \mathcal{E}_M &:= \{S_{j\ell}; S_{j\ell} \not\subset \partial\Omega; T_j \text{ and } T_\ell \text{ are of different type (regular, singular)}\}. \end{aligned} \quad (2.6)$$

and call them regular, singular and mixed edges, respectively. We also denote by

$$\mathcal{E}_B := \{S_{j\ell}; S_{j\ell} \subset \partial\Omega;\} \quad (2.7)$$

and call them the boundary edges.

Furthermore we denote

$$N_{jI} := \{\ell | T_\ell \text{ is neighboring triangle to } T_j, \text{ and } T_\ell \subset \Omega\},$$

and

$$N_{jR} := \{\ell | T_\ell \text{ is neighboring triangle to } T_j, \text{ and } T_\ell \in \mathcal{T}_R\},$$

$$N_{jS} := \{\ell | T_\ell \text{ is neighboring triangle to } T_j, \text{ and } T_\ell \in \mathcal{T}_S\}.$$

$$N_{jG} := \{\ell | T_\ell \text{ is neighboring triangle to } T_j, \text{ and } T_\ell \in \mathcal{T}_G\},$$

**3. The scheme.** Let  $c_h(x) := c_j$ ,  $f_h(x) := f_j$  for  $x \in T_j \in \mathcal{T}_h$ , be piecewise constant approximations of  $c$ ,  $f$ , respectively, defined by

$$c_j := \frac{1}{|T_j|} \int_{T_j} c, \quad f_j := \frac{1}{|T_j|} \int_{T_j} f. \quad (3.1)$$

Let  $u_h(x) = u_j$  for  $x \in T_j \in \mathcal{T}_h \cup \mathcal{T}_G$ , be a piecewise constant solution of the discrete problem

$$(L_h u_h)_j = f_j, \quad \text{if } T_j \in \mathcal{T}_h, \quad (3.2)$$

$$u_\ell = -u_j, \quad \text{if } S_{j\ell} \subset \partial\Omega, \text{ and } T_\ell \in \mathcal{T}_G \text{ is the ghost triangle to } T_j \subset \Omega, \quad (3.3)$$

where the discrete operator is given by

$$(L_h u_h)_j := -\frac{\varepsilon}{|T_j|} \sum_{\ell \in N_j} (u_\ell - u_j) \gamma_{j\ell} + \frac{1}{|T_j|} \sum_{\ell \in N_j} g_{j\ell}(\mathcal{U}_{j\ell}, \mathcal{U}_{\ell j}) + c_j u_j. \quad (3.4)$$

The first term in (3.4) approximates the value of the diffusion term  $-\varepsilon\Delta v$  in  $\bar{x}_j$ , while  $\sum_{\ell \in N_j} g_{j\ell}(\mathcal{U}_{j\ell}, \mathcal{U}_{\ell j})$  approximates the values of the convective term  $\operatorname{div}(bv)$  along  $S_{j\ell}$ . Here,  $g_{j\ell}$  stands for an upwind finite volume flux, and,

$$\mathcal{U}_{j\ell} = \mathcal{U}_{j\ell}(u_j, u_\ell), \quad \mathcal{U}_{\ell j} = \mathcal{U}_{\ell j}(u_\ell, u_j) \quad (3.5)$$

will be defined more precisely later. A particular scheme of the type (3.2)–(3.5) is then chosen by the particular choice of functions  $\mathcal{U}_{j\ell}$  and  $g_{j\ell}$ .

**EXAMPLE 3.1. (General numerical flux)** In general, we suppose that the upwind finite volume flux  $g_{j\ell}(u, v)$  is a *Lipschitz continuous* function, i.e., we suppose that there is a constant  $c > 0$  such that

$$|g_{j\ell}(u, v) - g_{j\ell}(u', v')| \leq ch (|u - u'| + |v - v'|). \quad (3.6)$$

Furthermore we suppose that  $g_{j\ell}$  satisfies the following three basic properties:

$$g_{j\ell}(u, u) = u \int_{S_{j\ell}} bn_{j\ell} ds, \quad (3.7)$$

$$g_{j\ell}(u, v) = -g_{\ell j}(v, u), \quad (3.8)$$

$$\frac{\partial}{\partial u} g_{j\ell}(u, v) \geq 0 \geq \frac{\partial}{\partial v} g_{j\ell}(u, v), \quad (3.9)$$

which are referred to as *consistency*, *conservativity*, and *monotonicity* of the numerical flux  $g_{j\ell}$ , respectively. (See [33] or [34] for more discussion on general upwind finite volume numerical fluxes.) Moreover, due to (3.7) and  $\operatorname{div} b(x) = 0$ , we have that (cf. (3.13)):

$$\sum_{\ell \in N_j} g_{j\ell}(u, u) = 0 \quad \text{for all } j. \quad (3.10)$$

**EXAMPLE 3.2. (First order Engquist-Osher scheme)** As a particular example of the numerical flux we choose the Engquist-Osher type upwind finite volume flux  $g_{j\ell}$  defined by

$$g_{j\ell}(u, v) := b_{j\ell}^+ u + b_{j\ell}^- v, \quad b_{j\ell}^\pm := \int_{S_{j\ell}} (bn_{j\ell})^\pm ds. \quad (3.11)$$

It can be easily shown that this particular numerical flux satisfies (3.6)–(3.10). The easiest choice of  $\mathcal{U}_{j\ell}$  in (3.5), namely

$$\mathcal{U}_{j\ell} := u_j, \quad \mathcal{U}_{\ell j} := u_\ell, \quad (3.12)$$

used together with (3.11) in (3.4) defines a *first order* numerical scheme.

**REMARK 3.3.** Due to the properties of  $b$  we have for all  $T_j \in \mathcal{T}_h$

$$\sum_{\ell \in N_j} (b_{j\ell}^+ + b_{j\ell}^-) = \sum_{\ell \in N_j} b_{j\ell} = \sum_{\ell \in N_j} \int_{S_{j\ell}} bn_{j\ell} ds = \int_{\partial T_j} bn_{j\ell} ds = \int_{T_j} \operatorname{div} b dx = 0, \quad (3.13)$$

and, for all  $S_{j\ell} \in \mathcal{E}$ ,

$$b_{\ell j} = -b_{j\ell}, \quad b_{\ell j}^+ = -b_{j\ell}^-, \quad b_{\ell j}^- = -b_{j\ell}^+. \quad (3.14)$$

**EXAMPLE 3.4. (Higher order scheme using MUSCL type reconstruction)** Let  $T_k, T_\ell, T_m$  be all neighboring triangles to  $T_j$  with centers of gravity  $x_k, x_\ell, x_m, x_j$ , respectively. Let  $w \in L^\infty(\Omega)$  with  $w|_{T_j} \in C^0(T_j)$  and  $w_i := w(x_i)$  for  $i = k, \ell, m, j$ , respectively. Let (cf. [19])

$$\begin{aligned} R_k^w &\text{ be a plane passing through } (x_\ell, w_\ell), (x_m, w_m), (x_j, w_j), \\ R_\ell^w &\text{ be a plane passing through } (x_k, w_k), (x_m, w_m), (x_j, w_j), \\ R_m^w &\text{ be a plane passing through } (x_k, w_k), (x_\ell, w_\ell), (x_j, w_j). \end{aligned}$$

Define an index  $i$  by

$$|\nabla R_i^w| = \min \{ |\nabla R_k^w|, |\nabla R_\ell^w|, |\nabla R_m^w| \} \quad (3.15)$$

and put

$$G_j^w := \nabla R_i^w. \quad (3.16)$$

If  $w_j \geq \max\{w_k, w_\ell, w_m\}$  or  $w_j \leq \min\{w_k, w_\ell, w_m\}$ , we say that  $w_j$  is a local extremum. Let the coefficients  $\alpha_j = \alpha_j^w \in \{0, 1\}$  be such that

$$\alpha_j^w = \begin{cases} 0 & \text{if } w_j \text{ is the local extremum,} \\ 1 & \text{otherwise.} \end{cases} \quad (3.17)$$

Then define

$$L_j^w(x) := w_j + \alpha_j^w G_j^w(x - x_j). \quad (3.18)$$

Finally, the higher order MUSCL type Engquist-Osher scheme is defined by (3.4) with the numerical flux (3.11) and

$$\mathcal{U}_{j\ell} := L_j^u(x_{j\ell}), \quad \mathcal{U}_{\ell j} := L_\ell^u(x_{j\ell}). \quad (3.19)$$

It can be shown that, on the regular grid, the reconstruction operator  $L_j^u$  defined by (3.18) has the following properties.

LEMMA 3.5. *For all  $T_j \in \mathcal{T}_R$  we have*

$$(a) \quad |L_j^u(x_\ell) - u_j| \leq |u_j - u_\ell| \quad \text{for all } \ell \in N_j, \quad (3.20)$$

$$(b) \quad |L_j^u(x_{j\ell}) - u_j| \leq \frac{1}{2}|u_j - u_\ell| \quad \text{for all } \ell \in N_j, \quad (3.21)$$

$$(c) \quad (u_\ell - L_j^u(x_\ell))(u_j - u_\ell) \leq 0 \quad \text{for all } \ell \in N_j. \quad (3.22)$$

We will give the proof of this lemma in the following section.

**4. Main result.** We will use the scheme (3.2)–(3.4) with the following definition of the numerical flux:

- If  $S_{j\ell} \in \mathcal{E}_R$  we use the higher order flux using MUSCL type reconstruction, i.e. we set (cf. (3.11), (3.18)–(3.19))

$$g_{j\ell}(\mathcal{U}_{j\ell}, \mathcal{U}_{\ell j}) := b_{j\ell}^+ L_j^u(x_{j\ell}) + b_{j\ell}^- L_\ell^u(x_{j\ell}). \quad (4.1)$$

- If  $S_{j\ell} \in \mathcal{E}_M$  or  $S_{j\ell} \in \mathcal{E}_S$  we use the first order flux, i.e. we set (cf. (3.11), (3.12))

$$g_{j\ell}(\mathcal{U}_{j\ell}, \mathcal{U}_{\ell j}) := b_{j\ell}^+ u_j + b_{j\ell}^- u_\ell. \quad (4.2)$$

- If  $S_{j\ell} \in \mathcal{E}_B$  we use

$$g_{j\ell}(\mathcal{U}_{j\ell}, \mathcal{U}_{\ell j}) := b_{j\ell}^+ u_j + b_{j\ell}^- u_\ell \quad (4.3)$$

where in this case  $u_\ell$  is the value in the ghost cell of the cell  $T_j$  satisfying  $u_\ell = -u_j$  (cf. (3.3)).

The main result of this paper is formulated in the following theorem.

**THEOREM 4.1.** *Let  $u_h(x) = u_j$  for  $x \in T_j \in \mathcal{T}_h$ , be a piecewise constant numerical solution of the discrete problem (3.2)–(3.4) with a numerical flux satisfying (3.11), (3.18)–(3.19), (4.1)–(4.3) and let the Assumption 2.4 hold. We define*

$$z_h := I_h v - u_h \quad (4.4)$$

where

$$I_h v(x) := v(\bar{x}_j) = v_j \quad \text{if } x \in T_j \in \mathcal{T}_h. \quad (4.5)$$

Then, defining

$$\|z_h\|_\varepsilon^2 := \varepsilon \gamma \sum_{\mathcal{E} \cup \mathcal{E}_B} (z_j - z_\ell)^2 + c_0 \sum_{T_j \in \mathcal{T}_h} z_j^2 |T_j|, \quad (4.6)$$

we have the following error estimate for any  $\delta > 0$ :

$$\|z_h\|_\varepsilon^2 \leq c \left( \varepsilon h^2 + h^{4-2\delta} + \frac{h^3}{\varepsilon} \right) \|v\|_{2,2}^2 + c \frac{h^4}{\varepsilon} \sum_{T_j \in \mathcal{T}_R} R_j^2 |T_j|. \quad (4.7)$$

If, moreover,  $v \in W^{3,2}(\Omega)$ , we have

$$\|z_h\|_\varepsilon^2 \leq c \left( \varepsilon h^3 + h^{4-2\delta} + \frac{h^3}{\varepsilon} \right) \|v\|_{3,2}^2 + c \frac{h^4}{\varepsilon} \sum_{T_j \in \mathcal{T}_R} R_j^2 |T_j|. \quad (4.8)$$

Here,  $R_j := \frac{1}{|T_j|} \sum_{\ell \in N_j} (u_\ell - u_j) \gamma_{j\ell}$ .

**REMARK 4.2.**

- Note that if  $T_\ell$  is the mirrored ghost triangle to  $T_j$ , we have  $v(\bar{x}_\ell) = -v(\bar{x}_j)$  (see Assumption 2.4), and also  $u_\ell = -u_j$  (see (3.3)). Therefore,

$$z_\ell = v_\ell - u_\ell = -v_j + u_j = -z_j, \quad (4.9)$$

if  $T_\ell$  is the mirrored ghost triangle to  $T_j$ .

- In the case when the first order scheme is used in the whole domain we get (for comparison) the following result:

$$\|z_h\|_\varepsilon^2 \leq c \left( \varepsilon h^2 + h^{4-2\delta} + \frac{h^2}{\varepsilon} \right) \|v\|_{2,2}^2. \quad (4.10)$$

For the higher order MUSCL type scheme we thus gain  $h^{1/2}$  inside the estimate of the norm of  $\|z_h\|_\varepsilon$  for the term corresponding (as it shows) to the convective part of the equation, compared to the first order scheme: compare  $\frac{h^3}{\varepsilon}$  to  $\frac{h^2}{\varepsilon}$  in the estimate of the norm of  $\|z_h\|_\varepsilon^2$ .

- For the context of the definition of  $R_j$  see (8.6). It follows from Lemma 8.4 that the sum  $\sum_{T_j \in \mathcal{T}_R} R_j^2 |T_j|$  is of the same order in  $\varepsilon$  as  $\|v\|_{2,2}^2(\Omega)$ , cf. (8.14) and (8.20), namely we have both

$$\sum_{T_j \in \mathcal{T}_R} R_j^2 |T_j| \leq \frac{c}{\varepsilon^3} \|f\|_{L^2(\Omega)}^2 \quad \text{and} \quad \|v\|_{W^{2,2}(\Omega)}^2 \leq \frac{c}{\varepsilon^3} \|f\|_{L^2(\Omega)}^2,$$

the second estimate being sharp due to Remark 8.5. In such a way, the estimate (4.7) can be put in form of

$$\|z_h\|_\varepsilon^2 \leq c \left( \varepsilon h^2 + h^{4-2\delta} + \frac{h^3}{\varepsilon} \right) \|v\|_{2,2}^2 + c \frac{h^4}{\varepsilon^4} \|f\|_{L^2(\Omega)}^2$$

or

$$\|z_h\|_\varepsilon^2 \leq c \left( \varepsilon h^2 + h^{4-2\delta} + \frac{h^3}{\varepsilon} \right) \frac{1}{\varepsilon^3} \|f\|_{L^2(\Omega)}^2 + c \frac{h^4}{\varepsilon^4} \|f\|_{L^2(\Omega)}^2$$

while in the case of the first order scheme we would get analogously instead of (4.10) the following estimate:

$$\|z_h\|_\varepsilon^2 \leq c \left( \varepsilon h^2 + h^{4-2\delta} + \frac{h^2}{\varepsilon} \right) \frac{1}{\varepsilon^3} \|f\|_{L^2(\Omega)}^2.$$

- For the particular numerical calculation for which  $\varepsilon \approx h$ , we get, using (4.10) and (4.7), the error estimates of the order  $O(\sqrt{h})$  and  $O(h)$  in the cases of first order and higher order scheme, respectively. If  $\varepsilon \approx \sqrt{h}$ , we get in the corresponding cases the error estimates of the order  $O(h^{3/4})$  and  $O(h^{5/4})$ , respectively.
- In the estimate (4.7), the norm  $\|v\|_{2,2}^2$  still depends singularly on  $\varepsilon$ . Due to Lemma 8.4 it behaves like  $\frac{1}{\varepsilon^3}$ . This means altogether the term  $\frac{h^3}{\varepsilon} \|v\|_{2,2}^2$  behaves like  $\frac{h^3}{\varepsilon^4}$ . These types of estimate are usual also for stabilized finite elements but for numerical computations on a non adaptive grid they are of limited importance. Nevertheless numerical experiments indicate that adaptive grids in the neighbourhood of the boundary layer allow acceptable resolutions.

In the remaining part of the paper we will give the proof of Theorem 4.1.

**5. The basic estimates.** First we give a proof of Lema 3.5.

*Proof of Lemma 3.5 from page 7:* Of course, for  $T_j \in \mathcal{T}_R$ , (b) follows immediately from (a). To prove (a), let us consider the following geometrical situation. Without loss of generality we assume that  $x_j = (0, 0)$ ,  $x_\ell = (1, 0)$ ,  $x_m = (\cos \alpha, \sin \alpha)$ ,  $x_k = (\cos \alpha, -\sin \alpha)$ ,  $\alpha = \frac{2}{3}\pi$ . Therefore,

$$\sin \alpha = \frac{\sqrt{3}}{2} \quad \text{and} \quad \cos \alpha = -\frac{1}{2}. \quad (5.1)$$

Let  $u_j, u_\ell, u_m, u_k$  be arbitrary. Denote  $P_i = (x_i, u_i)$  for  $i = j, \ell, m, k$ . The planes  $p_k, p_\ell, p_m$  through the set of points  $(P_j, P_\ell, P_m)$ ,  $(P_j, P_m, P_k)$ ,  $(P_j, P_\ell, P_k)$ , respectively, are given by

$$p_k(y) = u_j + (u_\ell - u_j)y_1 + \frac{(u_m - u_j) - \cos \alpha(u_\ell - u_j)}{\sin \alpha} y_2, \quad (5.2)$$

$$p_\ell(y) = u_j + \frac{(u_m - u_j) + (u_k - u_j)}{2 \cos \alpha} y_1 + \frac{(u_m - u_j) - (u_k - u_j)}{2 \sin \alpha} y_2, \quad (5.3)$$

$$p_m(y) = u_j + (u_\ell - u_j)y_1 - \frac{(u_k - u_j) - \cos \alpha(u_\ell - u_j)}{\sin \alpha} y_2, \quad (5.4)$$

where  $y = (y_1, y_2)$ . If there is a local extremum in  $x_j$  we have (see (3.18))  $L_j^u(x_i) = u_j$  for  $i = k, \ell, m$  and (3.20) holds. Therefore we can assume that there is no local extremum in  $x_j$ , i.e.,  $\alpha_j^u = 1$ . Furthermore we assume without loss of generality that (see (3.18))

$$\nabla L_j^u = \nabla p_m,$$

which means that

$$|\nabla p_m| \leq |\nabla p_\ell|, \quad (5.5)$$

$$|\nabla p_m| \leq |\nabla p_k|. \quad (5.6)$$

Considering  $L_j^u(x_i) - u_j$  for  $i = k, \ell, m$ , we will discuss two cases.

First case: if  $i = \ell$ , or  $i = k$ , we obtain

$$\begin{aligned} L_j^u(x_\ell) - u_j &= \nabla p_m \cdot (1, 0) = u_\ell - u_j, \\ L_j^u(x_k) - u_j &= \nabla p_m \cdot (\cos \alpha, -\sin \alpha) = u_k - u_j, \end{aligned}$$

and (3.20) follows.

Second case: if  $i = m$ , we have

$$L_j^u(x_m) - u_j = \nabla p_m \cdot (\cos \alpha, \sin \alpha) = (u_j - u_k) + (u_j - u_\ell),$$

using (5.1). Therefore, in order to prove (3.20) we have to show that

$$|(u_j - u_k) + (u_j - u_\ell)| \leq |u_j - u_m|. \quad (5.7)$$

Using (5.1) we obtain for the gradients of  $p_k, p_\ell, p_m$ ,

$$\begin{aligned} |\nabla p_k|^2 &= \frac{4}{3} [(u_\ell - u_j)^2 + (u_m - u_j)^2 + (u_\ell - u_j)(u_m - u_j)] \\ |\nabla p_\ell|^2 &= \frac{4}{3} [(u_m - u_j)^2 + (u_k - u_j)^2 + (u_m - u_j)(u_k - u_j)] \\ |\nabla p_m|^2 &= \frac{4}{3} [(u_k - u_j)^2 + (u_\ell - u_j)^2 + (u_k - u_j)(u_\ell - u_j)]. \end{aligned} \quad (5.8)$$

It follows that (5.5), (5.6) are equivalent to

$$(u_m - u_\ell)(u_m + u_k + u_\ell - 3u_j) \geq 0, \quad (5.9)$$

$$(u_m - u_k)(u_m + u_k + u_\ell - 3u_j) \geq 0, \quad (5.10)$$

respectively.

We will further distinguish three cases:

- (i)  $u_m + u_k + u_\ell - 3u_j = 0$ . This implies  $(u_j - u_k) + (u_j - u_\ell) = u_m - u_j$  and (5.7) follows.
- (ii)  $u_m + u_k + u_\ell - 3u_j > 0$ . Then (5.10) and (5.9) imply  $u_m \geq u_k$  and  $u_m \geq u_\ell$ , respectively. Since  $u_j$  is not a local maximum, we also have  $u_m \geq u_j$ . Then

$$\begin{aligned} u_j - u_k + u_j - u_\ell &= 3u_j - u_k - u_\ell - u_m + u_m - u_j \\ &< u_m - u_j = |u_m - u_j|. \end{aligned}$$

On the other hand,  $u_j \geq \min(u_k, u_\ell)$ , since  $u_j$  is not a local minimum. Therefore,  $u_j + u_m \geq u_k + u_\ell$  and

$$u_j - u_k + u_j - u_\ell \geq u_j - u_m = -|u_m - u_j|.$$

Altogether, (5.7) follows.

- (iii)  $u_m + u_k + u_\ell - 3u_j < 0$ . In this case we proceed quite analogously as in the case (ii) which finishes the proof of Lemma 3.5, part (a).

Finally let us prove (c). We have to show

$$\left(u_i - L_j^u(x_i)\right)(u_j - u_i) \leq 0, \quad i = k, \ell, m. \quad (5.11)$$

Using the same notation as in the part (a), we see that  $L_j^u(x_i) = u_i$  for  $i = k$  or  $i = \ell$ , and (5.11) follows. For  $i = m$  we have

$$\left(u_m - L_j^u(x_m)\right)(u_j - u_m) = (u_m + u_k + u_\ell - 3u_j)(u_j - u_m). \quad (5.12)$$

Discussing the cases (i), (ii), (iii) as in the part (a), we see that (5.11) follows. The Lemma 3.5 is proven.  $\square$

We continue with a rather technical result. Sometimes we have to change the sum over all triangles into a sum over all edges of the triangulation. The general rules for this are given by the following lemma.

LEMMA 5.1. *Let  $A_{j\ell} \in \mathbb{R}$  for all  $j, T_j \in \mathcal{T}_h$ . Then we have*

$$\begin{aligned} \sum_{T_j \in \mathcal{T}_h} \sum_{\ell \in N_j} A_{j\ell} &= \sum_{\text{edges} \in \mathcal{E}_R} \tilde{A}_{j\ell} + \sum_{\text{edges} \in \mathcal{E}_S} \tilde{A}_{j\ell} + \sum_{\text{edges} \in \mathcal{E}_M} \tilde{A}_{j\ell} + \sum_{\text{edges} \in \mathcal{E}_B} A_{j\ell} \\ &= \sum_{\text{edges} \in \mathcal{E}} \tilde{A}_{j\ell} + \sum_{\text{edges} \in \mathcal{E}_B} A_{j\ell} \end{aligned} \quad (5.13)$$

where  $\tilde{A}_{j\ell} := A_{j\ell} + A_{\ell j}$  and  $\mathcal{E} = \mathcal{E}_R \cup \mathcal{E}_S \cup \mathcal{E}_M$ .

*Proof:*

$$\begin{aligned} \sum_{T_j \in \mathcal{T}_h} \sum_{\ell \in N_j} A_{j\ell} &= \sum_{T_j \in \mathcal{T}_R} \sum_{\ell \in N_j} A_{j\ell} + \sum_{T_j \in \mathcal{T}_S} \sum_{\ell \in N_j} A_{j\ell} \\ &= \sum_{T_j \in \mathcal{T}_R} \sum_{\ell \in N_{jR}} A_{j\ell} + \sum_{T_j \in \mathcal{T}_R} \sum_{\ell \in N_{jS}} A_{j\ell} + \sum_{T_j \in \mathcal{T}_R} \sum_{\ell \in N_{jG}} A_{j\ell} \\ &\quad + \sum_{T_j \in \mathcal{T}_S} \sum_{\ell \in N_{jR}} A_{j\ell} + \sum_{T_j \in \mathcal{T}_S} \sum_{\ell \in N_{jS}} A_{j\ell} + \sum_{T_j \in \mathcal{T}_S} \sum_{\ell \in N_{jG}} A_{j\ell} \\ &= \sum_{\text{edges} \in \mathcal{E}_R} \tilde{A}_{j\ell} + \sum_{\text{edges} \in \mathcal{E}_S} \tilde{A}_{j\ell} + \sum_{\text{edges} \in \mathcal{E}_M} \tilde{A}_{j\ell} + \sum_{\text{edges} \in \mathcal{E}_B} A_{j\ell} \end{aligned}$$

where  $\tilde{A}_{j\ell} := A_{j\ell} + A_{\ell j}$ .  $\square$

REMARK 5.2. We often abbreviate  $\sum_{\mathcal{E}} \equiv \sum_{\text{edges} \in \mathcal{E}}$  and so on.

In what follows we prove the discrete energy estimate for the higher order scheme.

LEMMA 5.3. *Let  $u_h$  be the numerical solution defined by the scheme (3.2)–(3.4) with a numerical flux satisfying (3.11), (3.18)–(3.19), (4.1)–(4.3) and let Assumption 2.4 hold. Then there is a constant  $c > 0$  such that for all  $\varepsilon > 0$  and all  $h > 0$*

$$\varepsilon \gamma \sum_{\mathcal{E} \cup \mathcal{E}_B} (u_j - u_\ell)^2 + c_0 \sum_{T_j \in \mathcal{T}_h} u_j^2 |T_j| \leq c \sum_{T_j \in \mathcal{T}_h} f_j^2 |T_j|. \quad (5.14)$$

*Proof:* We multiply  $(L_h u_h)_j = f_j$  by  $|T_j| u_j$  and sum up over all  $j$  such that  $T_j \in \mathcal{T}_h$ . Then we obtain

$$-\varepsilon \sum_j \sum_{\ell \in N_j} (u_\ell - u_j) \gamma_{j\ell} u_j + \sum_j \sum_{\ell \in N_j} g_{j\ell} (\mathcal{U}_{j\ell}, \mathcal{U}_{\ell j}) u_j + \sum_j c_j u_j^2 |T_j| = \sum_j f_j u_j |T_j|. \quad (5.15)$$

For the first term on the left hand side in (5.15) we get using  $\gamma_{j\ell} = \gamma_{\ell j}$  and (3.3),

$$\begin{aligned} -\varepsilon \sum_j \sum_{\ell \in N_j} (u_\ell - u_j) \gamma_{j\ell} u_j &= -\varepsilon \sum_j \sum_{\ell \in N_{jI}} (u_\ell - u_j) \gamma_{j\ell} u_j - \varepsilon \sum_j \sum_{\ell \in N_{jG}} (u_\ell - u_j) \gamma_{j\ell} u_j \\ &= -\varepsilon \sum_{\mathcal{E}} ((u_\ell - u_j) \gamma_{j\ell} u_j + (u_j - u_\ell) \gamma_{\ell j} u_\ell) + \varepsilon \sum_j \sum_{\ell \in N_{jG}} 2u_j^2 \gamma_{j\ell} \\ &= \varepsilon \sum_{\mathcal{E}} (u_\ell - u_j)^2 \gamma_{j\ell} + \frac{\varepsilon}{2} \sum_j \sum_{\ell \in N_{jG}} (u_\ell - u_j)^2 \gamma_{j\ell} \\ &\geq \frac{\varepsilon \gamma}{2} \sum_{\mathcal{E} \cup \mathcal{E}_B} (u_\ell - u_j)^2. \end{aligned} \quad (5.16)$$

For the second term on the left hand side of (5.15) we get

$$\begin{aligned} \sum_j \sum_{\ell \in N_j} g_{j\ell}(\mathcal{U}_{j\ell}, \mathcal{U}_{\ell j}) u_j &= \sum_j \sum_{\ell \in N_j} (b_{j\ell}^+ u_j + b_{j\ell}^- u_\ell) u_j \\ &\quad + \sum_{T_R} \sum_{\ell \in N_{jR}} (b_{j\ell}^+ \alpha_j^u G_j^u(x_{j\ell} - x_j) + b_{j\ell}^- \alpha_\ell^u G_\ell^u(x_{j\ell} - x_\ell)) u_j \\ &=: B_1 + B_2 \end{aligned}$$

Due to (3.13) we have  $\sum_{\ell \in N_j} (b_{j\ell}^+ + b_{j\ell}^-) u_j^2 = u_j^2 \sum_{\ell \in N_j} b_{j\ell} = 0$  for all  $j$ , and we can proceed, using also (3.14) and (3.3), as follows:

$$\begin{aligned} B_1 &= \sum_j \sum_{\ell \in N_j} (b_{j\ell}^+ u_j^2 + b_{j\ell}^- u_\ell u_j) = \sum_j \sum_{\ell \in N_j} (b_{j\ell}^+ u_j^2 + b_{j\ell}^- u_\ell u_j - \frac{1}{2}(b_{j\ell}^+ + b_{j\ell}^-) u_j^2) \\ &= \frac{1}{2} \sum_j \sum_{\ell \in N_j} (b_{j\ell} u_\ell + b_{j\ell}^+(u_j - u_\ell) - b_{j\ell}^-(u_j - u_\ell)) u_j = \frac{1}{2} \sum_j \sum_{N_{jI}} \dots + \frac{1}{2} \sum_j \sum_{N_{jG}} \dots \\ &= \frac{1}{2} \sum_{\mathcal{E}} (b_{j\ell} u_\ell u_j + b_{\ell j} u_j u_\ell) \\ &\quad + \frac{1}{2} \sum_{\mathcal{E}} (b_{j\ell}^+(u_j - u_\ell) u_j - b_{j\ell}^-(u_j - u_\ell) u_j - b_{j\ell}^-(u_\ell - u_j) u_\ell + b_{j\ell}^+(u_\ell - u_j) u_\ell) \\ &\quad + \frac{1}{2} \sum_j \sum_{N_{jG}} \dots \\ &= 0 + \frac{1}{2} \sum_{\mathcal{E}} (b_{j\ell}^+(u_j - u_\ell)^2 - b_{j\ell}^-(u_j - u_\ell)^2) + \frac{1}{2} \sum_j \sum_{N_{jG}} (-3b_{j\ell}^- u_j + b_{j\ell}^+ u_j) u_j \\ &\geq \frac{1}{2} \sum_{\mathcal{E}} (b_{j\ell}^+(u_j - u_\ell)^2 - b_{j\ell}^-(u_j - u_\ell)^2) = \frac{1}{2} \sum_{\mathcal{E}} (b_{j\ell}^+ - b_{j\ell}^-) (u_j - u_\ell)^2. \end{aligned} \tag{5.17}$$

Now let us treat the term  $B_2$ .

$$\begin{aligned} B_2 &= \sum_{T_R} \sum_{\ell \in N_{jR}} (b_{j\ell}^+ \alpha_j^u G_j^u(x_{j\ell} - x_j) + b_{j\ell}^- \alpha_\ell^u G_\ell^u(x_{j\ell} - x_\ell)) u_j \\ &= \sum_{\mathcal{E}_R} ((b_{j\ell}^+ \alpha_j^u G_j^u(x_{j\ell} - x_j) + b_{j\ell}^- \alpha_\ell^u G_\ell^u(x_{j\ell} - x_\ell)) u_j \\ &\quad + (b_{\ell j}^+ \alpha_\ell^u G_\ell^u(x_{\ell j} - x_\ell) + b_{\ell j}^- \alpha_j^u G_j^u(x_{\ell j} - x_j)) u_\ell) \\ &= \sum_{\mathcal{E}_R} (b_{j\ell}^+ \alpha_j^u G_j^u(x_{j\ell} - x_j) (u_j - u_\ell) + b_{j\ell}^- \alpha_\ell^u G_\ell^u(x_{j\ell} - x_\ell) (u_j - u_\ell)) \\ &\geq - \sum_{\mathcal{E}_R} (b_{j\ell}^+ \alpha_j^u |G_j^u(x_{j\ell} - x_j)| |u_j - u_\ell| + |b_{j\ell}^- \alpha_\ell^u G_\ell^u(x_{j\ell} - x_\ell)| |u_j - u_\ell|). \end{aligned}$$

Now since

$$\alpha_j^u |G_j^u(x_{j\ell} - x_j)| = \frac{1}{2} \alpha_j^u |G_j^u(x_\ell - x_j)| = \frac{1}{2} |(L_j^u(x_\ell) - u_j)| \leq \frac{1}{2} |u_\ell - u_j|$$

(see Lemma 3.5), we can continue

$$\geq -\frac{1}{2} \sum_{\mathcal{E}_R} (b_{j\ell}^+ |u_j - u_\ell|^2 + |b_{j\ell}^-| |u_j - u_\ell|^2) = -\frac{1}{2} \sum_{\mathcal{E}_R} (b_{j\ell}^+ - b_{j\ell}^-) (u_j - u_\ell)^2. \tag{5.18}$$

Finally, using the estimate  $\sum_j f_j u_j |T_j| \leq \frac{c_0}{2} \sum_j u_j^2 |T_j| + c \sum_j f_j^2 |T_j|$  and (5.15)–(5.18), we get (5.14).  $\square$

In the next lemma we give the basic local estimates for the exact solution  $v$ .

LEMMA 5.4. *For  $v \in W^{2,2}(\Omega)$  the following estimates hold.*

a) *For all  $S_{j\ell}$ , and for all  $x, y \in T_j \cup T_\ell$  we have*

$$|v(x) - v(y)| \leq ch |v|_{W^{2,2}(Q_{j\ell})} + c|v|_{W^{1,2}(Q_{j\ell})}. \quad (5.19)$$

*Here,  $Q_{j\ell}$  is a rectangular domain containing  $T_j \cup T_\ell$  and a finite fixed number of neighboring triangles. The number of these triangles is independent of  $h$  (for more details see the proof).*

b) *For all  $S_{j\ell}$  we have*

$$|\kappa_{j\ell}(v)| := \left| \frac{v_\ell - v_j}{d_{j\ell}} - \frac{1}{|S_{j\ell}|} \int_{S_{j\ell}} \partial_n v \right| \leq c |v|_{W^{2,2}(Q_j)}. \quad (5.20)$$

*Here,  $Q_j$  is a finite union of triangles containing the rectangular  $S_{j\ell} \times [\overline{x_{j\ell}}, \overline{x_j}]$ .*

c) *For  $S_{j\ell} \in \mathcal{E}_R$ , and if the solution  $v$  moreover satisfies  $v \in W^{3,2}(\Omega)$ , we have*

$$|\kappa_{j\ell}(v)| := \left| \frac{v_\ell - v_j}{d_{j\ell}} - \frac{1}{|S_{j\ell}|} \int_{S_{j\ell}} \partial_n v \right| \leq ch |v|_{W^{3,2}(T_j \cup T_\ell)}. \quad (5.21)$$

d) *For all  $T_j \in \mathcal{T}_h$  we have*

$$\left| \int_{T_j} (v(\overline{x_j}) - v(x)) dx \right| \leq ch^{3-\delta} \|v\|_{W^{2,2}(T_j)}. \quad (5.22)$$

for all  $\delta > 0$ .

*Proof of a):* Suppose  $T_j \in \mathcal{T}_h$  and that either  $T_\ell \in T_h$  (in this case we have  $S_{j\ell} \in \mathcal{E}$ ) or  $T_\ell \in \mathcal{T}_G$  (in this case we have  $S_{j\ell} \in \mathcal{E}_B$ ). Without loss of generality we can suppose that there is a local cartesian coordinate system  $(x, y)$  such that the origin of this system coincides with  $x_{j\ell}$ , the  $x$  axis is aligned with the line through  $\overline{x_\ell}$  and  $\overline{x_j}$  and that the vertices of  $S_{j\ell}$  are given by  $(0, -\alpha)$  and  $(0, \alpha)$ . Let also  $x = (x_1, x_2)$ ,  $y = (y_1, y_2) \in T_j \cup T_\ell$ . We write

$$v(x) - v(y) = \underbrace{v(x_1, x_2) - v(x_1, y_2)}_{=: R_1} + \underbrace{v(x_1, y_2) - v(y_1, y_2)}_{=: R_1}. \quad (5.23)$$

First we will estimate  $R_1$ . To this point, consider the rectangular  $Q_j = [-3h, 3h] \times [-\alpha, \alpha]$  and a smooth cut-off 1D function  $\phi$  such that  $\phi = 0$  on  $(-3h, -2h)$  and on  $(2h, 3h)$ , and  $\phi = 1$  on  $(-h, h)$ , with  $0 \leq \phi \leq 1$  and  $|\phi'| \leq \frac{c}{h}$ . Then we have

$$\begin{aligned} R_1 &= \int_{y_2}^{x_2} \partial_2 v(x_1, s) ds \\ &= \int_{y_2}^{x_2} \left[ \partial_2 v(x_1, s) \phi(x_1) - \partial_2 v(-3h, s) \phi(-3h) \right] ds \\ &= \int_{y_2}^{x_2} \int_{-3h}^{x_1} \partial_t (\partial_2 v(t, s) \phi(t)) dt ds \\ &= \int_{y_2}^{x_2} \int_{-3h}^{x_1} \partial_t \partial_2 v(t, s) \phi(t) dt ds + \int_{y_2}^{x_2} \int_{-3h}^{x_1} \partial_2 v(t, s) \phi'(t) dt ds. \end{aligned}$$

Therefore,

$$\begin{aligned} |R_1| &\leq \int_{\min(x_2, y_2)}^{\max(x_2, y_2)} \int_{-3h}^{3h} \left( |\partial_{12} v(t, s) \phi(t)| + |\partial_2 v(t, s) \phi'(t)| \right) dt ds \\ &\leq \int_{Q_j} |\partial_{12} v(t, s)| dt ds + \frac{c}{h} \int_{Q_j} |\partial_2 v(t, s)| dt ds \\ &\leq ch |v|_{W^{2,2}(Q_j)} + |v|_{W^{1,2}(Q_j)}. \end{aligned}$$

For  $R_2$  we proceed in a similar way, using the cut-off function defined on  $[-h, h] \times -[3\alpha, 3\alpha]$ .

*Proof of b):* Suppose  $T_j \in \mathcal{T}_h$  and that either  $T_\ell \in T_h$  (in this case we have  $S_{j\ell} \in \mathcal{E}$ ) or  $T_\ell \in \mathcal{T}_G$  (in this case we have  $S_{j\ell} \in \mathcal{E}_B$ ). Consider the same local cartesian coordinate system as in case a). Then let  $\bar{x}_\ell =: (a, 0)$  and  $\bar{x}_j =: (b, 0)$ ,  $a < b$ . Using the notation  $v_j := v(\bar{x}_j)$ ,  $v_\ell := v(\bar{x}_\ell)$  we obtain in both cases, using  $\gamma_{j\ell} = \frac{|S_{j\ell}|}{d_{j\ell}} = \frac{2\alpha}{b-a}$ ,

$$\begin{aligned}
(v_\ell - v_j)\gamma_{j\ell} - \int_{S_{j\ell}} \partial_n v &= (v(a, 0) - v(b, 0))\frac{2\alpha}{b-a} + \int_{-\alpha}^{\alpha} \partial_1 v(0, y) dy \\
&= \frac{2\alpha}{b-a} \int_b^a \partial_1 v(x, 0) dx + \int_{-\alpha}^{\alpha} \partial_1 v(0, y) dy \\
&= \frac{1}{b-a} \int_{-\alpha}^{\alpha} \int_a^b -\partial_1 v(x, 0) dx dy + \frac{1}{b-a} \int_a^b \int_{-\alpha}^{\alpha} \partial_1 v(0, y) dy dx \\
&= \frac{1}{b-a} \int_{-\alpha}^{\alpha} \int_a^b \left[ \partial_1 v(0, y) - \partial_1 v(x, y) + \partial_1 v(x, y) - \partial_1 v(x, 0) \right] dx dy \\
&= \frac{1}{b-a} \int_{-\alpha}^{\alpha} \int_a^b \left[ \int_x^0 \partial_{11} v(t, y) dt + \int_0^y \partial_{12} v(x, s) ds \right] dx dy.
\end{aligned}$$

This implies

$$\begin{aligned}
\left| (v_\ell - v_j)\gamma_{j\ell} - \int_{S_{j\ell}} \partial_n v \right| &\leq \frac{1}{b-a} \int_{-\alpha}^{\alpha} \int_a^b \int_a^b |\partial_{11} v(t, y)| dt dx dy \\
&\quad + \frac{1}{b-a} \int_{-\alpha}^{\alpha} \int_a^b \int_{-\alpha}^{\alpha} |\partial_{12} v(x, s)| ds dx dy \\
&\leq \int_{\tilde{Q}_j} |\partial_{11} v(x, s)| ds dx + \frac{2\alpha}{b-a} \int_{\tilde{Q}_j} |\partial_{12} v(x, s)| ds dx \\
&\leq ch|v|_{W^{2,2}(Q_j)}
\end{aligned}$$

where  $Q_j$  is a union of triangles which contains the rectangular  $[a, b] \times [-\alpha, \alpha]$ .

*Proof of c):* Since the constant  $c > 0$  should not depend on  $T_j$ ,  $T_\ell$ ,  $h$ , we use the following transformation mapping. Let  $T := T_{j\ell} = T_j \cup T_\ell$  be a fixed rhombus in the computational domain and  $\hat{T}$  the reference rhombus with vertices  $P_1 = (1, 0)$ ,  $P_2 = (0, 1)$ ,  $P_3 = (-1, 0)$ ,  $P_4 = (0, -1)$ . Suppose  $\hat{T}$  is mapped onto  $T$  by a 1-1 mapping

$$\begin{aligned}
F : \hat{T} &\rightarrow T, \\
F(\hat{x}) &= A\hat{x} + a,
\end{aligned}$$

where  $A \in \mathbb{R}^{2 \times 2}$  is an invertible matrix and  $a \in \mathbb{R}^2$ , such that

- the oriented segment  $S$  of length 2 connecting points  $P_3$  and  $P_1$  is mapped onto the oriented segment  $S_{j\ell}$ . I.e., denoting  $\hat{\xi} := (2, 0)^T$ , we have  $|A\hat{\xi}| = |S_{j\ell}|$ .
- if  $n = n_{j\ell} = \frac{x_\ell - x_j}{|x_\ell - x_j|}$  is the outward (with respect to  $T_j$ ) unit normal vector to  $S_{j\ell}$ , then

$$\hat{n} \equiv \hat{n}(\hat{x}) := A^{-1}n(F(\hat{x})) = A^{-1}n = \frac{A^{-1}x_\ell - A^{-1}x_j}{|x_\ell - x_j|} = \frac{\hat{x}_\ell - \hat{x}_j}{|\hat{x}_\ell - \hat{x}_j|}$$

and therefore

$$|A\hat{n}| = 1.$$

- The properties of  $A$  imply

$$\|A\| \leq ch, \quad \|A^{-1}\| \leq ch^{-1}, \quad |\det A|^{-1} \leq ch^{-2}.$$

A map with these properties exists (see Heinrich [27, Section 5.2.2], and Ciarlet [11]), since all involved triangles are equilateral. Using the above notation, the following transformation rule holds:

$$\int_{S_{j\ell}} v \, d\sigma(x) = \int_S (v \circ F) \frac{|A\hat{\xi}|}{|\hat{\xi}|} \, d\sigma(\hat{x}).$$

Furthermore for  $\hat{v}(\hat{x}) := v(F(\hat{x}))$ , we have

$$\nabla_{\hat{x}} \hat{v}(\hat{x}) = A^T \cdot \nabla v(x), \quad x = F(\hat{x}),$$

and therefore

$$\begin{aligned} \frac{1}{|S_{j\ell}|} \int_{S_{j\ell}} \partial_n v &= \frac{1}{|S_{j\ell}|} \int_{S_{j\ell}} n(x) \nabla v(x) \, d\sigma \\ &= \frac{1}{|A\hat{\xi}|} \int_S A\hat{n}(A^T)^{-1} \nabla_{\hat{x}} \hat{v} \frac{|A\hat{\xi}|}{|\hat{\xi}|} \, d\sigma(\hat{x}) = \frac{A\hat{n}}{|\hat{\xi}|} \int_S (A^T)^{-1} \nabla_{\hat{x}} \hat{v} \, d\sigma(\hat{x}). \end{aligned}$$

We again denote  $\kappa_{j\ell}(v) := \frac{v_\ell - v_j}{d_{j\ell}} - \frac{1}{|S_{j\ell}|} \int_{S_{j\ell}} \partial_n v$ . (cf. (5.20)) Note that since  $W^{2,2}$  is imbedded into  $C^{0,\alpha}$  for any  $\alpha \in (0, 1)$ ,  $\kappa_{j\ell}(v)$  is well defined. Hence we can define

$$H(\hat{v}) := \frac{\hat{v}(\hat{x}_\ell) - \hat{v}(\hat{x}_j)}{|A(\hat{x}_\ell - \hat{x}_j)|} - \frac{A\hat{n}(A^T)^{-1}}{|\hat{\xi}|} \int_S \nabla_{\hat{x}} \hat{v} \, d\sigma(\hat{x}).$$

We see that

$$H(\hat{v}) = \kappa_{j\ell}(v).$$

Hence,

$$\begin{aligned} |H(\hat{v})| &\leq \frac{|\hat{v}(\hat{x}_\ell) - \hat{v}(\hat{x}_j)|}{|A(\hat{x}_\ell - \hat{x}_j)|} + \frac{|A\hat{n}| |(A^T)^{-1}|}{|\hat{\xi}|} \int_S |\nabla_{\hat{x}} \hat{v}| \, d\sigma(\hat{x}) \\ &\leq ch^{-1} |\hat{v}|_{C(\hat{T})} + ch^{-1} \|\nabla \hat{v}\|_{L^1(S)} \\ &\leq ch^{-1} \|\hat{v}\|_{W^{2,2}(\hat{T})} \leq ch^{-1} \|\hat{v}\|_{W^{k,2}(\hat{T})} \end{aligned}$$

where  $k \geq 2$ . Next we show that  $H$  vanishes for all polynomials  $p$  of degree  $\leq k$ . Let us assume without restriction that the vortices of the two neighboring triangles which build the rhombus are, as before, given by  $P_1 := (1, 0)$ ,  $P_2 := (0, 1)$ ,  $P_3 := (-1, 0)$ ,  $P_4 := (0, -1)$ . Let

$$p(y_1, y_2) = by_1^2 + cy_2^2 + dy_1 + ey_2 + f$$

and let  $(x_1, 0)$ ,  $(-x_1, 0)$  be the centers of gravity of the triangles  $P_1P_2P_4$  and  $P_2P_3P_4$ , respectively. Then we have

$$\begin{aligned} H(\hat{p}) = \kappa_{j\ell}(p) &= \frac{p(x_1, 0) - p(-x_1, 0)}{|(x_1, 0) - (-x_1, 0)|} - \frac{1}{2} \int_{-\frac{1}{2}}^{-\frac{1}{2}} n \nabla p \\ &= \frac{2d \cdot x_1}{2 \cdot x_1} - \frac{1}{2} \int_{-\frac{1}{2}}^{-\frac{1}{2}} \begin{pmatrix} 1 \\ 0 \end{pmatrix} \cdot \begin{pmatrix} d \\ 2cy_2 + e \end{pmatrix} dy_2 \\ &= d - \frac{1}{2} \cdot 2d = 0. \end{aligned}$$

Due to the properties of  $H$  we can use the Bramble–Hilbert lemma (see for example [11]) to obtain

$$|H(\widehat{v})| \leq c \|H\|_{2,2}^* |\widehat{v}|_{W^{k,2}(\widehat{T})} \leq c h^{-1} |\widehat{v}|_{W^{k,2}(\widehat{T})}.$$

Finally,

$$\begin{aligned} |\kappa_{j\ell}(v)| &= |H(\widehat{v})| \leq c h^{-1} |\widehat{v}|_{W^{k,2}(\widehat{T})} \\ &\leq c h^{-1} |\det A|^{-1/2} \|A\|^k |v|_{W^{k,2}(T)} \\ &\leq c h^{-1} h^{-1} h^k |v|_{W^{k,2}(T)} \end{aligned}$$

which proves for  $k = 2$  again the assertion b) of the Lemma (along singular edges) and for  $k = 3$  the assertion c) of the Lemma.

*Proof of d):* Let  $\overline{x}_j$  be the intersection of the perpendicular bisectors in  $T_j$  and

$$G(v) := \int_{T_j} v \, dx - v(\overline{x}_j) |T_j|.$$

Since the last term is an integration formula which is exact for constant functions and since the imbedding

$$W^{1,2+r}(T_j) \hookrightarrow L^\infty(T_j)$$

exists for any  $r > 0$ , we can apply [20, Satz 7.4] or [1, Lemma 4.3.8] and obtain

$$G(v) \leq c h^{3-\frac{2}{2+r}} |v|_{W^{1,2+r}(T_j)} \leq c h^{3-\delta} \|v\|_{W^{2,2}(T_j)}$$

for arbitrary  $\delta > 0$ .

This proves d).  $\square$

LEMMA 5.5. *As a consequence, we have for all  $v \in W^{2,2}(\Omega)$ ,  $T_j \in \mathcal{T}_R$  and  $E(v) := \sum_{\ell \in N_j} (v_\ell - v_j)$ ,*

$$|E(v)| \leq c h |v|_{W^{2,2}(\widetilde{T}_j)}, \quad (5.24)$$

where  $\widetilde{T}_j := T_j \cup \left( \bigcup_{\ell \in N_j} T_\ell \right)$ . For  $T_j \in \mathcal{T}_S$ , we immediately get by (5.19),

$$|E(v)| \leq c h |v|_{W^{2,2}(Q_j)} + c |v|_{W^{1,2}(Q_j)}. \quad (5.25)$$

Here,  $Q_j$  is a rectangular domain containing  $\widetilde{T}_j$  and a finite fixed number of neighboring triangles (cf. (5.19)).

*Proof:*

$$E(v) = E(v_j) = \sum_{\ell \in N_j} (v_\ell - v_j) = (v_\ell - v_j) + (v_k - v_j) + (v_m - v_j). \quad (5.26)$$

Consider now  $T := T_j \cup T_\ell \cup T_k \cup T_m$  and a reference triangle  $\widehat{T} = \widehat{T}_1 \cup \widehat{T}_2 \cup \widehat{T}_3 \cup \widehat{T}_4$ , such that  $\widehat{T}$  is mapped onto  $T$  by a 1-1 mapping

$$\begin{aligned} F : \widehat{T} &\rightarrow T, \\ F(\widehat{x}) &= A\widehat{x} + a \end{aligned}$$

and  $F(\widehat{T}_i) = T_{p_i}$  for  $i = 1, 2, 3, 4$ ,  $p_i = j, \ell, k, m$ , respectively. As in the proof of Lemma 5.4, c)  $A \in \mathbb{R}^{2 \times 2}$  is an invertible matrix and  $a \in \mathbb{R}^2$ . Define

$$\begin{aligned} \widehat{E} : W^{2,2}(\widehat{T}) &\rightarrow \mathbb{R}, \\ \widehat{E}(\widehat{v}) &= (\widehat{v}_\ell - \widehat{v}_j) + (\widehat{v}_k - \widehat{v}_j) + (\widehat{v}_m - \widehat{v}_j). \end{aligned}$$

Then

$$|\hat{E}(\hat{v})| \leq c \|\hat{v}\|_{C(\hat{T})} \leq c \|\hat{v}\|_{W^{2,2}(\hat{T})}.$$

Moreover,  $\hat{E}$  vanishes for all polynomials  $\hat{p}$  of degree  $\leq 1$ . Indeed, we have for such  $\hat{p}$  that

$$\hat{E}(\hat{p}) = \nabla \hat{p}[(\hat{x}_\ell - \hat{x}_j) + (\hat{x}_k - \hat{x}_j) + (\hat{x}_m - \hat{x}_j)] = 0.$$

Using then the Bramble–Hilbert lemma, one gets

$$|\hat{E}(\hat{v})| \leq c |\hat{v}|_{W^{2,2}(\hat{T})}.$$

Finally,

$$|E(v)| = |\hat{E}(\hat{v})| \leq c |\hat{v}|_{W^{2,2}(\hat{T})} \leq ch |v|_{W^{2,2}(T)}. \quad (5.27)$$

□

**6. The basic strategy in proving the main result.** The main technical step in the whole proof is to consider the term  $(L_h(I_h v) - L_h u_h, z_h) := \sum_j (L_h(I_h v) - L_h u_h)_j |T_j| z_j$  in the following form.

LEMMA 6.1.

$$(L_h(I_h v) - L_h u_h, z_h) = (\Psi_H, z_h) + (\Psi_K, z_h) + (\Psi_N, z_h), \quad (6.1)$$

where

$$\begin{aligned} \Psi_{Hj} &= -\frac{\varepsilon}{|T_j|} \sum_{\ell \in N_j} |S_{j\ell}| \left( \frac{v_\ell - v_j}{d_{j\ell}} - \frac{1}{|S_{j\ell}|} \int_{S_{j\ell}} \partial_n v \right), \\ \Psi_{Kj} &= \frac{1}{|T_j|} \sum_{\ell \in N_j} \left( g_{j\ell}(\mathcal{V}_{j\ell}, \mathcal{V}_{\ell j}) - \int_{S_{j\ell}} n_{j\ell} b v \right), \\ \Psi_{Nj} &= \frac{1}{|T_j|} \int_{T_j} c(v_j - v) \end{aligned}$$

where  $\mathcal{V}_{j\ell} = \mathcal{U}_{j\ell}(v_j, v_\ell)$ ,  $\mathcal{V}_{\ell j} = \mathcal{U}_{\ell j}(v_\ell, v_j)$ , and

$$(\Psi_A, z_h) := \sum_j \Psi_{Aj} z_j |T_j|, \quad \text{for } A = H, K, N.$$

*Proof:* We have due to (3.4) and (4.5),

$$(L_h I_h v)_j = -\frac{\varepsilon}{|T_j|} \sum_{\ell \in N_j} |S_{j\ell}| \left( \frac{v_\ell - v_j}{d_{j\ell}} \right) + \frac{1}{|T_j|} \sum_{\ell \in N_j} g_{j\ell}(\mathcal{V}_{j\ell}, \mathcal{V}_{\ell j}) + \frac{1}{|T_j|} \int_{T_j} c v_j.$$

Moreover,

$$(L_h u_h)_j = f_j = \frac{1}{|T_j|} \int_{T_j} f = \frac{1}{|T_j|} \int_{T_j} L v,$$

while (2.1) implies

$$\frac{1}{|T_j|} \int_{T_j} L v = -\frac{\varepsilon}{|T_j|} \sum_{\ell \in N_j} \int_{S_{j\ell}} \partial_n v + \frac{1}{|T_j|} \sum_{\ell \in N_j} \int_{S_{j\ell}} b v n_{j\ell} + \frac{1}{|T_j|} \int_{T_j} c v,$$

which completes the proof.  $\square$

In what follows, we split the sums in (6.1) into two parts,

$$\sum_j = \sum_{T_j \in \mathcal{T}_R} + \sum_{T_j \in \mathcal{T}_S}.$$

In the “regular” part of the sum we have regular triangles and the higher order approximation, in the “singular” part of the sum (“on the strips”) we have general triangles and the first order approximation. We thus get

$$(L_h(I_h v) - L_h u_h, z_h) = (\Psi_H, z_h)_R + (\Psi_H, z_h)_S + (\Psi_K, z_h)_R + (\Psi_K, z_h)_S + (\Psi_N, z_h)_R + (\Psi_N, z_h)_S$$

and will proceed by estimating the terms on the right-hand side both from above and from below.

## 7. The estimates from above.

**7.1. The estimate of  $(\Psi_N, z_h)$  from above.** For the estimate of  $(\Psi_N, z_h)$  (approximating the zero-order term) from above we obtain the following results.

LEMMA 7.1. *We have on the regular triangles*

$$\sum_{T_j \in \mathcal{T}_R} |\Psi_{Nj}|^2 |T_j| \leq ch^4 \|v\|_{2,2,R}^2, \quad (7.1)$$

$$(\Psi_N, z_h)_R \leq ch^4 \|v\|_{2,2,R}^2 + \frac{c_0}{8} \sum_{T_j \in \mathcal{T}_R} z_j^2 |T_j| \quad (7.2)$$

where  $\|v\|_{2,2,R}^2 := \sum_{T_j \in \mathcal{T}_R} \|v\|_{W^{2,2}(T_j)}^2$ .

*Proof:* Using [1, Lemma 4.3.8] we obtain that  $\int_{T_j} (v(\bar{x}_j) - v(x)) dx \leq ch^3 \|v\|_{W^{2,2}(T_j)}$  on the regular part  $\mathcal{T}_R$  of the grid, and therefore

$$\begin{aligned} \Psi_{Nj} &= \frac{1}{|T_j|} \int_{T_j} (v(\bar{x}_j) - v(x)) c(x) dx \\ &= \frac{1}{|T_j|} \left( c(\bar{x}_j) \int_{T_j} (v(\bar{x}_j) - v(x)) dx + \int_{T_j} (v(\bar{x}_j) - v(x)) (c(x) - c(\bar{x}_j)) dx \right) \\ &\leq ch \|v\|_{W^{2,2}(T_j)} + \sup_{x \in T_j} |v(\bar{x}_j) - v(x)| \frac{1}{|T_j|} \int_{T_j} |c(x) - c(\bar{x}_j)| dx \\ &\leq ch \|v\|_{W^{2,2}(Q_{j\ell})} \end{aligned}$$

using (5.19) and the smoothness of  $c(x)$ . Then (7.1) follows, since

$$\sum_{T_j \in \mathcal{T}_R} |\Psi_{Nj}|^2 |T_j| \leq c \sum_{T_j \in \mathcal{T}_R} h^2 \|v\|_{W^{2,2}(Q_{j\ell})}^2 |T_j| \leq ch^4 \|v\|_{2,2,R}^2.$$

Now this implies

$$\begin{aligned} (\Psi_N, z_h)_R &\leq \sum_{T_j \in \mathcal{T}_R} |\Psi_{Nj}| |z_j| |T_j| \leq c \sum_{T_j \in \mathcal{T}_R} |\Psi_{Nj}|^2 |T_j| + \frac{c_0}{8} \sum_{T_j \in \mathcal{T}_R} |T_j| |z_j|^2 \\ &\leq ch^4 \|v\|_{2,2,R}^2 + \frac{c_0}{8} \sum_{T_j \in \mathcal{T}_R} |T_j| |z_j|^2 \end{aligned}$$

which is (7.2).  $\square$

LEMMA 7.2. *We have on singular triangles*

$$\sum_{T_j \in \mathcal{T}_S} |\Psi_j|^2 |T_j| \leq ch^{4-2\delta} \|v\|_{2,2,S}^2 \quad (7.3)$$

$$(\Psi_N, z_h)_S \leq ch^{4-2\delta} \|v\|_{2,2,S}^2 + \frac{c_0}{8} \sum_{T_j \in \mathcal{T}_S} z_j^2 |T_j| \quad (7.4)$$

for any  $\delta > 0$ , where  $\|v\|_{2,2,S}^2 := \sum_{T_j \in \mathcal{T}_S} \|v\|_{W^{2,2}(T_j)}^2$ .

*Proof:* Similarly as in the proof of Lemma 7.1 we have

$$\Psi_{Nj} = \frac{1}{|T_j|} \left( c(x_j) \int_{T_j} (v(\bar{x}_j) - v(x)) dx + \int_{T_j} (v(x_j) - v(x)) (c(x) - c(\bar{x}_j)) dx \right).$$

The second term can be estimated as in the proof of Lemma 7.1. For the first one we use (5.22) and obtain

$$\begin{aligned} \Psi_{Nj} &= \frac{1}{|T_j|} \left( c(x_j) \int_{T_j} (v(\bar{x}_j) - v(x)) dx + \int_{T_j} (v(\bar{x}_j) - v(x)) (c(x) - c(\bar{x}_j)) dx \right) \\ &\leq ch^{1-\delta} \|v\|_{W^{2,2}(Q_{j\ell})}. \end{aligned}$$

This implies

$$\sum_S |\psi_{Nj}|^2 |T_j| \leq c \sum_S h^{2(1-\delta)} \|v\|_{W^{2,2}(Q_{j\ell})}^2 |T_j| \leq ch^{4-2\delta} \|v\|_{2,2,S}^2$$

and therefore, as before,

$$\begin{aligned} (\Psi_N, z_h)_S &\leq c \sum_{T_j \in \mathcal{T}_S} |\Psi_{Nj}|^2 |T_j| + \frac{c_0}{8} \sum_{T_j \in \mathcal{T}_S} |T_j| |z_j|^2 \\ &\leq ch^{4-2\delta} \|v\|_{2,2,S}^2 + \frac{c_0}{8} \sum_{T_j \in \mathcal{T}_S} |T_j| |z_j|^2. \end{aligned}$$

This finishes the proof of the lemma.  $\square$

Putting the results of Lemmata 7.1, 7.2 together, we obtain

LEMMA 7.3. *We have (on the whole domain)*

$$(\Psi_N, z_h) \leq ch^{4-2\delta} \|v\|_{2,2}^2 + \frac{c_0}{8} \sum_{T_j \in \mathcal{T}_h} z_j^2 |T_j|. \quad (7.5)$$

**7.2. The estimates of  $(\Psi_H, z_h)$  from above.** For the estimate of  $(\Psi_H, z_h)$  (approximating the diffusion part) from above we obtain the following result.

LEMMA 7.4. *For  $v \in W^{2,2}(\Omega)$  we have*

$$(\Psi_H, z_h) \leq c\varepsilon h \|v\|_{2,2} \left( \sum_{\mathcal{E} \cup \mathcal{E}_B} (z_j - z_\ell) \right)^{1/2} \quad (7.6)$$

$$\leq c\varepsilon h^2 \|v\|_{2,2}^2 + \frac{\varepsilon\gamma}{8} \sum_{\mathcal{E} \cup \mathcal{E}_B} (z_j - z_\ell)^2. \quad (7.7)$$

If moreover  $v \in W^{3,2}(\Omega)$ , we have

$$(\Psi_H, z_h) \leq c\varepsilon h^3 \|v\|_{3,2}^2 + \frac{\varepsilon\gamma}{8} \sum_{\mathcal{E} \cup \mathcal{E}_B} (z_j - z_\ell)^2. \quad (7.8)$$

*Proof:* We keep using the notation  $\kappa_{j\ell} := \kappa_{j\ell}(v) := \frac{v_\ell - v_j}{d_{j\ell}} - \frac{1}{|S_{j\ell}|} \int_{S_{j\ell}} \partial_n v$ , (cf. (5.20)). Note that  $\kappa_{j\ell} = -\kappa_{\ell j}$ . We have

$$(\Psi_H, z_h) = \sum_j \Psi_{Hj} z_j |T_j| = -\varepsilon \sum_j \sum_{N_{jI}} \kappa_{j\ell} |S_{j\ell}| z_j - \varepsilon \sum_j \sum_{N_{jG}} \kappa_{j\ell} |S_{j\ell}| z_j. \quad (7.9)$$

Now,

$$\begin{aligned} -\varepsilon \sum_j \sum_{N_{jI}} \kappa_{j\ell} |S_{j\ell}| z_j &= -\varepsilon \sum_{\mathcal{E}} |S_{j\ell}| (\kappa_{j\ell} z_j + \kappa_{\ell j} z_\ell) = \varepsilon \sum_{\mathcal{E}} |S_{j\ell}| \kappa_{j\ell} (z_\ell - z_j) \\ &\leq c\varepsilon h \left( \sum_{\mathcal{E}} |\kappa_{j\ell}|^2 \right)^{1/2} \left( \sum_{\mathcal{E}} (z_\ell - z_j)^2 \right)^{1/2} \\ &\stackrel{(5.20)}{\leq} c\varepsilon h \left( \sum_{\mathcal{E}} |v|_{W^{2,2}(Q_j)}^2 \right)^{1/2} \left( \sum_{\mathcal{E}} (z_\ell - z_j)^2 \right)^{1/2} \\ &\leq c\varepsilon h |v|_{W^{2,2}(\Omega)} \left( \sum_{\mathcal{E}} (z_\ell - z_j)^2 \right)^{1/2}, \\ -\varepsilon \sum_j \sum_{N_{jG}} \kappa_{j\ell} |S_{j\ell}| z_j &= -\frac{\varepsilon}{2} \sum_j \sum_{N_{jG}} \kappa_{j\ell} |S_{j\ell}| (z_j - z_\ell) = -\frac{\varepsilon}{2} \sum_{\mathcal{E}_B} \kappa_{j\ell} |S_{j\ell}| (z_j - z_\ell), \end{aligned} \quad (7.10)$$

the last-but-one equality holding due to (4.9). The last term can now be estimated as in (7.10) and (7.6) follows.

In order to prove (7.8) we consider  $(\Psi_H, z_h) = (\Psi_H, z_h)_R + (\Psi_H, z_h)_S$ . The second term can be estimated as in (7.10). For the first one we use (5.21) and proceed as in (7.10):

$$\begin{aligned} (\Psi_H, z_h) &= (\Psi_H, z_h)_R + (\Psi_H, z_h)_S \\ &\leq c\varepsilon h^4 \|v\|_{3,2}^2 + \frac{\varepsilon\gamma}{16} \sum_{\mathcal{E} \cup \mathcal{E}_B} (z_j - z_\ell)^2 + c\varepsilon h^2 \|v\|_{2,2,S}^2 + \frac{\varepsilon\gamma}{16} \sum_{\mathcal{E} \cup \mathcal{E}_B} (z_j - z_\ell)^2 \\ &\leq c\varepsilon h^4 \|v\|_{3,2}^2 + \frac{\varepsilon\gamma}{8} \sum_{\mathcal{E} \cup \mathcal{E}_B} (z_j - z_\ell)^2 + c\varepsilon h^3 \|v\|_{3,2,\Omega}^2, \end{aligned}$$

using (2.5), and the proof follows.  $\square$

**7.3. The estimates of  $(\Psi_K, z_h)$  from above.** For  $(\Psi_K, z_h)$  (approximating the convective term) we obtain the following lemma.

LEMMA 7.5. *We have*

$$(\Psi_K, z_h)_R \leq c \frac{h^3}{\varepsilon} \|v\|_{2,2}^2 + \frac{\varepsilon\gamma}{8} \sum_{\mathcal{E} \cup \mathcal{E}_B} (z_j - z_\ell)^2. \quad (7.11)$$

*Proof:* Considering first  $S_{j\ell} \in \mathcal{E}_R$ , we use that in this case  $g_{j\ell}(\mathcal{V}_{j\ell}, \mathcal{V}_{\ell j}) = g_{j\ell}(L_j^v(x_{j\ell}), L_\ell^v(x_{j\ell}))$  to write

$$\begin{aligned} g_{j\ell}(L_j^v(x_{j\ell}), L_\ell^v(x_{j\ell})) &- \int_{S_{j\ell}} n_{j\ell} b v \\ &= \int_{S_{j\ell}} (n_{j\ell} b)^+ (L_j^v(x_{j\ell}) - v) + \int_{S_{j\ell}} (n_{j\ell} b)^- (L_\ell^v(x_{j\ell}) - v) \\ &=: T + U. \end{aligned} \quad (7.12)$$

We consider  $T$  only, the term  $U$  can be treated similarly. We have

$$\begin{aligned} T &= \left(n_{j\ell}b(x_{j\ell})\right)^+ \int_{S_{j\ell}} (L_j^v(x_{j\ell}) - v) + \int_{S_{j\ell}} \left[ (n_{j\ell}b)^+ - \left(n_{j\ell}b(x_{j\ell})\right)^+ \right] (L_j^v(x_{j\ell}) - v) \\ &=: T_1 + T_2, \end{aligned} \quad (7.13)$$

with  $L_j^v(x_{j\ell}) = v_j + \alpha_j^v G_j^v(x_{j\ell} - x_j)$ .

First let us consider the case  $\alpha_j^v = 1$ . Estimating  $T_1$ , we will consider two cases. Firstly, if the stencil of  $G_j^v$  contains  $\ell$ , we get

$$L_j^v(x_{j\ell}) - v = v_j + (v_{j\ell} - v_j) - v = v_{j\ell} - v.$$

Then,

$$\int_{S_{j\ell}} (v_{j\ell} - v) d\sigma(x) = \int_S (\hat{v}_{j\ell} - \hat{v}(\hat{x})) \frac{|A\hat{\xi}|}{|\hat{\xi}|} d\sigma(\hat{x}) =: H(\hat{v})$$

for  $\hat{v} \in W^{2,2}(\hat{T})$ , using the same mapping  $F : \hat{T} \rightarrow T$  as in the proof of Lemma 5.4. It follows that

$$H(\hat{p}) = 0 \quad \text{for all polynomials } \hat{p} \text{ of degree } \leq 1$$

and

$$|H(\hat{v})| \leq ch \|\hat{v}\|_{C(\hat{T})} \leq ch \|\hat{v}\|_{W^{2,2}(\hat{T})}.$$

Therefore, using the Bramble–Hilbert lemma,

$$|H(\hat{v})| \leq ch |\hat{v}|_{W^{2,2}(\hat{T})} \leq ch^2 |v|_{W^{2,2}(T)}.$$

Consequently,

$$|T_1| \leq ch^2 |v|_{W^{2,2}(T)}.$$

Secondly, but still for  $\alpha_j^v = 1$ , if the stencil of  $G_j^v$  does not contain  $\ell$ , we have

$$\begin{aligned} L_j^v(x_{j\ell}) - v &= v_j - G_j^v\left((x_{jk} - x_j) + (x_{jm} - x_j)\right) - v \\ &= (v_j - v_{jk}) + (v_j - v_{jm}) + (v_j - v). \end{aligned}$$

Using the same transformation as before, we can show that

$$H(\hat{v}) := \int_S \left( (\hat{v}_j - \hat{v}_{jk}) + (\hat{v}_j - \hat{v}_{jm}) + (\hat{v}_j - \hat{v}(\hat{x})) \right) \frac{|A\hat{\xi}|}{|\hat{\xi}|} d\sigma(\hat{x})$$

satisfies

$$|H(\hat{v})| \leq ch \|\hat{v}\|_{W^{2,2}(\hat{T})},$$

while for all polynomials  $\hat{p}$  of degree  $\leq 1$ ,

$$\begin{aligned} H(\hat{p}) &= \int_{S_{j\ell}} (\nabla p(x_j - x_{jk}) + \nabla p(x_j - x_{jm}) + p(x_j) - p(x)) d\sigma(x) \\ &= \int_{S_{j\ell}} (p(x_{j\ell}) - p(x)) d\sigma(x) = 0. \end{aligned}$$

Therefore, by the same argument as before,

$$|T_1| \leq ch^2 |v|_{W^{2,2}(T)}.$$

Finally, for general  $\alpha_j^v$  we obtain

$$\begin{aligned}
T_1 &= \left(n_{j\ell}b(x_{j\ell})\right)^+ \int_{S_{j\ell}} (L_j^v(x_{j\ell}) - v) \\
&= \left(n_{j\ell}b(x_{j\ell})\right)^+ \int_{S_{j\ell}} (v_j + \alpha_j^v G_j^v(x_{j\ell} - x_j) - v) \\
&\leq \left(n_{j\ell}b(x_{j\ell})\right)^+ \int_{S_{j\ell}} (v_j + G_j^v(x_{j\ell} - x_j) - v) + \left(n_{j\ell}b(x_{j\ell})\right)^+ \int_{S_{j\ell}} |\alpha_j^v - 1| |G_j^v(x_{j\ell} - x_j)|.
\end{aligned} \tag{7.14}$$

The integrand of the first integral corresponds to the case of  $\alpha_j^v = 1$  studied above and therefore the first integral can be estimated by  $ch^2|v|_{W^{2,2}(T)}$ . The second integral can be estimated as follows using (3.21):

$$\int_{S_{j\ell}} |\alpha_j^v - 1| |G_j^v(x_{j\ell} - x_j)| \leq \frac{1}{2} \int_{S_{j\ell}} |\alpha_j^v - 1| |v_\ell - v_j| \leq \frac{1}{2} \int_{S_{j\ell}} |\alpha_j^v - 1| \left| \sum_{l \in N_j} (v_\ell - v_j) \right|$$

since  $\alpha_j^v - 1 \neq 0$  implies that there is a local extremum in  $v_j$ . We then continue

$$\int_{S_{j\ell}} |\alpha_j^v - 1| |G_j^v(x_{j\ell} - x_j)| \leq c \int_{S_{j\ell}} |E(v_j)| \leq ch^2|v|_{W^{2,2}(\tilde{T}_j)}$$

where we used (5.24). We conclude that in any case we have using (3.20)

$$|T_1| \leq ch^2 \|v\|_{W^{2,2}(\tilde{T}_j)}.$$

The term  $T_2$  in (7.13) can be estimated as follows:

$$|T_2| \leq c|S_{j\ell}|h \sup_{S_{j\ell}} |L_j^v(x_{j\ell}) - v| \leq ch^2 \|v\|_{W^{2,2}(T)}. \tag{7.15}$$

The term  $U$  in (7.12) can be treated in the same way. This finishes the first part of the proof.

Now we will consider the general situation for any  $S_{j\ell}$ . We have

$$\begin{aligned}
(\Psi_K, z_h) &= \sum_j \sum_{N_j} (g_{j\ell}(\mathcal{V}_{j\ell}, \mathcal{V}_{\ell j}) - \int_{S_{j\ell}} n_{j\ell}bv) z_j \\
&= \sum_{\mathcal{T}_R} \sum_{N_j} \cdots + \sum_{\mathcal{T}_S} \sum_{N_j} \cdots \\
&= \sum_{\mathcal{T}_R} \sum_{N_{jR}} \cdots + \sum_{\mathcal{T}_R} \sum_{N_{jS}} \cdots + \sum_{\mathcal{T}_R} \sum_{N_{jG}} \cdots \\
&\quad + \sum_{\mathcal{T}_S} \sum_{N_{jR}} \cdots + \sum_{\mathcal{T}_S} \sum_{N_{jS}} \cdots + \sum_{\mathcal{T}_S} \sum_{N_{jG}} \cdots.
\end{aligned} \tag{7.16}$$

Using the first part of the proof, we get for the first case

$$\begin{aligned}
\sum_{\mathcal{T}_R} \sum_{N_{jR}} \cdots &= \sum_{\mathcal{T}_R} \sum_{N_{jR}} (g_{j\ell}(L_j^v(x_{j\ell}), L_\ell^v(x_{j\ell})) - \int_{S_{j\ell}} n_{j\ell}bv) z_j \\
&= \sum_{\mathcal{E}_R} (g_{j\ell}(L_j^v(x_{j\ell}), L_\ell^v(x_{j\ell})) - \int_{S_{j\ell}} n_{j\ell}bv) (z_j - z_\ell) \\
&\leq c \sum_{\mathcal{E}_R} h^2 \|v\|_{W^{2,2}(T_{j\ell})} (z_j - z_\ell) \\
&\leq c \frac{h^4}{\varepsilon} \|v\|_{W^{2,2}(\Omega)}^2 + \frac{\varepsilon\gamma}{8} \sum_{\mathcal{E}_R} (z_j - z_\ell)^2.
\end{aligned} \tag{7.17}$$

Secondly,

$$\begin{aligned}
 \sum_{\mathcal{T}_R} \sum_{N_{jS}} \cdots + \sum_{\mathcal{T}_S} \sum_{N_{jR}} \cdots &= \sum_{\mathcal{T}_R} \sum_{N_{jS}} \left( g_{j\ell}(v_j, v_\ell) - \int_{S_{j\ell}} n_{j\ell} b v \right) z_j \\
 &+ \sum_{\mathcal{T}_S} \sum_{N_{jR}} \left( g_{j\ell}(v_j, v_\ell) - \int_{S_{j\ell}} n_{j\ell} b v \right) z_j \\
 &= \sum_{\mathcal{E}_M} \left( g_{j\ell}(v_j, v_\ell) - \int_{S_{j\ell}} n_{j\ell} b v \right) (z_j - z_\ell)
 \end{aligned} \tag{7.18}$$

Since  $g_{j\ell}(v_j, v_\ell) = b_{j\ell}^+ v_j + b_{j\ell}^- v_\ell$  for  $S_{j\ell} \in \mathcal{E}_M$ , we proceed by estimating

$$\begin{aligned}
 &\left| b_{j\ell}^+ v_j + b_{j\ell}^- v_\ell - \int_{S_{j\ell}} n_{j\ell} b v \right| \\
 &\leq \left| \int_{S_{j\ell}} (bn_{j\ell})^+ \underbrace{(v_j - v)}_{\text{use (5.19)}} + (bn_{j\ell})^- \underbrace{(v_\ell - v)}_{\text{use (5.19)}} \right| \\
 &\leq c \int_{S_{j\ell}} |bn_{j\ell}| (h \|v\|_{W^{2,2}(Q_{j\ell})} + \|v\|_{W^{1,2}(Q_{j\ell})}) \\
 &\leq c h^2 \|v\|_{W^{2,2}(Q_{j\ell})} + c h \|v\|_{W^{1,2}(Q_{j\ell})},
 \end{aligned} \tag{7.19}$$

and continue in (7.18) to obtain

$$\begin{aligned}
 &\sum_{\mathcal{T}_R} \sum_{N_{jS}} \cdots + \sum_{\mathcal{T}_S} \sum_{N_{jR}} \cdots \\
 &\leq c \sum_{\mathcal{E}_M} h^2 \|v\|_{W^{2,2}(Q_{j\ell})} (z_j - z_\ell) + c \sum_{\mathcal{E}_M} h \|v\|_{W^{1,2}(Q_{j\ell})} (z_j - z_\ell)
 \end{aligned} \tag{7.20}$$

$$\leq c \frac{h^4}{\varepsilon} \|v\|_{W^{2,2}(\mathcal{T}_S)}^2 + c \frac{h^2}{\varepsilon} \|v\|_{W^{1,2}(\mathcal{T}_S)}^2 + \frac{\varepsilon \gamma}{8} \sum_{\mathcal{E}_M} (z_j - z_\ell)^2, \tag{7.21}$$

and, since according to (2.5) we have  $\|v\|_{W^{1,2}(\mathcal{T}_S)}^2 \leq ch \|v\|_{W^{2,2}(\Omega)}^2$ , we obtain in this case finally

$$\sum_{\mathcal{T}_R} \sum_{N_{jS}} \cdots + \sum_{\mathcal{T}_S} \sum_{N_{jR}} \cdots \leq c \frac{h^3}{\varepsilon} \|v\|_{2,2}^2 + \frac{\varepsilon \gamma}{8} \sum_{\mathcal{E}_M} (z_j - z_\ell)^2. \tag{7.22}$$

Thirdly, we consider

$$\begin{aligned}
 \sum_{\mathcal{T}_S} \sum_{N_{jS}} \cdots &= \sum_{\mathcal{T}_S} \sum_{N_{jS}} \left( g_{j\ell}(v_j, v_\ell) - \int_{S_{j\ell}} n_{j\ell} b v \right) z_j \\
 &= \sum_{\mathcal{E}_S} \left( g_{j\ell}(v_j, v_\ell) - \int_{S_{j\ell}} n_{j\ell} b v \right) (z_j - z_\ell)
 \end{aligned} \tag{7.23}$$

and proceed as in (7.19), (7.22) to get

$$\sum_{\mathcal{T}_S} \sum_{N_{jS}} \cdots \leq c \frac{h^3}{\varepsilon} \|v\|_{2,2}^2 + \frac{\varepsilon \gamma}{8} \sum_{\mathcal{E}_S} (z_j - z_\ell)^2. \tag{7.24}$$

Finally we will estimate the terms with the ghost cells. We have

$$\begin{aligned} \sum_{\mathcal{T}_R} \sum_{N_{jG}} \cdots &= \sum_{\mathcal{T}_R} \sum_{N_{jG}} \left( g_{j\ell}(v_j, v_\ell) - \int_{S_{j\ell}} n_{j\ell} b v \right) z_j \\ &= \frac{1}{2} \sum_{\mathcal{T}_R} \sum_{N_{jG}} \left( g_{j\ell}(v_j, v_\ell) - \int_{S_{j\ell}} n_{j\ell} b v \right) (z_j - z_\ell) \end{aligned}$$

and therefore we can proceed as before to get  $c \frac{h^3}{\varepsilon} \|v\|_{2,2}^2 + \frac{\varepsilon\gamma}{8} \sum_{\mathcal{E}_B} (z_j - z_\ell)^2$ . The other terms in (7.16) can be treated similarly, and the result follows.  $\square$

**7.4. The final estimate from above.** Putting together the estimates (7.5), (7.7), (7.8), (7.11), we get the following result.

**THEOREM 7.6. (Estimate from above)** *Under the assumptions of Theorem 4.1 there exists a constant  $c > 0$  independent of  $\varepsilon$  such that for  $v \in W^{2,2}(\Omega)$  we have*

$$\begin{aligned} (L_h I_h v - L_h u_h, z_h) &\leq c \left( \varepsilon h^2 + h^{4-2\delta} + \frac{h^3}{\varepsilon} \right) \|v\|_{2,2}^2 \\ &\quad + \frac{\varepsilon\gamma}{4} \sum_{\mathcal{E} \cup \mathcal{E}_B} (z_j - z_\ell)^2 + \frac{c_0}{8} \sum_{T_j \in \mathcal{T}_h} z_j^2 |T_j|. \end{aligned} \quad (7.25)$$

If moreover  $v \in W^{3,2}(\Omega)$ , we have

$$\begin{aligned} (L_h I_h v - L_h u_h, z_h) &\leq c \left( \varepsilon h^3 + h^{4-2\delta} + \frac{h^3}{\varepsilon} \right) \|v\|_{3,2}^2 \\ &\quad + \frac{\varepsilon\gamma}{4} \sum_{\mathcal{E} \cup \mathcal{E}_B} (z_j - z_\ell)^2 + \frac{c_0}{8} \sum_{T_j \in \mathcal{T}_h} z_j^2 |T_j|. \end{aligned} \quad (7.26)$$

**8. The estimates from below.** In this part of the paper we will prove an estimate from below.

**THEOREM 8.1. (Estimate from below)** *Under the assumptions of Theorem 4.1 there exists a constant  $c > 0$  independent of  $\varepsilon$  such that for  $v \in W^{2,2}(\Omega)$  we have*

$$\begin{aligned} &(L_h I_h v - L_h u_h, z_h) \\ &\geq \frac{\varepsilon\gamma}{2} \sum_{\mathcal{E}} (z_\ell - z_j)^2 + 2\varepsilon\gamma \sum_{\mathcal{E}_B} z_j^2 + c_0 \sum_j z_j^2 |T_j| + \frac{1}{2} \sum_{\mathcal{E}_M \cup \mathcal{E}_S} (b_{j\ell}^+ - b_{j\ell}^-) (z_j - z_\ell)^2 \\ &\quad - c \frac{h^4}{\varepsilon} \|v\|_{W^{2,2}(\Omega)}^2 - c \frac{h^4}{\varepsilon} \sum_{\mathcal{T}_R} R_j^2 |T_j| \end{aligned} \quad (8.1)$$

where  $\gamma = \min \gamma_{j\ell}$  and  $R_j := \frac{1}{|T_j|} \sum_{\ell \in N_j} (u_\ell - u_j) \gamma_{j\ell}$ .

*Proof:* Using the notation  $\sum_j \equiv \sum_{T_j \in \mathcal{T}_h}$  and the definition of  $L_h$  one can write

$$(L_h I_h v - L_h u_h, z_h) = A_1 + A_2 + A_3$$

where, denoting as before  $\mathcal{V}_{j\ell} = \mathcal{U}_{j\ell}(v_j, v_\ell)$ ,  $\mathcal{V}_{\ell j} = \mathcal{U}_{\ell j}(v_\ell, v_j)$ ,

$$\begin{aligned} A_1 &= \sum_j \left( -\varepsilon \sum_{\ell \in N_j} (v_\ell - v_j) \gamma_{j\ell} + \varepsilon \sum_{\ell \in N_j} (u_\ell - u_j) \gamma_{j\ell} \right) z_j \geq \varepsilon\gamma \sum_{\mathcal{E}} (z_\ell - z_j)^2 + 2\varepsilon\gamma \sum_{\mathcal{E}_B} z_j^2, \\ A_2 &= \sum_j \sum_{\ell \in N_j} \left( g_{j\ell}(\mathcal{V}_{j\ell}, \mathcal{V}_{\ell j}) - g_{j\ell}(\mathcal{U}_{j\ell}, \mathcal{U}_{\ell j}) \right) z_j, \\ A_3 &= \sum_j c_j (v_j - u_j) z_j |T_j| \geq c_0 \sum_j z_j^2 |T_j|. \end{aligned} \quad (8.2)$$

It remains to treat  $A_2$ . Recall that

$$g_{j\ell}(\mathcal{U}_{j\ell}, \mathcal{U}_{\ell j}) = \begin{cases} b_{j\ell}^+ L_j^u(x_{j\ell}) + b_{j\ell}^- L_\ell^u(x_{j\ell}) & \text{for } S_{j\ell} \in \mathcal{E}_R, \\ b_{j\ell}^+ u_j + b_{j\ell}^- u_\ell & \text{for } S_{j\ell} \in \mathcal{E}_S \cup \mathcal{E}_M \cup \mathcal{E}_B. \end{cases} \quad (8.3)$$

For  $S_{j\ell} \in \mathcal{E}_R$  we can use the fact that

$$L_j^u(x_{j\ell}) - u_j = \frac{1}{2}(L_j^u(x_\ell) - u_j) = \frac{1}{2}(L_j^u(x_\ell) - u_\ell) + \frac{1}{2}(u_\ell - u_j)$$

(and similarly for  $L_\ell^u(x_{j\ell}) - u_\ell$ ), and write

$$\begin{aligned} g_{j\ell}(\mathcal{U}_{j\ell}, \mathcal{U}_{\ell j}) &= b_{j\ell}^+ u_j + b_{j\ell}^- u_\ell + b_{j\ell}^+ (L_j^u(x_{j\ell}) - u_j) + b_{j\ell}^- (L_\ell^u(x_{j\ell}) - u_\ell) \\ &= b_{j\ell}^+ u_j + b_{j\ell}^- u_\ell + \frac{1}{2} b_{j\ell}^+ (L_j^u(x_\ell) - u_\ell) + \frac{1}{2} b_{j\ell}^+ (u_\ell - u_j) \\ &\quad + \frac{1}{2} b_{j\ell}^- (L_\ell^u(x_j) - u_j) + \frac{1}{2} b_{j\ell}^- (u_j - u_\ell) \end{aligned}$$

and similarly for  $g_{j\ell}(\mathcal{V}_{j\ell}, \mathcal{V}_{\ell j})$ . Therefore,

$$\begin{aligned} A_2 &= \sum_j \sum_{\ell \in N_j} \left( g_{j\ell}(\mathcal{V}_{j\ell}, \mathcal{V}_{\ell j}) - g_{j\ell}(\mathcal{U}_{j\ell}, \mathcal{U}_{\ell j}) \right) z_j \\ &= \sum_j \sum_{\ell \in N_j} (b_{j\ell}^+ v_j + b_{j\ell}^- v_\ell) z_j + \frac{1}{2} \sum_{\mathcal{T}_R} \sum_{N_{jR}} \left( b_{j\ell}^+ (L_j^v(x_\ell) - v_\ell) + b_{j\ell}^- (L_\ell^v(x_j) - v_j) \right) z_j \\ &\quad + \frac{1}{2} \sum_{\mathcal{T}_R} \sum_{N_{jR}} \left( b_{j\ell}^+ (v_\ell - v_j) + b_{j\ell}^- (v_j - v_\ell) \right) z_j \\ &\quad - \sum_j \sum_{\ell \in N_j} (b_{j\ell}^+ u_j + b_{j\ell}^- u_\ell) z_j - \frac{1}{2} \sum_{\mathcal{T}_R} \sum_{N_{jR}} \left( b_{j\ell}^+ (L_j^u(x_\ell) - u_\ell) + b_{j\ell}^- (L_\ell^u(x_j) - u_j) \right) z_j \\ &\quad - \frac{1}{2} \sum_{\mathcal{T}_R} \sum_{N_{jR}} \left( b_{j\ell}^+ (u_\ell - u_j) + b_{j\ell}^- (u_j - u_\ell) \right) z_j, \end{aligned}$$

and hence,

$$\begin{aligned} A_2 &= \sum_j \sum_{\ell \in N_j} (b_{j\ell}^+ z_j + b_{j\ell}^- z_\ell) z_j \\ &\quad + \frac{1}{2} \sum_{\mathcal{T}_R} \sum_{N_{jR}} \left( b_{j\ell}^+ (L_j^v(x_\ell) - v_\ell) + b_{j\ell}^- (L_\ell^v(x_j) - v_j) \right) z_j \\ &\quad - \frac{1}{2} \sum_{\mathcal{T}_R} \sum_{N_{jR}} \left( b_{j\ell}^+ (L_j^u(x_\ell) - u_\ell) + b_{j\ell}^- (L_\ell^u(x_j) - u_j) \right) z_j \\ &\quad + \frac{1}{2} \sum_{\mathcal{T}_R} \sum_{N_{jR}} \left( b_{j\ell}^+ (z_\ell - z_j) + b_{j\ell}^- (z_j - z_\ell) \right) z_j \\ &=: W_1 + W_2 + W_3 + W_4. \end{aligned}$$

For  $W_1$  we proceed by the same way as we did treating the term  $B_1$  on page 12 to get (cf. (5.17))

$$W_1 \geq \frac{1}{2} \sum_{\mathcal{E}} (b_{j\ell}^+ - b_{j\ell}^-) (z_j - z_\ell)^2, \quad (8.4)$$

while for  $W_4$  we get

$$\begin{aligned} W_4 &= \frac{1}{2} \sum_{T_R} \sum_{N_{jR}} \left( (b_{j\ell}^+ - b_{j\ell}^-)(z_\ell - z_j) \right) z_j = \frac{1}{2} \sum_{\mathcal{E}_R} (b_{j\ell}^+ - b_{j\ell}^-) \left( (z_\ell - z_j)z_j + (z_j - z_\ell)z_\ell \right) \\ &= -\frac{1}{2} \sum_{\mathcal{E}_R} (b_{j\ell}^+ - b_{j\ell}^-)(z_j - z_\ell)^2. \end{aligned}$$

Therefore,

$$W_1 + W_4 \geq \frac{1}{2} \sum_{\mathcal{E}_S \cup \mathcal{E}_M} (b_{j\ell}^+ - b_{j\ell}^-)(z_j - z_\ell)^2. \quad (8.5)$$

In order to estimate terms  $W_2$  and  $W_3$  we use the definition (3.18) to see that

$$u_\ell - L_j^u(x_\ell) = u_\ell - u_j - \alpha_j^u G_j^u(x_\ell - x_j).$$

Let us discuss the three possible cases. If  $\alpha_j^u = 1$  and stencil of  $G_j^u$  contains  $x_\ell$  then  $u_\ell - L_j^u(x_\ell) = 0$ . If  $\alpha_j^u = 1$  and stencil of  $G_j^u$  does not contain  $x_\ell$ , we have

$$\begin{aligned} u_\ell - L_j^u(x_\ell) &= u_\ell - u_j - G_j^u(x_\ell - x_j) \\ &= u_\ell - u_j + G_j^u(x_m - x_j + x_k - x_j) \\ &= u_\ell - u_j + u_m - u_j + u_k - u_j \\ &= \left( \frac{1}{|T_j|} \sum_{k \in N_j} (u_k - u_j) \gamma_{jk} \right) \frac{|T_j|}{\gamma_j} =: R_j \frac{|T_j|}{\gamma_j}. \end{aligned} \quad (8.6)$$

In the (8.6) we take the advantage of the fact that for  $T_j \in \mathcal{T}_R$  we have  $\gamma_{jk} = \text{const}_j =: \gamma_j$  for all  $k \in N_j$ .

Finally, if  $\alpha_j^u = 0$ , then  $u_\ell - L_j^u(x_\ell) = u_\ell - u_j$ . However,  $\alpha_j^u = 0$  implies that there is local extremum in  $u_j$ , or more precisely, that  $u_j$  is extremal out of the values  $u_j, u_\ell, \ell \in N_j$ . Therefore we have in this case

$$|u_\ell - u_j| \leq \left| \sum_{k \in N_j} (u_k - u_j) \right| = |R_j| \frac{|T_j|}{\gamma_j}.$$

We conclude that in any case,

$$|u_\ell - L_j^u(x_\ell)| \leq |R_j| \frac{|T_j|}{\gamma_j} \quad (8.7)$$

and the same estimate holds when replacing  $j$  by  $\ell$  and vice versa.

By the same considerations we get that

$$|v_\ell - L_j^v(x_\ell)| \leq \left| \sum_{\ell \in N_j} (v_\ell - v_j) \right|. \quad (8.8)$$

In order to estimate the last term, we define

$$E(v) = E(v_j) := \sum_{\ell \in N_j} (v_\ell - v_j) = (v_\ell - v_j) + (v_k - v_j) + (v_m - v_j). \quad (8.9)$$

Consider now  $T := T_j \cup T_\ell \cup T_k \cup T_m$  and a reference triangle  $\hat{T} = \hat{T}_1 \cup \hat{T}_2 \cup \hat{T}_3 \cup \hat{T}_4$ , such that  $\hat{T}$  is mapped onto  $T$  by a 1-1 mapping

$$\begin{aligned} F : \hat{T} &\rightarrow T, \\ F(\hat{x}) &= A\hat{x} + a \end{aligned}$$

and  $F(\hat{T}_i) = T_{p_i}$  for  $i = 1, 2, 3, 4$ ,  $p_i = j, \ell, k, m$ , respectively. Note that  $\hat{T}$  is also a regular, larger triangle. As in the proof of Lemma 5.4 part c),  $A \in \mathbb{R}^{2 \times 2}$  is an invertible matrix and  $a \in \mathbb{R}^2$ . Define

$$\begin{aligned} \hat{E} &: W^{2,2}(\hat{T}) \rightarrow \mathbb{R}, \\ \hat{E}(\hat{v}) &= (\hat{v}_\ell - \hat{v}_j) + (\hat{v}_k - \hat{v}_j) + (\hat{v}_m - \hat{v}_j). \end{aligned}$$

Then

$$|\hat{E}(\hat{v})| \leq c \|\hat{v}\|_{C(\hat{T})} \leq c \|\hat{v}\|_{W^{2,2}(\hat{T})}.$$

Moreover,  $\hat{E}$  vanishes for all polynomials  $\hat{p}$  of degree  $\leq 1$ . Indeed, we have for such  $\hat{p}$  that

$$\hat{E}(\hat{p}) = \nabla \hat{p}[(\hat{x}_\ell - \hat{x}_j) + (\hat{x}_k - \hat{x}_j) + (\hat{x}_m - \hat{x}_j)] = 0.$$

Using then the Bramble–Hilbert lemma, one gets

$$|\hat{E}(\hat{v})| \leq c |\hat{v}|_{W^{2,2}(\hat{T})}.$$

Finally,

$$|E(v)| = |\hat{E}(\hat{v})| \leq c |\hat{v}|_{W^{2,2}(\hat{T})} \leq ch |v|_{W^{2,2}(T)} \quad (8.10)$$

using the properties of  $F$  similar to those in the proof of Lemma 5.4, part c). Therefore we obtain using (8.8), (5.27),

$$|v_\ell - L_j^v(x_\ell)| \leq ch |v|_{W^{2,2}(T)} \quad (8.11)$$

with  $c > 0$  independent of  $h$  and  $T = T_j \cup \bigcup_{j \in N_\ell} T_\ell$ , and the same estimate holds for  $|v_j - L_\ell^v(x_j)|$ .

Using now (8.11) we see that

$$\begin{aligned} W_2 &= \frac{1}{2} \sum_{T_R} \sum_{N_{jR}} \left( b_{j\ell}^+ (L_j^v(x_\ell) - v_\ell) + b_{j\ell}^- (L_\ell^v(x_j) - v_j) \right) z_j \\ &= \frac{1}{2} \sum_{\mathcal{E}_R} \left( b_{j\ell}^+ (L_j^v(x_\ell) - v_\ell) + b_{j\ell}^- (L_\ell^v(x_j) - v_j) \right) (z_j - z_\ell) \\ &\geq -\frac{1}{2} \sum_{\mathcal{E}_R} \left( b_{j\ell}^+ |L_j^v(x_\ell) - v_\ell| |z_j - z_\ell| + |b_{j\ell}^-| |L_\ell^v(x_j) - v_j| |z_j - z_\ell| \right) \\ &\geq -c h^2 \sum_{\mathcal{E}_R} |v|_{W^{2,2}(T)} |z_j - z_\ell| \\ &\geq -c \frac{h^4}{\varepsilon} \sum_{\mathcal{E}_R} |v|_{W^{2,2}(T)}^2 - \frac{\varepsilon \gamma}{4} \sum_{\mathcal{E}_R} (z_j - z_\ell)^2 \\ &\geq -c \frac{h^4}{\varepsilon} \|v\|_{W^{2,2}(\Omega)}^2 - \frac{\varepsilon \gamma}{4} \sum_{\mathcal{E}_R} (z_j - z_\ell)^2. \end{aligned} \quad (8.12)$$

Similarly, (8.7) implies that

$$\begin{aligned} W_3 &\geq -\frac{1}{2} \sum_{\mathcal{E}_R} \left( b_{j\ell}^+ |L_j^u(x_\ell) - u_\ell| |z_j - z_\ell| + |b_{j\ell}^-| |L_\ell^u(x_j) - u_j| |z_j - z_\ell| \right) \\ &\geq -c h^2 \sum_{\mathcal{E}_R} \left( |R_j| \sqrt{|T_j|} + |R_\ell| \sqrt{|T_\ell|} \right) |z_j - z_\ell| \\ &\geq -c \frac{h^4}{\varepsilon} \sum_{T_R} R_j^2 |T_j| - \frac{\varepsilon \gamma}{4} \sum_{\mathcal{E}_R} (z_j - z_\ell)^2. \end{aligned} \quad (8.13)$$

Now, estimates (8.5), (8.12), (8.13) imply

$$A_2 \geq \frac{1}{2} \sum_{\mathcal{E}_S \cup \mathcal{E}_M} (b_{j\ell}^+ - b_{j\ell}^-)(z_j - z_\ell)^2 - \frac{\varepsilon\gamma}{2} \sum_{\mathcal{E}_R} (z_j - z_\ell)^2 - c \frac{h^4}{\varepsilon} \|v\|_{W^{2,2}(\Omega)}^2 - c \frac{h^4}{\varepsilon} \sum_{\mathcal{T}_R} R_j^2 |T_j|,$$

which together with (8.2) finishes the proof.  $\square$

LEMMA 8.2. *Let  $R_j = \frac{1}{|T_j|} \sum_{\ell \in N_j} (u_\ell - u_j) \gamma_{j\ell}$  for  $T_j \in \mathcal{T}_R$  (cf. (8.6)). Then there is a constant  $c > 0$  independent of  $\varepsilon$  and  $h$ , such that*

$$\sum_{\mathcal{T}_R} R_j^2 |T_j| \leq \frac{c}{\varepsilon^3} \|f\|_{L^2(\Omega)}^2. \quad (8.14)$$

*Proof:* From the definition of the numerical scheme in (3.4) we obtain

$$\begin{aligned} R_j &= \frac{1}{\varepsilon |T_j|} \left( \varepsilon \sum_{\ell \in N_j} (u_\ell - u_j) \gamma_{j\ell} \right) \\ &\quad \text{(see (3.4) and use } (L_h u_h)_j = f_j \text{ with } T_j \in \mathcal{T}_R) \\ &= \frac{1}{\varepsilon |T_j|} \left( \sum_{\ell \in N_j} g_{j\ell} (L_j^u(x_{j\ell}), L_\ell^u(x_{j\ell})) + c_j u_j |T_j| - f_j |T_j| \right). \end{aligned} \quad (8.15)$$

We have for  $T_j \in \mathcal{T}_R$ ,

$$\begin{aligned} &\left| \sum_{\ell \in N_j} g_{j\ell} (L_j^u(x_{j\ell}), L_\ell^u(x_{j\ell})) \right| = \left| \sum_{\ell \in N_j} [b_{j\ell}^+ (L_j^u(x_{j\ell})) + b_{j\ell}^- (L_\ell^u(x_{j\ell}))] \right| \\ &= \left| \sum_{\ell \in N_j} [b_{j\ell}^+ (L_j^u(x_{j\ell}) - u_j) + b_{j\ell}^- (L_\ell^u(x_{j\ell}) - u_\ell) + u_j b_{j\ell}^+ + u_\ell b_{j\ell}^-] \right| \\ &\quad \text{using (3.20)} \\ &\leq ch \sum_{\ell \in N_j} |u_\ell - u_j| + \left| \sum_{\ell \in N_j} (b_{j\ell}^+ + b_{j\ell}^-) u_j + b_{j\ell}^- (u_\ell - u_j) \right| \\ &\leq ch \sum_{\ell \in N_j} |u_\ell - u_j|, \end{aligned} \quad (8.16)$$

since  $\sum_{\ell \in N_j} (b_{j\ell}^+ + b_{j\ell}^-) u_j = 0$ . Therefore

$$\begin{aligned} |R_j| &\leq \frac{c}{\varepsilon h^2} \left( h \sum_{\ell \in N_j} |u_\ell - u_j| + |u_j| |T_j| + |f_j| |T_j| \right) \\ &\leq \frac{c}{\varepsilon} \left( \frac{1}{h} \sum_{\ell \in N_j} |u_\ell - u_j| + |u_j| + |f_j| \right) \end{aligned} \quad (8.17)$$

and

$$\begin{aligned} \sum_{T_j \in \mathcal{T}_R} R_j^2 |T_j| &\leq \frac{c}{\varepsilon^2} \sum_{T_j \in \mathcal{T}_R} \left( \sum_{N_{jR}} |u_\ell - u_j|^2 + u_j^2 |T_j| + f_j^2 |T_j| \right) \\ &\leq \frac{c}{\varepsilon^3} \left( \varepsilon \sum_{\mathcal{E}_R} |u_\ell - u_j|^2 \right) + \frac{c}{\varepsilon^2} \sum_j \left( u_j^2 |T_j| + f_j^2 |T_j| \right) \\ &\leq \frac{c}{\varepsilon^3} \sum_j f_j^2 |T_j|, \end{aligned} \quad (8.18)$$

using (5.14). This implies (8.14).  $\square$

REMARK 8.3. The terms

$$\|v\|_{W^{2,2}(\Omega)}^2 \quad \text{and} \quad \sum_{\mathcal{T}_R} R_j^2 |T_j| \quad (8.19)$$

contained on the right-hand side of the estimate (8.1) depend of course on  $\varepsilon$ . As we will see in the following Lemma, the sum  $\sum_{\mathcal{T}_R} R_j^2 |T_j|$  is of the same order in  $\varepsilon$  as is the norm  $\|v\|_{W^{2,2}(\Omega)}^2$ .

LEMMA 8.4. *Let  $v \in W^{2,2}(\Omega)$  be the solution of (2.1)–(2.2). Then*

$$\|v\|_{W^{2,2}(\Omega)}^2 \leq \frac{C}{\varepsilon^3} \|f\|_{L^2(\Omega)}^2, \quad (8.20)$$

*Proof:* We have the energy estimate

$$\varepsilon \int_{\Omega} |\nabla v|^2 + c_0 \int_{\Omega} |v|^2 \leq C \int_{\Omega} |f|^2 \quad (8.21)$$

and therefore

$$\|v\|_{W^{2,2}(\Omega)}^2 \leq C \int_{\Omega} (\Delta v)^2 \leq \frac{C}{\varepsilon^2} \int_{\Omega} (b \nabla v + cv - f)^2 \leq \frac{C}{\varepsilon^3} \int_{\Omega} f^2.$$

REMARK 8.5. The order of  $\varepsilon$  in (8.20) is sharp. This follows from the following example. For  $x, y \in \mathbb{R}$  let

$$\begin{aligned} a &:= \left(1 - \exp\left(-\frac{1}{\varepsilon}\right)\right)^{-1}, & v_1(x) &:= a \left(1 - \exp\left(-\frac{1-x}{\varepsilon}\right)\right) - 1 + x, & v_2(y) &:= y(1-y), \\ v(x, y) &:= v_1(x)v_2(y), & f(x, y) &:= v(x, y) + v_1(x)(2\varepsilon + 1 - 2y) + v_2(y). \end{aligned}$$

Then we have

$$\begin{aligned} -\varepsilon \Delta v + \partial_x v + \partial_y v + v &= f & \text{in } \Omega &:= ]0, 1[ \times ]0, 1[, \\ v &= 0 & \text{on } \partial\Omega, \end{aligned}$$

and

$$\|\partial_x^2 v\|_{L^2(\Omega)} > \varepsilon^{-\frac{3}{2}} \quad \text{and} \quad \|f\|_{L^2(\Omega)} \leq c \quad \text{for all } \varepsilon > 0. \quad (8.22)$$

It is not difficult to verify these results.

**9. The final estimate.** Putting together the estimates (7.25), (8.1) and using the definition of  $\|z_h\|_{\varepsilon}$  (see (4.6)), we obtain the main estimates (4.7) and (4.8) of Theorem 4.1. The result for the first order scheme (4.10) can be obtained using only the parts of the estimates (7.25), (8.1) which corresponds to the first order parts of the scheme.

**10. Acknowledgement.** The authors want to thank the unknown referees for their valuable comments.

#### REFERENCES

- [1] J. BEY, *Finite-Volumen- und Mehrgitter-Verfahren für elliptische Randwertprobleme*, Teubner Stuttgart, Leipzig, 1998.
- [2] D. BOUCHE, J.-M. GHIDAGLIA AND F.-P. PASCAL, *Error estimate for the upwind finite volume method for the nonlinear scalar conservation law*, J. Comput. Appl. Math. 235, No. 18 (2011), 5394–5410.

- [3] F. BOUCHUT AND B. PERTHAME, *Kruzkov's estimates for scalar conservation laws revisited*, Trans. Am. Math. Soc. 350, No. 7 (1998), 2847–2870.
- [4] A. BRADJI AND J. FUHRMANN, *Some abstract error estimates of a finite volume scheme for a nonstationary heat equation on general nonconforming multidimensional spatial meshes*, Appl. Math. 58, No. 1 (2013), 1–38.
- [5] F. BREZZI, T.J.R. HUGHES, L.D. MARINI, A. RUSSO AND E. SÜLI, *A priori error analysis of residual-free bubbles for advection-diffusion problems*, SIAM J. Numer. Anal. 36, No. 6 (1999), 1933–1948.
- [6] Z. CAI, J. DOUGLAS JR. AND M. PARK, *Development and analysis of higher order finite volume methods for elliptic equations*, Adv. Comput. Math. 19 (2003), 3–33.
- [7] C. CHAINAIS-HILLAIRET, *Second-order finite-volume schemes for a nonlinear hyperbolic equation: Error estimate*, Math. Methods Appl. Sci. 23, No. 5 (2000), 467–490.
- [8] L. CHEN, *A new class of high order finite volume methods for second order elliptic equations*, SIAM J. Numer. Anal. 47 (2010), 4021–4043.
- [9] Z. CHEN, J. WU AND Y. XU, *Higher-order finite volume methods for elliptic boundary value problems*, Adv. Comput. Math. 37 (2012), 191–253.
- [10] Z. CHEN, Y. XU AND Y. ZHANG, *A construction of higher-order finite volume methods*, Math. Comp. 84 (2015), 599–628.
- [11] P. G. CIARLET, *The finite element method for elliptic problems*, North-Holland, 1978.
- [12] B. COCKBURN, F. COQUEL AND P. LEFLOCH, *An error estimate for finite volume methods for multidimensional conservation laws*, Math. Comp. 63, No. 207 (1994), 77–103.
- [13] B. COCKBURN, F. COQUEL AND P. LEFLOCH, *Convergence of the finite volume method for multidimensional conservation laws*, SIAM J. Numer. Anal. 32 (1995), 687–705.
- [14] B. COCKBURN AND P.-A. GREMAUD, *A priori error estimates for numerical methods for scalar conservation laws. Part I: The general approach*, Math. Comp. 65, No. 214 (1996), 533–573.
- [15] B. COCKBURN AND C. W. SHU, *TVB Runge-Kutta projection discontinuous Galerkin finite element method for conservation laws. II: General framework*, Math. Comp., 52 (1989), 411–435.
- [16] B. COCKBURN AND W. ZHANG, *A posteriori error estimates for HDG methods*, J. Sci. Comput. 51, No. 3 (2012), 582–607.
- [17] F. COQUEL AND P. LEFLOCH, *Convergence of finite difference schemes for conservation laws in several space dimensions: the corrected antidiffusive flux approach*, Math. Comp. 57 (1991), 169–210.
- [18] L. CUETO-FELGUEROSO AND I. COLOMINAS, *High-order finite volume methods and multiresolution reproducing kernels*, Arch. Comput. Methods. Eng., DOI 10.1007/s11831-008-9017-y.
- [19] L. J. DURLOFSKY, B. ENGQUIST AND S. OSHER, *Triangle based adaptive stencils for the solution of hyperbolic conservation laws*, J. Comput. Phys. 98 (1992), 64–73.
- [20] G. DZIUK, *Theory and numerics of partial differential equations. (Theorie und Numerik partieller Differentialgleichungen)*, De Gruyter Studium, Berlin, 2010.
- [21] B. ENGQUIST AND S. OSHER, *One-sided difference approximations for nonlinear conservation laws*, Math. Comp. 36 (1981), 321–351.
- [22] K. ERIKSSON, D. ESTEP, P. HANSBO AND C. JOHNSON, *Computational differential equations*, Cambridge University Press, 1996.
- [23] R. EYMARD, R., T. GALLOUËT AND R. HERBIN, *A cell-centred finite-volume approximation for anisotropic diffusion operators on unstructured meshes in any space dimension*, IMA J. Numer. Anal. 26, No. 2 (2006), 326–353.
- [24] R. EYMARD, T. GALLOUËT AND R. HERBIN, *Cell centred discretisation of non linear elliptic problems on general multidimensional polyhedral grids*, J. Numer. Math. 17, No. 3 (2009), 173–193.
- [25] R. EYMARD, G. HENRY, R. HERBIN, F. HUBERT AND R. KLÖFKORN, *3D benchmark on discretization schemes for anisotropic diffusion problems on general grids*. In: Fořt, Jaroslav et al. (eds.), *Finite volumes for complex applications VI: Problems and perspectives*. FVCA 6, international symposium, Prague, Czech Republic, June 2011, Vol. 1 and 2. Springer Proceedings in Mathematics 4 (2011), 895–930.
- [26] M. FEISTAUER, J. FELCMAN AND M. LUKÁČOVÁ-MEDVIĐOVÁ, *On the convergence of a combined finite volume-finite element method for nonlinear convection-diffusion problems*, Num. Meth. PDE, 13 (1997), 1–28.
- [27] B. HEINRICH, *Finite difference methods on irregular networks*, Birkhäuser, Basel, 1987.
- [28] R. HERBIN, *An error estimate for a finite volume scheme for a diffusion-convection problem on triangular mesh*, Num. Meth. PDE, 11 (1995), 165–173.
- [29] P. HOUSTON, J.A. MACKENZIE, E. SÜLI AND G. WARNECKE, *A posteriori error analysis for numerical approximations of Friedrichs systems*, Numer. Math. 82, No. 3 (1999), 433–470.
- [30] L. IVAN, *Development of high-order CENO finite-volume schemes with block-based adaptive mesh refinement*, Thesis, University Toronto, 2011.
- [31] C. JOHNSON AND A. SZEPESSY, *Convergence of a finite element method for a nonlinear hyperbolic conservation law*, Math. Comp. 49 (1987), 427–444.
- [32] C. JOHNSON, *Numerical solution of partial differential equations by the finite element method*, Cambridge University Press, 1994.
- [33] D. KRÖNER, S. NOELLE AND M. ROKYTA, *Convergence of higher order upwind finite volume schemes on unstructured grids for scalar conservation laws in two space dimensions*, Num. Math., 71, No. 4 (1995), 527–560.
- [34] D. KRÖNER AND M. ROKYTA, *Convergence of upwind finite volume schemes for scalar conservation laws in 2D*, SIAM J. Numer. Anal., 31, No. 2 (1994), 324–343.
- [35] D. KUZMIN, *A guide to numerical methods for transport equations*, Preprint, Univ. Erlangen-Nürnberg, 2010.
- [36] G. LUBE, *Streamline diffusion finite element method for quasilinear elliptic problems*, Num. Math. 61 (1992), 335–357.

- [37] M. LUKÁČOVÁ–MEDVIĐOVÁ, *Combined finite element–finite volume method (convergence analysis)*, Comment. Math. Univ. Carolinae 38,3 (1997), 717–741.
- [38] M. OEVERMANN, R. KLEIN, *A Cartesian grid finite volume method for elliptic equations with variable coefficients and embedded interfaces*, J. Comput. Phys. 219 (2006), 749–769.
- [39] B. RIVIERE, *Discontinuous Galerkin methods for solving elliptic and parabolic equations. Theory and implementation*, SIAM, Philadelphia, 2008.
- [40] C. ROHDE, *Weakly coupled hyperbolic systems*, PhD Thesis, Universität Freiburg, 1996.
- [41] E. SÜLI, *A posteriori error analysis and adaptivity for finite element approximations of hyperbolic problems*, in: Kröner, Dietmar (ed.) et al., *An introduction to recent developments in theory and numerics for conservation laws*, Proceedings of the international school, Freiburg/Littenweiler, Germany, October 20–24, 1997. Springer Lect. Notes Comput. Sci. Eng. 5 (1999), 123–194.
- [42] J.-P. VILA, *Convergence and error estimates in finite volume schemes for general multi-dimensional scalar conservation laws. I: Explicit monotone schemes*, M<sup>2</sup>AN 28, No. 3 (1994), 267–295.
- [43] M. VLASÁK, V. DOLEJŠÍ AND J. HÁJEK, *A priori error estimates of an extrapolated space-time discontinuous Galerkin method for nonlinear convection-diffusion problems*, Numer. Meth. Part. D. E. 27, No. 6 (2011), 1456–1482.
- [44] W. WANG, J. GUZMAN, C.-W. SHU, *The multiscale discontinuous Galerkin method for solving a class of second order elliptic problems with rough coefficients*, Int. J. Numer. Anal. Model. 8 (2011), 28–47.
- [45] M. WIERSE, *Higher order upwind schemes on unstructured grids for the compressible Euler equations in time dependent geometries in 3D*, PhD Thesis, Universität Freiburg, 1994.
- [46] C. ZAMFIRESCU, *Survey of two-dimensional acute triangulations*, Discrete Math. 313(1), (2013), 35–49 (see also <http://czamfirescu.tricube.de/CTZamfirescu-08.pdf>).
- [47] Q. ZHANG AND C.-W. SHU, *Error estimates to smooth solutions of Runge-Kutta discontinuous Galerkin method for scalar conservation laws*, SIAM J. Numer. Anal. 42, No. 2 (2004), 641–666.