

NMSA409, topic 3: L_2 -properties

Let $T \subset \mathbb{R}$ be an open interval and consider a stochastic process $\{X_t, t \in T\}$ with continuous time and finite second moments.

Definition 3.1: We call the process $\{X_t, t \in T\}$ L_2 -continuous (mean square continuous) at the point $t_0 \in T$ if $\mathbb{E}|X_t - X_{t_0}|^2 \rightarrow 0$ for $t \rightarrow t_0$. The process is L_2 -continuous if it is L_2 -continuous at all points $t \in T$.

Theorem 3.1: A stochastic process $\{X_t, t \in T\}$ is L_2 -continuous if and only if its mean value $\mathbb{E}X_t$ is a continuous function on T and its autocovariance function $R(s, t)$ is continuous at points $[s, t]$ for which $s = t$.

Corollary 3.1: Centered weakly stationary process is L_2 -continuous if and only if its autocovariance function $R(t)$ is continuous at point 0.

Definition 3.2: We call the process $\{X_t, t \in T\}$ L_2 -differentiable (mean square differentiable) at the point $t_0 \in T$ if there is a random variable X'_{t_0} such that

$$\lim_{h \rightarrow 0} \mathbb{E} \left| \frac{X_{t_0+h} - X_{t_0}}{h} - X'_{t_0} \right|^2 = 0.$$

The random variable X'_{t_0} is called the *derivative in the L_2 (mean square) sense* of the process $\{X_t, t \in T\}$ at the point t_0 . The process is L_2 -differentiable if it is L_2 -differentiable at all points $t \in T$.

Theorem 3.2: A stochastic process $\{X_t, t \in T\}$ is L_2 -differentiable if and only if its mean value $\mathbb{E}X_t$ is differentiable and the second-order generalized partial derivative of the autocovariance function $R(s, t)$ exists and is finite at points $[s, t]$ for which $s = t$, i.e. there is a finite limit

$$\lim_{h, h' \rightarrow 0} \frac{1}{hh'} [R(t+h, t+h') - R(t, t+h') - R(t+h, t) + R(t, t)].$$

Remark: A sufficient condition for the existence of the second-order generalized partial derivative is the existence and continuity of the second-order partial derivatives $\frac{\partial^2 R(s, t)}{\partial s \partial t}$ and $\frac{\partial^2 R(s, t)}{\partial t \partial s}$.

Remark: Any L_2 -differentiable process is also L_2 -continuous.

Definition 3.3: Let $T = [a, b]$ be a bounded closed interval. For any $n \in \mathbb{N}$ let $D_n = \{t_{n,0}, \dots, t_{n,n}\}$ be a division of the interval $[a, b]$ where $a = t_{n,0} < t_{n,1} < \dots < t_{n,n} = b$. We define the partial sums I_n of the centered stochastic process $\{X_t, t \in T\}$ by the formula

$$I_n = \sum_{i=0}^{n-1} X_{t_{n,i}} (t_{n,i+1} - t_{n,i}), \quad n \in \mathbb{N}.$$

If there is a random variable I such that $\mathbb{E}|I_n - I|^2 \rightarrow 0$ for $n \rightarrow \infty$ and for each division of the interval $[a, b]$ such that $\max_{0 \leq i \leq n-1} (t_{n,i+1} - t_{n,i}) \rightarrow 0$ we call it the *Riemann integral* of the process $\{X_t, t \in T\}$ and denote it by $I = \int_a^b X_t dt$. For a non-centered process with the mean value $\mathbb{E}X_t$ we define the Riemann integral as

$$\int_a^b X_t dt = \int_a^b (X_t - \mathbb{E}X_t) dt + \int_a^b \mathbb{E}X_t dt$$

if the centered process $\{X_t - \mathbb{E}X_t, t \in T\}$ has a Riemann integral and the Riemann integral $\int_a^b \mathbb{E}X_t dt$ exists and is finite.

Theorem 3.3: A stochastic process $\{X_t, t \in [a, b]\}$ where $[a, b]$ is a bounded closed interval is Riemann-integrable if the Riemann integrals $\int_a^b \mathbb{E}X_t dt$ and $\int_a^b \int_a^b R(s, t) ds dt$ exist and are finite.

Exercise 3.1: Consider a stochastic process $X_t = \cos(t + B)$, $t \in \mathbb{R}$, where B is a random variable with the uniform distribution on the interval $(0, 2\pi)$. Is this process L_2 -continuous and L_2 -differentiable? Is it Riemann-integrable on a bounded closed interval $[a, b]$?

Exercise 3.2: Let $\{X_t, t \in \mathbb{R}\}$ be a process of independent identically distributed random variables with a mean value μ and a finite variance $\sigma^2 > 0$. What are the L_2 properties of such a process (including Riemann-integrability)?

Exercise 3.3: Consider the Poisson process with intensity λ . Determine the L_2 properties of the process, including Riemann-integrability.

Exercise 3.4: Determine the L_2 properties, including Riemann-integrability, of the Ornstein-Uhlenbeck process $\{U_t, t \geq 0\}$, defined by the formula

$$U_t = e^{-\alpha t/2} W_{\exp\{\alpha t\}}, \quad t \geq 0$$

where $\alpha > 0$ is a positive parameter and $\{W_t, t \geq 0\}$ is a Wiener process.

Exercise 3.5: Let $\{X_t, t \in \mathbb{R}\}$ be a centered stochastic process with the autocovariance function

$$R(t) = \exp\{\lambda(e^{it} - 1)\}, \quad t \in \mathbb{R},$$

where $\lambda > 0$. Determine the L_2 properties of the process, including Riemann-integrability.

Exercise 3.6: Let $\{W_t, t \geq 0\}$ be a Wiener process. We define $B_t = W_t - tW_1$, $t \in [0, 1]$. The stochastic process $\{B_t, t \in [0, 1]\}$ is called the *Brownian bridge*. Determine whether the process $\{B_t, t \in (0, 1)\}$ is L_2 -continuous and L_2 -differentiable. Does the Riemann integral $\int_0^1 B_t dt$ exist?

Exercise 3.7: *Integrated Wiener process* is defined as

$$X_t = \int_0^t W_\tau d\tau, \quad t \geq 0.$$

Using the properties of the Wiener process and L_2 -convergence prove that $X_t \sim N(0, v_t^2)$ for all $t \geq 0$ where $v_t^2 = \frac{1}{3}\sigma^2 t^3$ and σ^2 is the parameter of the Wiener process W_t . Use the fact that the L_2 -limit of a sequence of Gaussian random variables is a Gaussian random variable.