

### IV.3 Holomorphic functional calculus

**Proposition 16** (path integral with values in a Banach space). *Let  $\varphi : [a, b] \rightarrow \mathbb{C}$  be a continuous piecewise  $\mathcal{C}^1$ -smooth curve (i.e.,  $\varphi$  is a continuous mapping and there exists a partition of the interval  $[a, b]$  such that on each of its intervals the derivative  $\varphi'$  is continuous and has finite unilateral limits in the endpoints). Let  $X$  be a Banach space and let  $f : \langle \varphi \rangle \rightarrow X$  be a continuous mapping (where  $\langle \varphi \rangle = \varphi([a, b])$  is the range of  $\varphi$ ). Then the integral*

$$\int_{\varphi} f = \int_a^b f(\varphi(t))\varphi'(t) dt$$

*exists in the Bochner sense.*

**Remarks:**

- (1) Similarly as in the complex analysis, we will also consider the integral over a cycle (i.e., over a formal sum of closed piecewise  $\mathcal{C}^1$ -smooth curves).
- (2) To compute the path integral and to work with it we will use the weak version of the integral, i.e., the equivalence

$$x = \int_{\varphi} f \Leftrightarrow \forall x^* \in X^* : x^*(x) = \int_{\varphi} x^* \circ f.$$

- (3) To define the above-mentioned path integral and to work with it the Bochner theory is not necessary. The integral exists also in the Riemann sense. Where the Riemann integral of a function  $g : [a, b] \rightarrow X$  equals  $x \in X$  if and only if

$\forall \varepsilon > 0 \exists \delta > 0 \forall a = t_0 < t_1 < \dots < t_k = b$  partition of  $[a, b]$  :

$$\max_{1 \leq j \leq k} (t_j - t_{j-1}) < \delta \Rightarrow \forall u_1 \in [t_0, t_1], \dots, u_k \in [t_{k-1}, t_k] : \left\| x - \sum_{j=1}^k f(u_j)(t_j - t_{j-1}) \right\| < \varepsilon.$$

One can show that in our case the Riemann integral exists and, moreover, the equivalence from the preceding remark holds true. This approach can be found in the literature.

**Definition.** Let  $A$  be a Banach algebra with a unit  $e$ , let  $x \in A$  and let  $f$  be a function holomorphic on an open set  $\Omega \subset \mathbb{C}$  containing  $\sigma(x)$ . Let  $\Gamma$  be a „cycle around  $\sigma(x)$  in  $\Omega$ “ (i.e.,  $\Gamma$  is a cycle in  $\Omega$ ,  $\text{ind}_{\Gamma} z$  assumes only the values 0 or 1,  $\text{ind}_{\Gamma} z = 1$  for  $z \in \sigma(x)$  and  $\text{ind}_{\Gamma} z = 0$  for  $z \in \mathbb{C} \setminus \Omega$ ). We define the element  $\tilde{f}(x) \in A$  by the formula

$$\tilde{f}(x) = \frac{1}{2\pi i} \int_{\Gamma} f(\lambda)(\lambda e - x)^{-1} d\lambda.$$

**Remarks:**

- (1) We know from complex analysis that a cycle with the required properties exists.
- (2) The element  $\tilde{f}(x)$  is well defined due to Proposition 16.
- (3) It follows by the Cauchy theorem that the value  $\tilde{f}(x)$  does not depend on a concrete choice of the cycle  $\Gamma$ .
- (4) The mapping  $f \mapsto \tilde{f}(x)$  is called the **holomorphic functional calculus**, or the **Dunford functional calculus**.
- (5) Instead of  $\tilde{f}(x)$  one often writes just  $f(x)$ .

**Theorem 17** (properties of the holomorphic calculus). *Let  $A$  be a Banach algebra with a unit  $e$ , let  $x \in A$  and let  $\Omega \subset \mathbb{C}$  be an open set containing  $\sigma(x)$ .*

- (a) *The mapping  $f \mapsto \tilde{f}(x)$  is an algebraic homomorphism of the unital (commutative) algebra  $H(\Omega)$  into  $A$ .*
- (b)  *$\tilde{id}(x) = x$  and  $\tilde{1}(x) = e$ , where  $id(\lambda) = \lambda$  and  $1(\lambda) = 1$  for  $\lambda \in \Omega$ .*
- (c) *If  $p$  is a polynomial, then  $\tilde{p}(x) = p(x)$  where  $p(x)$  has the meaning from Lemma 11.*
- (d) *If  $\lambda \in \Omega$ , then  $\tilde{f}(\lambda e) = f(\lambda)e$ .*
- (e) *If  $f_n \rightarrow f$  locally uniformly on  $\Omega$  (where  $f_n \in H(\Omega)$  for each  $n \in \mathbb{N}$ ), then  $\tilde{f}_n(x) \rightarrow \tilde{f}(x)$  in  $A$ .*
- (f)  *$\tilde{f}(x) \in G(A)$  if and only if  $f(\lambda) \neq 0$  for each  $\lambda \in \sigma(x)$ .*
- (g)  *$\sigma(\tilde{f}(x)) = f(\sigma(x))$ .*
- (h)  *$(\tilde{g} \circ \tilde{f})(x) = \tilde{g}(\tilde{f}(x))$  whenever  $f \in H(\Omega)$ ,  $g \in H(\Omega')$ ,  $\Omega' \supset f(\sigma(x))$ .*
- (i) *If  $y \in A$  commutes with  $x$  (i.e.,  $xy = yx$ ), then  $y$  commutes with  $\tilde{f}(x)$  for each  $f \in H(\Omega)$ .*

**Remarks.** Let  $A$  be a unital Banach algebra and let  $x \in A$ .

- (1) If  $f$  and  $g$  are two holomorphic functions on a neighborhood of  $\sigma(x)$ , which coincide on a neighborhood of  $\sigma(x)$ , then  $\tilde{f}(x) = \tilde{g}(x)$ .
- (2) It may happen that  $f$  and  $g$  coincide on  $\sigma(x)$  and  $\tilde{f}(x) \neq \tilde{g}(x)$ .
- (3) The assignment  $f \mapsto \tilde{f}(x)$  need not be one-to-one. I.e., the equality  $\tilde{f}(x) = \tilde{g}(x)$  does not imply that  $f$  and  $g$  coincide on a neighborhood of  $\sigma(x)$ .
- (4) If  $\tilde{f}(x) = \tilde{g}(x)$ , then  $f|_{\sigma(x)} = g|_{\sigma(x)}$ .