

Proof of Lemma XI.4

(X, \mathcal{T}) - LCS. Then $X^* \subset X^\#$

(a) The topology $\sigma(X^\#, X)$ is Hausdorff

[clear, as X separates points of $X^\#$

$$- f \in X^\#, f \neq 0 \Rightarrow \exists x \in X - f(x) \neq 0$$

The topology $\sigma(X^*, X)$ on X^* is the subspace topology generated by $\sigma(X^\#, X)$

[clear from definitions]

(b) (X, \mathcal{T}) Hausdorff $\Rightarrow X^*$ is $\sigma(X^\#, X)$ -dense in $X^\#$

[(X, \mathcal{T}) Hausdorff $\Rightarrow X^*$ separates points of X
(a consequence of the H-B theorem)]

i.e. $(X^*)^\perp = \{0\}$. Thus $((X^*)^\perp)^\perp = X^\#$.

By the bipolar theorem we see that $X^\# = \overline{X^*}^{\sigma(X^\#, X)}$

(c) $A \subset X^*$. Then A is $\sigma(X^*, X)$ -relatively compact in X^*

\Leftrightarrow A is $\sigma(X^*, X)$ -bdcd

$$\overline{A}^{\sigma(X^*, X)} \subset X^*$$

\Rightarrow A rel. compact $\Rightarrow \overline{A}^{\sigma(X^*, X)}$ is $\sigma(X^*, X)$ -compact

This $\forall x \in X$ $f \mapsto f(x)$ is bdd on A , i.e., A is $\sigma(X^*, X)$ -bdcd

Moreover, $\overline{A}^{\sigma(X^{\#}, X)}$ is $\sigma(X^{\#}, X)$ -compact, thus $\sigma(X^{\#}, X)$ -compact
 thus $\sigma(X^{\#}, X)$ closed (as the topology is Hausdorff)

It follows $\overline{A}^{\sigma(X^{\#}, X)} = \overline{A}^{\sigma(X^{\#}, X)} \subset X$

Define $q_A(x) = \sup \{ |f(x)| ; f \in A \}$, $x \in X$

As A is $\sigma(X^{\#}, X)$ -bdcl, it is a well-defined
 seminorm. So, it is $\sup \mathcal{L}(X)$ -cts.

The $A_0 = \{ x \in X ; q_A(x) \leq 1 \}$ is a $\sigma(X^{\#}, X)$ -bdcl
 of 0. Therefore

$\overline{a \circ A}^{\sigma(X^{\#}, X)} = (A_0)^{\circ}$ is $\sigma(X^{\#}, X)$ -compact
 (Banach-Alaoglu)
 \uparrow
 Bipolar theorem

This also $\overline{A}^{\sigma(X^{\#}, X)}$ is $\sigma(X^{\#}, X)$ -compact,
 therefore $\sigma(X^{\#}, X)$ -compact

Lemma XI.6: X normed space $A \subset X^{\#}$ $\sigma(X^{\#}, X)$ bdcl
 $f \in X^{\#}$ The $\overline{A}^{\sigma(X^{\#}, X)}$
 $|f| \leq q_A \Leftrightarrow f \in a \circ A$

Proof:

Observe that $|f| \leq q_A$ means that $f \in (A_0)^{\circ}$
 and use the Bipolar theorem