

Let A be a unital C^* -algebra, e be its unit and $x \in A$ be a normal element. (i.e., $x^*x = xx^*$)

① Let $B := \overline{\text{alg}} \{e, x, x^*\}$

Then clearly B is a commutative C^* -algebra,
 B is a C^* -subalgebra of A
 $e \in B$, it is also unit of B

Thus, by Prop. XI.12 $\forall y \in B : \sigma_B(y) = \sigma_A(y)$. So, we will write only $\sigma(y)$.

② Let $h = \hat{x}$; i.e. $h(\varphi) = \varphi(x)$, $\varphi \in \Delta(B)$

Then h is a homeomorphism of $\Delta(B)$ onto $\sigma(x)$

$\Gamma \circ h = \hat{x}$ is cts on $\Delta(B)$

• $h(\Delta(B)) = \sigma(x)$ by Thm X.25 (e)

• h is one-to-one:

$$\varphi_1, \varphi_2 \in \Delta(B) : h(\varphi_1) = h(\varphi_2)$$

$$\text{Then: } \varphi_1(e) = \varphi_2(e) = 1$$

$$\varphi_1(x) = \varphi_2(x)$$

$$\varphi_1(x^*) = \overline{\varphi_1(x)} = \overline{\varphi_2(x)} = \varphi_2(x^*)$$



Prop. XI.8cc)

$\{y \in B : \varphi_1(y) = \varphi_2(y)\}$ is a closed algebra containing e, x, x^* , so it equals B

$$\text{Hence } \varphi_1 = \varphi_2$$

• $\Delta(B)$ compact, h cts and one-to-one \Rightarrow

h is a homeomorphism \downarrow

(3) Let $\Gamma: B \rightarrow \mathcal{L}(\Delta(B))$ be the Gelfand transform of B

For $f \in \mathcal{L}(\sigma(x))$ define $\tilde{f}(x) := \Gamma^{-1}(f \circ h)$

(4) $\Phi: f \mapsto \tilde{f}(x)$ is an isometric $*$ -isomorphism of $\mathcal{L}(\sigma(x))$ onto B

Γ is a homeomorphism $\Rightarrow f \mapsto f \circ h$ is an isometric $*$ -isomorphism of $\mathcal{L}(\sigma(x))$ onto $\mathcal{L}(\Delta(B))$

Γ is an isometric $*$ -isomorphism of B onto $\mathcal{L}(\Delta(B))$ by Theorem XI.9

So, Φ is such, as a composition of two such maps. \Downarrow

(5) $\tilde{1}(x) = 1$ (Φ preserves the unit)

$\tilde{c}d(x) = x$ ($\Gamma(x) = \hat{x} = h = \text{id} \circ h$)

(6) P is a polynomial $\Rightarrow \tilde{P}(x) = P(x)$

Γ Thus follows from (4) and (5) \Downarrow

(7) $\sigma(\tilde{f}(x)) = f(\sigma(x))$ for $f \in \mathcal{L}(\sigma(x))$.

Γ Φ is an isometric $*$ -isomorphism $\Rightarrow \Phi$ preserves spectrum
 $\Rightarrow \sigma(\tilde{f}(x)) = \sigma(\Phi(x)) = \sigma(x) = f(\sigma(x))$ \Downarrow

(8) If $y \in A$ commutes with x , it commutes with $\tilde{f}(x)$ for each $f \in \mathcal{L}(\sigma(x))$

$\Gamma \{z \in A; zy = yz\}$ is a closed subalgebra of A containing $1, x$ and also x^* (by Thm XI.13)
So, it contains B . $_$