

$$\int_0^{4\pi} \frac{1}{\sin^2 x + 2\cos^2 x} dx$$


celkem: $\frac{\sqrt{2}\pi}{2} (1+1+3\cdot 2)$
 $= 2\sqrt{2}\pi$

$t = \frac{1}{\sqrt{2}} x$

$$\int \frac{1}{\frac{t^2}{1+t^2} + 2 \frac{1}{1+t^2}} \frac{dt}{\sqrt{2}} = \int \frac{1}{t^2+2} dt = \int \frac{1}{2(\frac{t^2}{2}+1)} dt$$

$$= \frac{1}{2} \int \frac{1}{1+(\frac{t}{\sqrt{2}})^2} dt = \frac{1}{2} \cdot \sqrt{2} \arctan \frac{t}{\sqrt{2}} + C$$

$$= \left(\frac{\sqrt{2}}{2} \arctan \frac{\tan x}{\sqrt{2}} \right) + C \quad x \in \left(-\frac{\pi}{2} + 2\pi, \frac{\pi}{2} + 2\pi \right)$$



$$\int_0^{\pi/2} \rightarrow \int_0^{\infty} \frac{1}{t^2+2} dt = \left[\frac{\sqrt{2}}{2} \arctan \frac{t}{\sqrt{2}} \right]_0^{\infty}$$

$$= \frac{\sqrt{2}}{2} \cdot \frac{\pi}{2}$$

$$\int_{\pi/2}^{4\pi} \rightarrow \int_{-\infty}^0 \frac{1}{t^2+2} dt = \left[\frac{\sqrt{2}}{2} \arctan \frac{t}{\sqrt{2}} \right]_{-\infty}^0 = - \frac{\pi}{2} \cdot \frac{\sqrt{2}}{2} = - \frac{\sqrt{2}}{2} \frac{\pi}{2}$$

asoban!
3ks

$$\int_{-\infty}^{\infty} \frac{1}{t^2+2} dt = \left[\frac{\sqrt{2}}{2} \arctan \frac{t}{\sqrt{2}} \right]_{-\infty}^{\infty} = \frac{\sqrt{2}}{2} \left(\frac{\pi}{2} - \left(-\frac{\pi}{2} \right) \right)$$

$$10.04 \quad \int_{-\infty}^0 \frac{x}{x^3-1} dx =$$

$$\frac{1}{0!k} \int_{-\infty}^0 \frac{x}{x^3-1} dx =$$

$$= \int_{-\infty}^0 \frac{1}{3} \left(\frac{1}{x-1} + \frac{1}{2} \frac{2x-2}{x^2+x+1} \right) dx$$

$$x = A(x^2+x+1) + (Bx+C)(x-1)$$

$$x=1 \Rightarrow A=1/3 \Rightarrow B=-1/3, C=1/3$$

ahn. et.

$$= \frac{1}{3} \left[\ln \frac{|x-1|}{|x^2+x+1|} \right]_{-\infty}^0 + \left[\frac{1}{2} \cdot \frac{2}{\sqrt{3}} \arctan \frac{2x+1}{\sqrt{3}} \right]_{-\infty}^0$$

$$= 0 + \frac{1}{3} \left(\frac{\pi}{2} + \frac{\pi}{6} \right) = \underline{\underline{\frac{2\pi}{9}}}$$

$$10.03 \quad \int_4^{+\infty} \frac{x}{(x-1)(x-2)(x-3)} dx =$$

$$= \int_4^{\infty} \frac{1}{2} \left(\frac{1}{x-1} + \frac{-4}{x-2} + \frac{3}{x-3} \right)$$

$$x = A(x-2)(x-3) + B(x-1)(x-3) + C(x-1)(x-2)$$

$$x=1 \Rightarrow A=1/2$$

$$x=2 \Rightarrow B=-2$$

$$x=3 \Rightarrow C=3/2$$

$$= \frac{1}{2} \left[\ln \frac{(x-1)(x-3)^3}{(x-2)^4} \right]_4^{\infty} = -\frac{1}{2} \ln \frac{3 \cdot 1^3}{2^4}$$

$$= \underline{\underline{\frac{1}{2} \ln \frac{16}{3}}}$$

10.20. $\int_0^4 \frac{a \sin x}{\cos^2 x + 1} dx =$

$t = \cos x,$

$= \int_{-1}^1 \frac{1}{t^2 + 1} dt = [\arctan t]_{-1}^1 = \underline{\underline{\frac{\pi}{2}}}$


10.24. $\int_0^{+\infty} \frac{e^{-1/x}}{x^2} dx = \int_0^{+\infty} e^{-t} dt = [-e^{-t}]_0^{+\infty} = \underline{\underline{1}}$

10.23. $\int \frac{\sqrt{x}}{\sqrt{-(x^2+1)} \arctan x} dx = \int_{\arctan x}^{\pi/3} \frac{dt}{t} = \underline{\underline{\frac{4}{3} \ln \frac{4}{3}}}$

10.22. $\int_{-\infty}^0 x e^{-x^2} dx = \int_{-\infty}^0 \frac{1}{2} e^{-t} dt =$

$= [-\frac{1}{2} e^{-t}]_0^{-\infty} = \underline{\underline{-\frac{1}{2}}}$

10.22.



$$\frac{10.80.}{OK} \int_0^9 \frac{dx}{\sqrt{x+16} - \sqrt{x}} = \int_0^9 \frac{\sqrt{x+16} + \sqrt{x}}{16} dx$$

$$= \frac{1}{16} \left[\frac{(x+16)^{3/2}}{3/2} + \frac{x^{3/2}}{3/2} \right]_0^9 = \frac{1}{16} \left[\frac{121\sqrt{27}}{24} + \frac{27\sqrt{27}}{24} \right] = \frac{144}{16} = 9$$

$$\frac{10.24.}{?} \int_{-\infty}^{+\infty} \frac{e^x}{e^{2x} + 1} dx \quad t = e^x$$

$$= \int_0^{+\infty} \frac{dt}{t^2 + 1} = \left[\frac{2}{\sqrt{3}} \arctan \frac{2t+1}{\sqrt{3}} \right]_0^{+\infty} = \frac{2}{\sqrt{3}} \left(\frac{\pi}{2} - \frac{\pi}{6} \right) = \frac{2\sqrt{3}\pi}{9}$$

$$\frac{10.28.}{t=e^{-x}} \int_0^{+\infty} \frac{dx}{\sqrt{e^{2x} + 1}} = \int_0^{+\infty} \frac{dt}{\sqrt{1+t^2}}$$

$$= \int_0^1 \frac{dt}{\sqrt{1+t^2}} = \left[\ln \left(t + \sqrt{t^2+1} \right) \right]_0^1 = \ln(1+\sqrt{2})$$

$$\frac{10.22.}{t=e^x} \int_{-\infty}^{+\infty} \frac{e^x}{e^{2x} + 3e^x + 3} dx = \int_0^{+\infty} \frac{dt}{t^2 - 3t + 3}$$

$$= \int_0^{+\infty} \frac{dt}{(t - \frac{3}{2})^2 + \frac{3}{4}} = \left[\frac{2}{\sqrt{3}} \arctan \frac{2t-3}{\sqrt{3}} \right]_0^{+\infty} = \frac{2}{\sqrt{3}} \left(\frac{\pi}{2} - \frac{\pi}{3} \right) = \frac{\sqrt{3}\pi}{9}$$

$$\frac{10.26.}{t=e^x} \int_{-\infty}^{+\infty} \frac{e^x}{e^{2x} + e^{-2x}} dx = \int_{-\infty}^{+\infty} \frac{1}{t^2 + 1} dt$$

$$= \int_0^{+\infty} \frac{dt}{t^2 + 1} = \left[\arctan t \right]_0^{+\infty} = \frac{\pi}{2}$$

and as 10.26.

10.35. $\int_0^{\pi} \sin^2 x \cos 2x \, dx =$

$\int_0^{\pi} \frac{1}{4} \sin^2 2x \, dx =$
 $t = 2x$

$= \int_0^{2\pi} \frac{1}{8} \sin^2 t \, dt = \frac{1}{8} \left[\frac{t + \sin t \cos t}{2} \right]_0^{2\pi}$
 $= \frac{\pi}{8}$

10.36. $\int_{-\pi/2}^{\pi/2} x^3 \sin x \, dx =$

$\int_{-\pi/2}^{\pi/2} x^3 \cos x \, dx + \int_{-\pi/2}^{\pi/2} 3x^2 \cos x \, dx =$

$= \left[3x^2 \sin x \right]_{-\pi/2}^{\pi/2} - \int_{-\pi/2}^{\pi/2} 6x \sin x \, dx =$
 $\frac{3}{2} \pi^2$

$= \frac{3}{2} \pi^2 - 6 \int_{-\pi/2}^{\pi/2} \cos x \, dx = \frac{3}{2} \pi^2 - 12$

10.33. $\int_0^{\pi} \sin^3 x \, dx = \int_0^{\pi} (1 - \cos^2 x) \sin x \, dx$
 $t = \cos x$

$= \int_{-1}^1 (1 - u^2) \, du = \left[u - \frac{u^3}{3} \right]_{-1}^1 =$
 $= \frac{2}{3} + \frac{2}{3} = \frac{4}{3}$

10.32. $\int_{\pi/4}^{\pi/2} \log x \, dx = \int_0^{\pi/4} \frac{\sin x}{\cos x} \, dx$
 $t = \cos x$

$= \int_{\sqrt{2}/2}^1 \frac{dt}{t} = \left[\ln t \right]_{\sqrt{2}/2}^1 = -\ln \frac{1}{\sqrt{2}} = \ln \sqrt{2}$
 $= \frac{1}{2} \ln 2$

10.31. $\int_0^{\infty} \frac{dx}{(kx^2 + 2lx + 2)^2}$
 $t = kx$

$= \int_0^{\infty} \frac{dt}{(t^2 + 2t + 2)^2} = \int_0^{\infty} \frac{dt}{((t+1)^2 + 1)^2}$
 $\text{Let } t+1 = \tan u$

$= \int_{\pi/4}^{\pi/2} \cos^3 u \, du = \left[\frac{\sin u \cos^2 u}{2} \right]_{\pi/4}^{\pi/2} = \frac{1}{8} - \frac{1}{4}$

10.42. $\int_{-1}^1 x^2 e^{-x} dx =$ PP

$= [-x^2 e^{-x}]_{-1}^1 + \int_{-1}^1 2x e^{-x} dx =$ PP

$= (e - e^{-1}) + [-2x e^{-x}]_{-1}^1 + 2 \int_{-1}^1 e^{-x} dx =$

$= (e - e^{-1}) + (2e - 2e^{-1}) + 2 [-e^{-x}]_{-1}^1 =$

$= 4(e - e^{-1})$

10.41. $\int_0^{+\infty} (3x \cos x^3 - \frac{1}{2} \sin x^3) dx$

~~PP~~ ~~PP~~

$\int_0^{+\infty} \frac{1}{x^2} \sin x^3 =$ PP $[-\frac{1}{x} \sin x^3]_0^{+\infty} + \int_0^{+\infty} \frac{1}{x} 3x^2 \cos x^3$

$= 0 + \int_0^{+\infty} 3x \cos x^3 dx$

$\Rightarrow I = 0.$

Je n'ai rien vu, je prie pour moi !!

10.50. $\int_0^{\pi/4} \sqrt{\cos x - \cos^3 x} dx =$

$= \int_0^{\pi/4} \cos x \sin x dx =$

$t = \cos x$

$= \int_{\sqrt{3}/2}^1 \sqrt{t} dt = [\frac{2}{3/2} t^{3/2}]_{\sqrt{3}/2}^1 =$

$= \frac{2}{3} (1 - \sqrt{\frac{3}{8}})$

10.39. $\int_{\pi/35}^{\pi/7} \frac{1}{\sqrt{x}} \sin^2 \frac{1}{x} dx$

$t = \frac{1}{x}, dt = -\frac{1}{x^2} dx$

$= \int_{\pi}^{3\pi} \sin^2 t dt = 2 \cdot \int_{\pi}^{2\pi} \sin^2 t dt =$

$\int_{\pi}^{2\pi} \sin^2 t dt = 2 \cdot [\frac{t + \sin t \cos t}{2}]_{\pi}^{2\pi}$

MEMOIR !!
NOUVEAU !!

je ZPF = PF

509!

$$\begin{aligned}
 \text{10.41.} \quad & \int_0^1 x^2 \ln(1+x^2) dx \stackrel{DP}{=} \\
 & \int_0^1 \left[\frac{x^3}{3} \ln(1+x^2) \right]_0^1 - \int_0^1 \frac{2x^4}{3(1+x^2)} dx = \\
 & \quad = \frac{1}{3} \ln 2 \\
 & = \frac{1}{3} \ln 2 - \frac{2}{3} \int_0^1 \left(\frac{x^4+x^2}{1+x^2} - \frac{x^2+1}{x^2+1} + \frac{1}{x^2+1} \right) dx \\
 & = \frac{1}{3} \ln 2 - \frac{2}{3} \int_0^1 \left(x^2 - 1 + \frac{1}{x^2+1} \right) dx = \\
 & = \frac{1}{3} \ln 2 - \frac{2}{3} \left[\frac{x^3}{3} - x + \arctan x \right]_0^1 = \\
 & = \frac{1}{3} \ln 2 - \frac{2}{3} \left[-\frac{2}{3} + \frac{\pi}{4} \right] = \\
 & \underline{\underline{= \frac{1}{3} \ln 2 + \frac{4}{9} - \frac{\pi}{6}}}.
 \end{aligned}$$

$$\begin{aligned}
 \text{10.44.} \quad & \int_{-\pi}^{\pi/2} \frac{\cos^3 x}{\sqrt{\sin x}} dx = \int_{-1}^{-\frac{\sqrt{2}}{2}} + \int_{-\frac{\sqrt{2}}{2}}^{\frac{1}{2}} \\
 & \int_{-\pi}^{\pi/2} \frac{\cos^3 x}{\sqrt{\sin x}} dx = \int_{-1}^{-\frac{\sqrt{2}}{2}} \dots + \int_{-\frac{\sqrt{2}}{2}}^{\frac{1}{2}} \dots = \\
 & = \int_0^1 \frac{1-t^2}{3\sqrt{t}} dt = \left[\frac{t^{3/2}}{3/2} - \frac{t^{5/2}}{5/2} \right]_0^1 = \\
 & = \frac{2}{3} \frac{3}{2} - \frac{2}{5} \frac{3}{2} = \underline{\underline{\frac{9}{8}}}.
 \end{aligned}$$

$$\begin{aligned}
 \text{10.43.} \quad & \int_0^{+\infty} \frac{dx}{\sqrt{e^x-1}} = \\
 & \quad t = \sqrt{e^x-1}, \quad dt = \frac{1 \cdot e^x}{2\sqrt{e^x-1}} dx \\
 & \quad t^2+1 = e^x \\
 & = \int_0^{+\infty} \frac{2}{2(t^2+1)} dt = 2 [\arctan t]_0^{+\infty} = \underline{\underline{\pi}}.
 \end{aligned}$$

10.49. $\int_0^1 x \cos x^2 dx$

~~let~~ $x = \cos t$; $dx = -\sin t dt$

$\int_0^{\pi/2} t^2 \sin t dt = \int_0^{\pi/2} [t^2 \cos t]_0^{\pi/2} - \int_0^{\pi/2} 2t \cos t dt =$

$= -[-2t \sin t]_0^{\pi/2} - \int_0^{\pi/2} 2 \sin t dt =$

$= \pi - 2[-\cos t]_0^{\pi/2} = \underline{\underline{\pi - 2}}$

10.48. $\int_0^1 x \arcsin x dx =$

OK $x = \sin t$, $dx = \cos t dt$

$= \int_0^{\pi/2} t \sin t \cos t dt =$

$= \frac{1}{2} \int_0^{\pi/2} 2t \sin 2t dt = \frac{1}{8} \int_0^{\pi} u \sin u du =$

PP $= \frac{1}{8} [u \cos u]_0^{\pi} + \frac{1}{8} \int_0^{\pi} \cos u du = \frac{1}{8} \pi + \frac{1}{8} [\sin u]_0^{\pi} =$

10.51. $\int_1^{\infty} \frac{1 - \ln x}{x^2} dx$

OK $t = \ln x$, $dt/dx = 1/x$, $x = e^t$

$= \int_0^{\infty} \frac{1-t}{e^{2t}} dt = \int_0^{\infty} (e^{-t} - te^{-t}) dt =$

$= [-e^{-t}]_0^{\infty} - \int_0^{\infty} te^{-t} dt =$

$= 1 - [-te^{-t}]_0^{\infty} - \int_0^{\infty} e^{-t} dt = \underline{\underline{2}}$

10.56. $\int_1^e \frac{1 + \ln x}{x} dx$

OK $t = \ln x$, $dt = \frac{1}{x} dx$

$= \int_0^1 (1+t) dt = [t + \frac{t^2}{2}]_0^1 = \underline{\underline{3/2}}$

10.64. $\int_0^1 x \sqrt{1-x^2} dx =$

* $t = \sqrt{1-x^2}$, $dt = -2x dx$

$x^2 = 1-t$

$= \int_0^1 \frac{1}{2} (1-t)^2 \sqrt{t} dt =$
 $(1-2t+t^2) dt$

$= \frac{1}{2} \left[\frac{t^{3/2}}{3/2} - 2 \cdot \frac{t^{5/2}}{5/2} + \frac{t^{7/2}}{7/2} \right]_0^1 =$

$= \frac{1}{4} \left[\frac{1 \cdot 2}{3} - \frac{2 \cdot 4}{5} + \frac{2 \cdot 1}{7} \right] = \frac{35 - 42 + 14}{105} = \underline{\underline{\frac{8}{105}}}$

10.63. $\int_0^1 x^2 \sqrt{1-x^2} dx =$

OK $x = \sin t$, $dx = \cos t dt$

$= \int_0^{\pi/2} \sin^2 t \cos^2 t dt = \int_0^{\pi/2} (\sin^2 t - \sin^4 t) dt$

~~$= \int_0^{\pi/2} \sin^2 t dt - \int_0^{\pi/2} \sin^4 t dt$~~
 $= \frac{3}{16} \pi - \frac{5}{64} \pi = \underline{\underline{\frac{1}{32} \pi}}$

10.62. $\int_0^1 x^2 \sqrt{1-x^2} dx =$

* ~~$x = \sin t$~~ , $dx = \cos t dt$

$= \int_0^{\pi/2} \sin^2 t \cos^2 t dt = \int_0^{\pi/2} \frac{1}{4} \sin^2 2t dt =$

$= \int_0^{\pi/2} \frac{1}{8} \sin^2 u du =$
 $u = 2t, du = 2 dt$

$= \frac{1}{8} \left[\frac{u - \cos u \sin u}{2} \right]_0^{\pi/2} = \underline{\underline{\frac{1}{16} \pi}}$

17. $\int_0^{\pi/2} \sin^6 t dt = \int_0^{\pi/2} \sin^4 t \cos^2 t dt = \int_0^{\pi/2} \sin^4 t dt - \int_0^{\pi/2} \sin^6 t dt$

prok. $\int_0^{\pi/2} \sin^4 t dt = \int_0^{\pi/2} \sin^2 t (1 - \cos^2 t) dt = \int_0^{\pi/2} \sin^2 t dt - \int_0^{\pi/2} \sin^2 t \cos^2 t dt$

$\Rightarrow \int_0^{\pi/2} \sin^4 t dt = \frac{3}{8} \int_0^{\pi/2} \sin^2 t dt$

$\int_0^{\pi/2} \sin^2 t dt = \int_0^{\pi/2} \left(\frac{1 - \cos 2t}{2} \right) dt = \int_0^{\pi/2} \left(\frac{1}{2} - \frac{\cos 2t}{2} \right) dt$

$= \left[\frac{1}{8} \pi - \frac{1}{4} [\sin 2t]_0^{\pi/2} + \frac{1}{8} \left[\frac{2t + \sin 2t \cos 2t}{2} \right]_0^{\pi/2} \right] =$
 $= \frac{1}{8} \pi - 0 + \frac{1}{8} \frac{\pi}{2} = \underline{\underline{\frac{3}{16} \pi}}$

$$10.71 \int_0^1 \frac{x+1}{x+0} dx$$

$$x = \frac{-1}{1-t^2} = \frac{1}{t^2-1}$$

$$\frac{dx}{dt} = \frac{-2t}{(t^2-1)^2}$$

$$= \int_{\sqrt{2}}^{+\infty} t \cdot \frac{2t}{(t^2-1)^2} dt =$$

$$= \int_{\sqrt{2}}^{+\infty} \frac{1}{t-1} + \frac{1}{(t-1)^2} + \frac{-1}{t+1} + \frac{1}{(t+1)^2} dt$$

$$2t^2 = A(t-1)(t+1)^2 + B(t+1)^2 + C(t+1)(t-1)^2 + D(t-1)^2$$

$$t=1 \Rightarrow B = \frac{1}{2} \quad \text{or} \quad A+C=D$$


$$t=-1 \Rightarrow D = \frac{1}{2} \quad \text{or} \quad -A+B+C+D=0$$

$$2C - B - D = -1 \Rightarrow C = -\frac{3}{2}$$

$$A = \frac{1}{2}$$

$$\frac{1}{2} \left[\ln \left| \frac{t-1}{t+1} \right| - \frac{2t}{(t^2-1)} \right]_{\sqrt{2}}^{\infty} = \frac{1}{2} \left[\ln \left| \frac{\sqrt{2}-1}{\sqrt{2}+1} \right| + \frac{2\sqrt{2}}{2-1} \right]$$

$$= \frac{1}{2} \ln \left| \frac{\sqrt{2}+1}{\sqrt{2}-1} \right| + \sqrt{2} = \frac{1}{2} \cdot \ln (1+\sqrt{2})^2 + \sqrt{2} = \ln (1+\sqrt{2}) + \sqrt{2}$$



10.71

$$\frac{10. \int dx}{OK} \int_4^{+\infty} \frac{1}{x^2} \sqrt{\frac{x-2}{x-4}} dx =$$

$$t = \sqrt{\frac{x-2}{x-4}} \Rightarrow x = \frac{4t^2 - 4}{1-t^2}$$

$$\Rightarrow x = \frac{-4t^2 + 2}{1-t^2}$$

$$\frac{dx}{dt} = \frac{-8t(1-t^2) + 2t(-4t^2+2)}{(1-t^2)^2}$$

$$= \frac{-4t}{(1-t^2)^2}$$

$$= \int_1^{+\infty} \frac{4t}{(1-t^2)^2} dt$$

~~$$t = \sqrt{\frac{x-2}{x-4}} \Rightarrow dt = \frac{1}{2} \frac{1}{\sqrt{x-4}} dx = \frac{1}{2} \frac{1}{\sqrt{4t^2-2}} dx$$~~

~~$$dx = 2\sqrt{4t^2-2} dt$$~~

$$= \int_1^{+\infty} \frac{4t^2}{(4t^2-2)^2} dt = \int_1^{+\infty} \frac{4t^2}{(2t-\sqrt{2})^2 (2t+\sqrt{2})^2} dt$$

~~$$A \frac{1}{2t-\sqrt{2}} + \frac{B}{(2t-\sqrt{2})^2} + \frac{C}{2t+\sqrt{2}} + \frac{D}{(2t+\sqrt{2})^2}$$~~

$$4t^2 = A(2t-\sqrt{2})(2t+\sqrt{2})^2 + B(2t+\sqrt{2})^2 + C(2t-\sqrt{2})^2(2t+\sqrt{2}) + D(2t-\sqrt{2})^2$$

$$t = \frac{1}{2}\sqrt{2} \Rightarrow 8B = 2 \Rightarrow B = \frac{1}{4}$$

$$t = -\frac{1}{2}\sqrt{2} \Rightarrow 8D = 2 \Rightarrow D = \frac{1}{4}$$

$$8A + 8C = 0 \Rightarrow A = -C$$

$$-2\sqrt{2}A + 2\sqrt{2}C + 2D = 0$$

$$4\sqrt{2}C = -1$$

$$C = -\frac{1}{4\sqrt{2}}$$

$$A = \frac{1}{4\sqrt{2}}$$

~~$$\int_1^{+\infty} \left[\frac{1}{4\sqrt{2}} \ln \left| \frac{2t-\sqrt{2}}{2t+\sqrt{2}} \right| - \frac{1}{8} \frac{1}{(2t-\sqrt{2})} - \frac{1}{8} \frac{1}{(2t+\sqrt{2})} \right] dt$$~~

~~$$= \frac{1}{8\sqrt{2}} \ln \left| \frac{2+\sqrt{2}}{2-\sqrt{2}} \right| + \frac{1}{8} \left(\frac{1}{2-\sqrt{2}} + \frac{1}{2+\sqrt{2}} \right) = \frac{\sqrt{2}}{8} \ln(3+\sqrt{2}) + \frac{1}{4}$$~~

$$89. \int_0^{\pi/2} \frac{\sin x}{\cos^2 x + 3 \cos x + 4} dx$$

$$t = \cos x$$

$$= \int_0^1 \frac{dt}{t^2 + 3t + 4} = \int_0^1 \frac{dt}{(t + \frac{3}{2})^2 + \frac{7}{4}}$$

$$= \left[\frac{2}{\sqrt{7}} \arctan \left(\frac{2t+3}{\sqrt{7}} \right) \right]_0^1 =$$

$$= \frac{2}{\sqrt{7}} \left(\arctan \frac{5}{\sqrt{7}} - \arctan \frac{3}{\sqrt{7}} \right) = \frac{2}{\sqrt{7}} \cdot \arctan \frac{\frac{5}{\sqrt{7}} - \frac{3}{\sqrt{7}}}{1 + \frac{15}{7}}$$

$$= \frac{2}{\sqrt{7}} \arctan \frac{2\sqrt{7}}{17} = \arctan \frac{2\sqrt{7}}{17}$$

$$x = \arccos t, y = \arccos t$$

$$\arccos t = \arccos y = \arccos \left(\frac{x+y}{2} \right)$$

$$\arccos t = \arccos \left(\frac{x+y}{2} \right)$$

$$88. \int_{-\pi/4}^{\pi/4} \frac{dx}{\cos^3 x} =$$

$$= \int_{-1}^1 (t^2 + 1) dt = \left[\frac{t^3}{3} + t \right]_{-1}^1 =$$

$$t = \tan x \dots = \frac{4}{3} + \frac{4}{3} = \frac{8}{3}$$

$$87. \int_0^{\pi/4} \frac{dx}{\cos x + 2 \sin x + 3} = \frac{1}{\sqrt{5}} \left[\frac{1}{2} \ln \left| \frac{1-t}{1+t} \right| + \frac{1}{\sqrt{5}} \arctan \frac{t}{\sqrt{5}} \right]_0^{\pi/4}$$

$$t = \tan \frac{x}{2} \quad dx = \frac{2}{1+t^2} dt$$

$$\arcsin x = \frac{2t}{1+t^2}, \quad \cos x = \frac{1-t^2}{1+t^2}$$

$$= 2 \cdot \int_{-1}^1 \frac{1}{1+t^2} \dots dx = 2 \int_{-\infty}^{+\infty} \frac{1 - 2dt}{1-t^2+4t+3(1+t^2)}$$

$$= 4 \int_{-\infty}^{+\infty} \frac{dt}{2t^2+4t+4} =$$

$$= 2 \int_{-\infty}^{+\infty} \frac{dt}{(t+1)^2+1} =$$

$$= 2 \left[\arctan(t+1) \right]_{-\infty}^{+\infty} = 2\pi$$

$$\int_0^{\sqrt{1/2}} \frac{dx}{1+\sqrt{2}x}$$

$$t = \sqrt{2}x$$

$$\frac{dx}{dx} = \frac{1}{\sqrt{2}} = (1+\sqrt{2}^2x)$$

$$dx = \frac{dt}{1+t^2}$$

$$\int_0^{\sqrt{2}} \frac{dt}{(1+t)(1+t^2)} =$$

$$= \int_0^{\infty} \left[\frac{1}{2} \frac{1}{1+t} - \frac{1}{2} \frac{t-1}{t^2+1} \right] dt =$$

$$A(1+t^2) + (Bt+C)(1+t)$$

$$t=-1: A=\frac{1}{2} \quad | \quad t=i: 1=(B+iC)(1+i)$$

$$-3+iC = \frac{1}{2}$$

$$= \int_0^{\infty} \left[\frac{1}{2} \frac{1}{1+t} - \frac{1}{4} \frac{2t}{t^2+1} + \frac{1}{2} \frac{1}{t^2+1} \right] dt$$

$$= \left[\frac{1}{2} \ln \frac{(1+t)^2}{(t^2+1)} + \frac{1}{2} \arctan t \right]_0^{\infty} = \frac{1}{2} \ln 2$$

$$\int_{-\pi/2}^{\pi/2} \frac{1+\sin x}{1+\cos x} dx =$$

~~$$\int_{-\pi/2}^{\pi/2} \frac{1+\sin x - \cos x - \sin x \cos x}{\sin^2 x} dx$$~~

~~$$\int_{-\pi/2}^{\pi/2} \frac{1}{\sin^2 x} + \frac{1}{\sin x} - \frac{\cos x}{\sin^2 x} - \cos x dx$$~~

$$t = \tan \frac{x}{2}, \quad t \in (-1, 1)$$

~~$$\int_{-1}^1 \frac{1 + \frac{2t}{1+t^2}}{1 + \frac{1-t^2}{1+t^2}} \frac{2dt}{1+t^2} =$$~~

~~$$= \int_{-1}^1 \frac{t^2+1+2t}{(t^2+1)^2} \cdot \frac{2dt}{1+t^2} =$$~~

~~$$= \int_{-1}^1 \left(1 + \frac{2t}{t^2+1} \right) dt = \left[t + \ln(t^2+1) \right]_{-1}^1$$~~

~~$$= 1 + \ln 2 - (-1 + \ln 2) = \underline{2}$$~~