

Mathematics II - Matrices

23/24

`https:
//www.youtube.com/watch?v=zE7PKRjrid4`

`https:
//www.youtube.com/watch?v=zE7PKRjrid4`


Exercise




VI. Matrix calculus

							1	1	1	1	1									
				1	1	1	1	1	1	1	1									
			1	1			1					1	1							
		1	1											1	1					
		1	1			1		1						1	1					
		1	1			1		1						1	1					
		1	1											1	1					
			1	1		1	1	1						1	1					
			1	1			1							1	1					
			1	1										1	1					
			1	1	1									1	1	1				
	1	1	1	1										1	1	1				
1	1	1	1	1										1	1	1	1			
1	1	1	1	1										1	1	1	1			
1		1	1	1	1									1	1	1		1		
			1	1	1									1	1	1				
			1	1	1	1								1	1	1	1			
				1	1	1								1	1	1				
					1	1	1	1	1	1	1	1	1	1	1	1	1			

VI. Matrix calculus

							1	1	1	1	1							
				1	1	1	1	1	1	1	1							
		1	1			1						1	1					
1	1													1	1			
1	1			1		1								1	1			
1	1			1		1								1	1			
1	1													1	1			
		1	1			1	1	1				1	1					
		1	1			1						1	1					
		1	1											1	1			
	1	1	1	1										1	1	1		
1	1	1	1											1	1	1		
1	1	1	1											1	1	1	1	
1		1	1	1										1	1	1		1
			1	1	1									1	1	1		
				1	1	1								1	1	1		
					1	1	1							1	1	1		
		1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1





<https://www.pinterest.cl/pin/527273068861820414/>

VI.1. Basic operations with matrices

Definition

A table of numbers

$$\begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{pmatrix},$$

where $a_{ij} \in \mathbb{R}$, $i = 1, \dots, m$, $j = 1, \dots, n$, is called a **matrix of type $m \times n$** (shortly, an **m -by- n matrix**). We also write $(a_{ij})_{\substack{i=1..m \\ j=1..n}}$ for short.

An n -by- n matrix is called a **square matrix of order n** .

VI.1. Basic operations with matrices

Definition

A table of numbers

$$\begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{pmatrix},$$

where $a_{ij} \in \mathbb{R}$, $i = 1, \dots, m$, $j = 1, \dots, n$, is called a **matrix of type $m \times n$** (shortly, an **m -by- n matrix**). We also write $(a_{ij})_{\substack{i=1..m \\ j=1..n}}$ for short.

An n -by- n matrix is called a **square matrix of order n** .

The set of all m -by- n matrices is denoted by **$M(m \times n)$** .

Example

(3)

$$\begin{pmatrix} 6 \\ 7 \end{pmatrix} \begin{pmatrix} 2 & 2 & 8 \\ 5 & 0 & -2 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 2 \\ 4 & -4 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 \\ 0 & -3 & 1 \\ 4 & \pi & 3 \end{pmatrix}$$

Exercise

Find the type of the matrix

$$\begin{pmatrix} 6 & 11 & -2 \\ 23 & 31 & 5 \end{pmatrix}$$

A 2x3

B 3x2

C 6

Exercise

Find the type of the matrix

$$\begin{pmatrix} 6 & 11 & -2 \\ 23 & 31 & 5 \end{pmatrix}$$

A 2x3

B 3x2

C 6

A

Definition

Let

$$\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{pmatrix}.$$

The n -tuple $(a_{i1}, a_{i2}, \dots, a_{in})$, where $i \in \{1, 2, \dots, m\}$, is called the **i th row** of the matrix \mathbf{A} .

Definition

Let

$$A = \begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{pmatrix}.$$

The m -tuple $\begin{pmatrix} a_{1j} \\ a_{2j} \\ \vdots \\ a_{mj} \end{pmatrix}$, where $j \in \{1, 2, \dots, n\}$, is called the **j th column** of the matrix A .

Definition

We say that two matrices are **equal**, if they are of the same type and the corresponding elements are equal, i.e. if $\mathbf{A} = (a_{ij})_{\substack{i=1..m \\ j=1..n}}$ and $\mathbf{B} = (b_{uv})_{\substack{u=1..r \\ v=1..s}}$, then $\mathbf{A} = \mathbf{B}$ if and only if $m = r$, $n = s$ and $a_{ij} = b_{ij} \forall i \in \{1, \dots, m\}, \forall j \in \{1, \dots, n\}$.

Exercise

Are \mathbf{A} and \mathbf{B} equal?

$$\mathbf{A} = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 0 & 4 \\ -1 & -2 & 5 \end{pmatrix}$$

$$\mathbf{B} = \begin{pmatrix} 4 & 0 & 4 \\ 1 & 2 & 3 \\ -1 & -2 & 5 \end{pmatrix}$$

Definition

We say that two matrices are **equal**, if they are of the same type and the corresponding elements are equal, i.e. if $\mathbf{A} = (a_{ij})_{\substack{i=1..m \\ j=1..n}}$ and $\mathbf{B} = (b_{uv})_{\substack{u=1..r \\ v=1..s}}$, then $\mathbf{A} = \mathbf{B}$ if and only if $m = r$, $n = s$ and $a_{ij} = b_{ij} \forall i \in \{1, \dots, m\}, \forall j \in \{1, \dots, n\}$.

Exercise

Are \mathbf{A} and \mathbf{B} equal?

$$\mathbf{A} = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 0 & 4 \\ -1 & -2 & 5 \end{pmatrix}$$

$$\mathbf{B} = \begin{pmatrix} 4 & 0 & 4 \\ 1 & 2 & 3 \\ -1 & -2 & 5 \end{pmatrix}$$

No



Definition

Let $\mathbf{A}, \mathbf{B} \in M(m \times n)$, $\mathbf{A} = (a_{ij})_{\substack{i=1..m \\ j=1..n}}$, $\mathbf{B} = (b_{ij})_{\substack{i=1..m \\ j=1..n}}$, $\lambda \in \mathbb{R}$.

The **sum of the matrices \mathbf{A} and \mathbf{B}** is the matrix defined by

$$\mathbf{A} + \mathbf{B} = \begin{pmatrix} a_{11} + b_{11} & a_{12} + b_{12} & \dots & a_{1n} + b_{1n} \\ a_{21} + b_{21} & a_{22} + b_{22} & \dots & a_{2n} + b_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} + b_{m1} & a_{m2} + b_{m1} & \dots & a_{mn} + b_{mn} \end{pmatrix}.$$

Definition

Let $A, B \in M(m \times n)$, $A = (a_{ij})_{\substack{i=1..m \\ j=1..n}}$, $B = (b_{ij})_{\substack{i=1..m \\ j=1..n}}$, $\lambda \in \mathbb{R}$.

The **sum of the matrices A and B** is the matrix defined by

$$A + B = \begin{pmatrix} a_{11} + b_{11} & a_{12} + b_{12} & \dots & a_{1n} + b_{1n} \\ a_{21} + b_{21} & a_{22} + b_{22} & \dots & a_{2n} + b_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} + b_{m1} & a_{m2} + b_{m1} & \dots & a_{mn} + b_{mn} \end{pmatrix}.$$

The **product of the real number λ and the matrix A** (or the λ -multiple of the matrix A) is the matrix defined by

$$\lambda A = \begin{pmatrix} \lambda a_{11} & \lambda a_{12} & \dots & \lambda a_{1n} \\ \lambda a_{21} & \lambda a_{22} & \dots & \lambda a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ \lambda a_{m1} & \lambda a_{m2} & \dots & \lambda a_{mn} \end{pmatrix}.$$

Exercise

Let

$$A = \begin{pmatrix} 4 & 6 \\ 20 & 24 \end{pmatrix}$$

$$B = \begin{pmatrix} 2 & 5 \\ 3 & 7 \end{pmatrix}.$$

Find $A + B$

A 71

B

$$\begin{pmatrix} 6 & 9 \\ 7 & 11 \end{pmatrix}$$

C

$$\begin{pmatrix} 6 & 11 \\ 23 & 31 \end{pmatrix}$$

D

$$\begin{pmatrix} 26 & 62 \\ 112 & 268 \end{pmatrix}$$

E

$$\begin{pmatrix} 4 & 6 & 2 & 5 \\ 20 & 24 & 3 & 7 \end{pmatrix}$$

Exercise

Let

$$A = \begin{pmatrix} 4 & 6 \\ 20 & 24 \end{pmatrix}$$

$$B = \begin{pmatrix} 2 & 5 \\ 3 & 7 \end{pmatrix}.$$

Find $A + B$

A 71

B

$$\begin{pmatrix} 6 & 9 \\ 7 & 11 \end{pmatrix}$$

C

$$\begin{pmatrix} 6 & 11 \\ 23 & 31 \end{pmatrix}$$

D

$$\begin{pmatrix} 26 & 62 \\ 112 & 268 \end{pmatrix}$$

E

$$\begin{pmatrix} 4 & 6 & 2 & 5 \\ 20 & 24 & 3 & 7 \end{pmatrix}$$

C

Exercise

Let

$$A = \begin{pmatrix} 4 & 6 \\ 20 & 7 \end{pmatrix}$$

Find $5A$

A

$$\begin{pmatrix} 9 & 6 \\ 20 & 7 \end{pmatrix}$$

C

$$\begin{pmatrix} 20 & 6 \\ 20 & 7 \end{pmatrix}$$

B

$$\begin{pmatrix} 9 & 11 \\ 25 & 12 \end{pmatrix}$$

D

$$\begin{pmatrix} 20 & 30 \\ 100 & 35 \end{pmatrix}$$

Exercise

Let

$$A = \begin{pmatrix} 4 & 6 \\ 20 & 7 \end{pmatrix}$$

Find $5A$

A

$$\begin{pmatrix} 9 & 6 \\ 20 & 7 \end{pmatrix}$$

C

$$\begin{pmatrix} 20 & 6 \\ 20 & 7 \end{pmatrix}$$

B

$$\begin{pmatrix} 9 & 11 \\ 25 & 12 \end{pmatrix}$$

D

$$\begin{pmatrix} 20 & 30 \\ 100 & 35 \end{pmatrix}$$

D

Exercise

Let $\mathbf{A} = \begin{pmatrix} 1 & -2 \\ 2 & 4 \end{pmatrix}$, $\mathbf{B} = \begin{pmatrix} 3 & 1 \\ -2 & 0 \end{pmatrix}$, $\mathbf{C} = \begin{pmatrix} -4 & 0 \\ -2 & 1 \end{pmatrix}$ and
 $\mathbf{O} = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$.

1. Find $(\mathbf{A} + \mathbf{B}) + \mathbf{C}$ and $\mathbf{A} + (\mathbf{B} + \mathbf{C})$.
2. Find $\mathbf{A} + \mathbf{B}$ and $\mathbf{B} + \mathbf{A}$.
3. Find $\mathbf{A} + \mathbf{O}$ and $\mathbf{O} + \mathbf{A}$.
4. Find a matrix \mathbf{C}_A such that $\mathbf{A} + \mathbf{C}_A = \mathbf{O}$.
5. Find $2 \cdot 3\mathbf{A}$ and $2(3\mathbf{A})$.
6. Find $2 \cdot 3\mathbf{A}$ and $2(3\mathbf{A})$.
7. Find $(1 + 2)\mathbf{A}$ and $1\mathbf{A} + 2\mathbf{A}$.
8. Find $2\mathbf{A} + 2\mathbf{B}$ and $2(\mathbf{A} + \mathbf{B})$.

Proposition 1 (basic properties of the sum of matrices and of a multiplication by a scalar)

The following holds:

- $\forall \mathbf{A}, \mathbf{B}, \mathbf{C} \in M(m \times n): \mathbf{A} + (\mathbf{B} + \mathbf{C}) = (\mathbf{A} + \mathbf{B}) + \mathbf{C}$,
(associativity)
- $\forall \mathbf{A}, \mathbf{B} \in M(m \times n): \mathbf{A} + \mathbf{B} = \mathbf{B} + \mathbf{A}$, (commutativity)
- $\exists! \mathbf{O} \in M(m \times n) \forall \mathbf{A} \in M(m \times n): \mathbf{A} + \mathbf{O} = \mathbf{A}$, (existence of a zero element)
- $\forall \mathbf{A} \in M(m \times n) \exists \mathbf{C}_A \in M(m \times n): \mathbf{A} + \mathbf{C}_A = \mathbf{O}$, (existence of an opposite element)
- $\forall \mathbf{A} \in M(m \times n) \forall \lambda, \mu \in \mathbb{R}: (\lambda\mu)\mathbf{A} = \lambda(\mu\mathbf{A})$,
- $\forall \mathbf{A} \in M(m \times n): 1 \cdot \mathbf{A} = \mathbf{A}$,
- $\forall \mathbf{A} \in M(m \times n) \forall \lambda, \mu \in \mathbb{R}: (\lambda + \mu)\mathbf{A} = \lambda\mathbf{A} + \mu\mathbf{A}$,
- $\forall \mathbf{A}, \mathbf{B} \in M(m \times n) \forall \lambda \in \mathbb{R}: \lambda(\mathbf{A} + \mathbf{B}) = \lambda\mathbf{A} + \lambda\mathbf{B}$.

Remark

- The matrix \mathbf{O} from the previous proposition is called a **zero matrix** and all its elements are all zeros.


Remark

- The matrix \mathbf{O} from the previous proposition is called a **zero matrix** and all its elements are all zeros.
- The matrix \mathbf{C}_A from the previous proposition is called a **matrix opposite to \mathbf{A}** . It is determined uniquely, it is denoted by $-\mathbf{A}$, and it satisfies $-\mathbf{A} = (-a_{ij})_{\substack{i=1..m \\ j=1..n}}$ and $-\mathbf{A} = -1 \cdot \mathbf{A}$.

Definition

Let $A \in M(m \times n)$, $A = (a_{is})_{\substack{i=1..m, \\ s=1..n}}$, $B \in M(n \times k)$,
 $B = (b_{sj})_{\substack{s=1..n, \\ j=1..k}}$. Then the **product of matrices** A and B is defined
as a matrix $AB \in M(m \times k)$, $AB = (c_{ij})_{\substack{i=1..m, \\ j=1..k}}$, where

$$c_{ij} = \sum_{s=1}^n a_{is}b_{sj}.$$



Matrix multiplication

$$\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \\ a_{31} & a_{32} \\ a_{41} & a_{42} \end{pmatrix} \cdot \begin{pmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \end{pmatrix}$$

Matrix multiplication

$$\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \\ a_{31} & a_{32} \\ a_{41} & a_{42} \end{pmatrix} \cdot \begin{pmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \end{pmatrix} \\ = \begin{pmatrix} a_{11}b_{11} + a_{12}b_{21} & a_{11}b_{12} + a_{12}b_{22} & a_{11}b_{13} + a_{12}b_{23} \\ a_{21}b_{11} + a_{22}b_{21} & a_{21}b_{12} + a_{22}b_{22} & a_{21}b_{13} + a_{22}b_{23} \\ a_{31}b_{11} + a_{32}b_{21} & a_{31}b_{12} + a_{32}b_{22} & a_{31}b_{13} + a_{32}b_{23} \\ a_{41}b_{11} + a_{42}b_{21} & a_{41}b_{12} + a_{42}b_{22} & a_{41}b_{13} + a_{42}b_{23} \end{pmatrix}$$

Matrix multiplication

$$\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \\ a_{31} & a_{32} \\ a_{41} & a_{42} \end{pmatrix} \cdot \begin{pmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \end{pmatrix}$$
$$= \begin{pmatrix} a_{11}b_{11} + a_{12}b_{21} & a_{11}b_{12} + a_{12}b_{22} & a_{11}b_{13} + a_{12}b_{23} \\ a_{21}b_{11} + a_{22}b_{21} & a_{21}b_{12} + a_{22}b_{22} & a_{21}b_{13} + a_{22}b_{23} \\ a_{31}b_{11} + a_{32}b_{21} & a_{31}b_{12} + a_{32}b_{22} & a_{31}b_{13} + a_{32}b_{23} \\ a_{41}b_{11} + a_{42}b_{21} & a_{41}b_{12} + a_{42}b_{22} & a_{41}b_{13} + a_{42}b_{23} \end{pmatrix}$$

Matrix multiplication

$$\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \\ a_{31} & a_{32} \\ a_{41} & a_{42} \end{pmatrix} \cdot \begin{pmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \end{pmatrix}$$
$$= \begin{pmatrix} a_{11}b_{11} + a_{12}b_{21} & a_{11}b_{12} + a_{12}b_{22} & a_{11}b_{13} + a_{12}b_{23} \\ a_{21}b_{11} + a_{22}b_{21} & a_{21}b_{12} + a_{22}b_{22} & a_{21}b_{13} + a_{22}b_{23} \\ a_{31}b_{11} + a_{32}b_{21} & a_{31}b_{12} + a_{32}b_{22} & a_{31}b_{13} + a_{32}b_{23} \\ a_{41}b_{11} + a_{42}b_{21} & a_{41}b_{12} + a_{42}b_{22} & a_{41}b_{13} + a_{42}b_{23} \end{pmatrix}$$

Matrix multiplication

$$\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \\ a_{31} & a_{32} \\ a_{41} & a_{42} \end{pmatrix} \cdot \begin{pmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \end{pmatrix}$$
$$= \begin{pmatrix} a_{11}b_{11} + a_{12}b_{21} & a_{11}b_{12} + a_{12}b_{22} & a_{11}b_{13} + a_{12}b_{23} \\ a_{21}b_{11} + a_{22}b_{21} & a_{21}b_{12} + a_{22}b_{22} & a_{21}b_{13} + a_{22}b_{23} \\ a_{31}b_{11} + a_{32}b_{21} & a_{31}b_{12} + a_{32}b_{22} & a_{31}b_{13} + a_{32}b_{23} \\ a_{41}b_{11} + a_{42}b_{21} & a_{41}b_{12} + a_{42}b_{22} & a_{41}b_{13} + a_{42}b_{23} \end{pmatrix}$$

Matrix multiplication

$$\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \\ a_{31} & a_{32} \\ a_{41} & a_{42} \end{pmatrix} \cdot \begin{pmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \end{pmatrix}$$
$$= \begin{pmatrix} a_{11}b_{11} + a_{12}b_{21} & a_{11}b_{12} + a_{12}b_{22} & a_{11}b_{13} + a_{12}b_{23} \\ a_{21}b_{11} + a_{22}b_{21} & a_{21}b_{12} + a_{22}b_{22} & a_{21}b_{13} + a_{22}b_{23} \\ a_{31}b_{11} + a_{32}b_{21} & a_{31}b_{12} + a_{32}b_{22} & a_{31}b_{13} + a_{32}b_{23} \\ a_{41}b_{11} + a_{42}b_{21} & a_{41}b_{12} + a_{42}b_{22} & a_{41}b_{13} + a_{42}b_{23} \end{pmatrix}$$

Exercise

Find \mathbf{AB} , if

$$\mathbf{A} = \begin{pmatrix} 2 & 0 \\ -3 & 1 \end{pmatrix}$$

$$\mathbf{B} = \begin{pmatrix} 0 & -1 \\ 2 & 2 \end{pmatrix}$$

A $\begin{pmatrix} 3 & -1 \\ -2 & 2 \end{pmatrix}$

C $\begin{pmatrix} 0 & 0 \\ -6 & 2 \end{pmatrix}$

B $\begin{pmatrix} 0 & -2 \\ 2 & 5 \end{pmatrix}$

D something else

E \mathbf{AB} is not well defined

Exercise

Find \mathbf{AB} , if

$$\mathbf{A} = \begin{pmatrix} 2 & 0 \\ -3 & 1 \end{pmatrix}$$

$$\mathbf{B} = \begin{pmatrix} 0 & -1 \\ 2 & 2 \end{pmatrix}$$

A $\begin{pmatrix} 3 & -1 \\ -2 & 2 \end{pmatrix}$

C $\begin{pmatrix} 0 & 0 \\ -6 & 2 \end{pmatrix}$

B $\begin{pmatrix} 0 & -2 \\ 2 & 5 \end{pmatrix}$

D something else

E \mathbf{AB} is not well defined

B

Exercise

Find \mathbf{AB} , if

$$\mathbf{A} = \begin{pmatrix} 2 & 1 \\ 3 & 2 \end{pmatrix}$$

$$\mathbf{B} = \begin{pmatrix} 4 \\ -1 \end{pmatrix}$$

A $\begin{pmatrix} 5 \\ 2 \end{pmatrix}$

B $\begin{pmatrix} 10 & 7 \end{pmatrix}$

C $\begin{pmatrix} 8 & 4 \\ -3 & -2 \end{pmatrix}$

D $\begin{pmatrix} 7 \\ 10 \end{pmatrix}$

E \mathbf{AB} is not well defined

Exercise

Find \mathbf{AB} , if

$$\mathbf{A} = \begin{pmatrix} 2 & 1 \\ 3 & 2 \end{pmatrix}$$

$$\mathbf{B} = \begin{pmatrix} 4 \\ -1 \end{pmatrix}$$

A $\begin{pmatrix} 5 \\ 2 \end{pmatrix}$

B $\begin{pmatrix} 10 & 7 \end{pmatrix}$

C $\begin{pmatrix} 8 & 4 \\ -3 & -2 \end{pmatrix}$

D $\begin{pmatrix} 7 \\ 10 \end{pmatrix}$

E \mathbf{AB} is not well defined

D

Multiplication properties

Exercise

Let

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

$$I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

Find

1. AI

2. IA

Multiplication properties

Exercise

Let

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

$$I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

Find

1. AI

2. IA

Exercise

Let

$$A = \begin{pmatrix} 2 & -1 \\ 1 & 4 \end{pmatrix}$$

$$B = \begin{pmatrix} 0 & 0 \\ 3 & -3 \end{pmatrix}$$

Find

1. AB

2. BA



Theorem 2 (properties of the matrix multiplication)

Let $m, n, k, l \in \mathbb{N}$. Then:

- (i) $\forall \mathbf{A} \in M(m \times n) \forall \mathbf{B} \in M(n \times k) \forall \mathbf{C} \in M(k \times l)$:
$$\mathbf{A}(\mathbf{BC}) = (\mathbf{AB})\mathbf{C}, \quad (\text{associativity of multiplication})$$
- (ii) $\forall \mathbf{A} \in M(m \times n) \forall \mathbf{B}, \mathbf{C} \in M(n \times k)$:
$$\mathbf{A}(\mathbf{B} + \mathbf{C}) = \mathbf{AB} + \mathbf{AC}, \quad (\text{distributivity from the left})$$
- (iii) $\forall \mathbf{A}, \mathbf{B} \in M(m \times n) \forall \mathbf{C} \in M(n \times k)$:
$$(\mathbf{A} + \mathbf{B})\mathbf{C} = \mathbf{AC} + \mathbf{BC}, \quad (\text{distributivity from the right})$$
- (iv) $\exists ! \mathbf{I} \in M(n \times n) \forall \mathbf{A} \in M(n \times n): \mathbf{IA} = \mathbf{AI} = \mathbf{A}$. (existence and uniqueness of an *identity matrix* \mathbf{I})

Remark

Warning! The matrix multiplication is not commutative.

Definition

A **transpose** of a matrix

$$\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & a_{m3} & \cdots & a_{mn} \end{pmatrix}$$

is the matrix

$$\mathbf{A}^T = \begin{pmatrix} a_{11} & a_{21} & \cdots & a_{m1} \\ a_{12} & a_{22} & \cdots & a_{m2} \\ a_{13} & a_{23} & \cdots & a_{m3} \\ \vdots & \vdots & \ddots & \vdots \\ a_{1n} & a_{2n} & \cdots & a_{mn} \end{pmatrix},$$

i.e. if $\mathbf{A} = (a_{ij})_{\substack{i=1..m \\ j=1..n}}$, then $\mathbf{A}^T = (b_{uv})_{\substack{u=1..n \\ v=1..m}}$, where $b_{uv} = a_{vu}$ for each $u \in \{1, \dots, n\}$, $v \in \{1, 2, \dots, m\}$.

Definition

A **transpose** of a matrix

$$\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & a_{m3} & \cdots & a_{mn} \end{pmatrix}$$

is the matrix

$$\mathbf{A}^T = \begin{pmatrix} a_{11} & a_{21} & \cdots & a_{m1} \\ a_{12} & a_{22} & \cdots & a_{m2} \\ a_{13} & a_{23} & \cdots & a_{m3} \\ \vdots & \vdots & \ddots & \vdots \\ a_{1n} & a_{2n} & \cdots & a_{mn} \end{pmatrix},$$

i.e. if $\mathbf{A} = (a_{ij})_{\substack{i=1..m \\ j=1..n}}$, then $\mathbf{A}^T = (b_{uv})_{\substack{u=1..n \\ v=1..m}}$, where $b_{uv} = a_{vu}$ for each $u \in \{1, \dots, n\}$, $v \in \{1, 2, \dots, m\}$.

Definition

A **transpose** of a matrix

$$\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & a_{m3} & \dots & a_{mn} \end{pmatrix}$$

is the matrix

$$\mathbf{A}^T = \begin{pmatrix} a_{11} & a_{21} & \dots & a_{m1} \\ a_{12} & a_{22} & \dots & a_{m2} \\ a_{13} & a_{23} & \dots & a_{m3} \\ \vdots & \vdots & \ddots & \vdots \\ a_{1n} & a_{2n} & \dots & a_{mn} \end{pmatrix},$$

i.e. if $\mathbf{A} = (a_{ij})_{\substack{i=1..m \\ j=1..n}}$, then $\mathbf{A}^T = (b_{uv})_{\substack{u=1..n \\ v=1..m}}$, where $b_{uv} = a_{vu}$ for each $u \in \{1, \dots, n\}$, $v \in \{1, 2, \dots, m\}$.

Definition

A **transpose** of a matrix

$$\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & a_{m3} & \dots & a_{mn} \end{pmatrix}$$

is the matrix

$$\mathbf{A}^T = \begin{pmatrix} a_{11} & a_{21} & \dots & a_{m1} \\ a_{12} & a_{22} & \dots & a_{m2} \\ a_{13} & a_{23} & \dots & a_{m3} \\ \vdots & \vdots & \ddots & \vdots \\ a_{1n} & a_{2n} & \dots & a_{mn} \end{pmatrix},$$

i.e. if $\mathbf{A} = (a_{ij})_{\substack{i=1..m \\ j=1..n}}$, then $\mathbf{A}^T = (b_{uv})_{\substack{u=1..n \\ v=1..m}}$, where $b_{uv} = a_{vu}$ for each $u \in \{1, \dots, n\}$, $v \in \{1, 2, \dots, m\}$.

Definition

A **transpose** of a matrix

$$\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & a_{m3} & \cdots & a_{mn} \end{pmatrix}$$

is the matrix

$$\mathbf{A}^T = \begin{pmatrix} a_{11} & a_{21} & \cdots & a_{m1} \\ a_{12} & a_{22} & \cdots & a_{m2} \\ a_{13} & a_{23} & \cdots & a_{m3} \\ \vdots & \vdots & \ddots & \vdots \\ a_{1n} & a_{2n} & \cdots & a_{mn} \end{pmatrix},$$

i.e. if $\mathbf{A} = (a_{ij})_{\substack{i=1..m \\ j=1..n}}$, then $\mathbf{A}^T = (b_{uv})_{\substack{u=1..n \\ v=1..m}}$, where $b_{uv} = a_{vu}$ for each $u \in \{1, \dots, n\}$, $v \in \{1, 2, \dots, m\}$.

Example

$$\mathbf{D} = \begin{pmatrix} 1 & 2 \\ 4 & -4 \\ 2 & 3 \end{pmatrix}$$

$$\mathbf{D}^T = \begin{pmatrix} 1 & 4 & 2 \\ 2 & -4 & 3 \end{pmatrix}$$

$$\mathbf{F} = \begin{pmatrix} 1 & 2 & 3 \\ 0 & -3 & 1 \\ 4 & \pi & 3 \end{pmatrix}$$

$$\mathbf{F}^T = \begin{pmatrix} 1 & 0 & 4 \\ 2 & -3 & \pi \\ 3 & 1 & 3 \end{pmatrix}$$

Exercise

$$\mathbf{A} = \begin{pmatrix} 2 & 3 & 1 \\ 0 & -1 & 3 \\ -2 & 0 & 4 \end{pmatrix}.$$

Find \mathbf{A}^T ?

A

$$A^T = \begin{pmatrix} 2 & 3 & 1 \\ 0 & -1 & 3 \\ -2 & 0 & 4 \end{pmatrix}$$

C

$$A^T = \begin{pmatrix} -2 & 0 & 4 \\ 0 & -1 & 3 \\ 2 & 3 & 1 \end{pmatrix}$$

B

$$A^T = \begin{pmatrix} 2 & 0 & -2 \\ 3 & -1 & 0 \\ 1 & 3 & 4 \end{pmatrix}$$

D

$$A^T = \begin{pmatrix} 1 & 3 & 4 \\ 3 & -1 & 0 \\ 2 & 0 & -2 \end{pmatrix}$$

Exercise

$$\mathbf{A} = \begin{pmatrix} 2 & 3 & 1 \\ 0 & -1 & 3 \\ -2 & 0 & 4 \end{pmatrix}.$$

Find \mathbf{A}^T ?

A

$$A^T = \begin{pmatrix} 2 & 3 & 1 \\ 0 & -1 & 3 \\ -2 & 0 & 4 \end{pmatrix}$$

C

$$A^T = \begin{pmatrix} -2 & 0 & 4 \\ 0 & -1 & 3 \\ 2 & 3 & 1 \end{pmatrix}$$

B

$$A^T = \begin{pmatrix} 2 & 0 & -2 \\ 3 & -1 & 0 \\ 1 & 3 & 4 \end{pmatrix}$$

D

$$A^T = \begin{pmatrix} 1 & 3 & 4 \\ 3 & -1 & 0 \\ 2 & 0 & -2 \end{pmatrix}$$

B



Exercise

Let

$$A = \begin{pmatrix} 1 & -1 \\ 0 & 2 \end{pmatrix}$$

$$B = \begin{pmatrix} 3 & 0 \\ 5 & -1 \end{pmatrix}$$

Find

1. $(AB)^T$

2. $A^T B^T$

3. $B^T A^T$

Theorem 3 (properties of the transpose of a matrix)

Platí:

- (i) $\forall \mathbf{A} \in M(m \times n): (\mathbf{A}^T)^T = \mathbf{A},$
- (ii) $\forall \mathbf{A}, \mathbf{B} \in M(m \times n): (\mathbf{A} + \mathbf{B})^T = \mathbf{A}^T + \mathbf{B}^T,$
- (iii) $\forall \mathbf{A} \in M(m \times n) \forall \mathbf{B} \in M(n \times k): (\mathbf{AB})^T = \mathbf{B}^T \mathbf{A}^T.$

Theorem 3 (properties of the transpose of a matrix)

Plati:

- (i) $\forall \mathbf{A} \in M(m \times n): (\mathbf{A}^T)^T = \mathbf{A},$
- (ii) $\forall \mathbf{A}, \mathbf{B} \in M(m \times n): (\mathbf{A} + \mathbf{B})^T = \mathbf{A}^T + \mathbf{B}^T,$
- (iii) $\forall \mathbf{A} \in M(m \times n) \forall \mathbf{B} \in M(n \times k): (\mathbf{AB})^T = \mathbf{B}^T \mathbf{A}^T.$

Definition

We say that the matrix $\mathbf{A} \in M(n \times n)$ is **symmetric** if $\mathbf{A} = \mathbf{A}^T$.

Exercise

Let \mathbf{A} and \mathbf{B} are matrices of the type 2×3 . Which of these operations are NOT well defined?

A $\mathbf{A} + \mathbf{B}$

B $\mathbf{A}^T \mathbf{B}$

C \mathbf{BA}

D \mathbf{AB}^T

E \mathbf{AB}

Exercise

Let \mathbf{A} and \mathbf{B} are matrices of the type 2×3 . Which of these operations are NOT well defined?

A $\mathbf{A} + \mathbf{B}$

B $\mathbf{A}^T \mathbf{B}$

C \mathbf{BA}

D \mathbf{AB}^T

E \mathbf{AB}

C, E

Exercise

We want to multiply matrices $\mathbf{A} \times \mathbf{B}$. We need:

- A \mathbf{A} and \mathbf{B} needs to have the same number of rows.
- B \mathbf{A} and \mathbf{B} needs to have the same number of columns.
- C the number of rows of \mathbf{A} needs to be the same as the number of columns of \mathbf{B}
- D the number of columns of \mathbf{A} needs to be the same as the number of rows of \mathbf{B}

Exercise

We want to multiply matrices $\mathbf{A} \times \mathbf{B}$. We need:

- A \mathbf{A} and \mathbf{B} needs to have the same number of rows.
- B \mathbf{A} and \mathbf{B} needs to have the same number of columns.
- C the number of rows of \mathbf{A} needs to be the same as the number of columns of \mathbf{B}
- D the number of columns of \mathbf{A} needs to be the same as the number of rows of \mathbf{B}

D

Exercise

Let \mathbf{A} is a matrix of the type 2×3 and \mathbf{B} is of the type 3×6 .
Find the type of \mathbf{AB} :

A 2×6

C 3×3

E 3×6

B 6×2

D 2×3

Exercise

Let **A** is a matrix of the type 2×3 and **B** is of the type 3×6 .
Find the type of **AB**:

A 2×6

C 3×3

E 3×6

B 6×2

D 2×3

A

Exercise

Let **A** is a matrix of the type 2×3 and **B** is of the type 3×6 .
Find the type of **AB**:

A 2×6

C 3×3

E 3×6

B 6×2

D 2×3

A

Exercise (True or False?)

Let **A** and **B** be square matrices of the same dimension. Then

$$(\mathbf{A} + \mathbf{B}) \times (\mathbf{A} + \mathbf{B}) = \mathbf{A}^2 + 2\mathbf{AB} + \mathbf{B}^2.$$

Exercise

Let **A** is a matrix of the type 2×3 and **B** is of the type 3×6 .
Find the type of **AB**:

A 2×6

C 3×3

E 3×6

B 6×2

D 2×3

A

Exercise (True or False?)

Let **A** and **B** be square matrices of the same dimension. Then

$$(\mathbf{A} + \mathbf{B}) \times (\mathbf{A} + \mathbf{B}) = \mathbf{A}^2 + 2\mathbf{AB} + \mathbf{B}^2.$$

False

$$\begin{aligned}(\mathbf{A} + \mathbf{B}) \times (\mathbf{A} + \mathbf{B}) &= \mathbf{A}^2 + \mathbf{AB} + \mathbf{BA} + \mathbf{B}^2 \\ &\neq \mathbf{A}^2 + \mathbf{AB} + \mathbf{AB} + \mathbf{B}^2 = \mathbf{A}^2 + 2\mathbf{AB} + \mathbf{B}^2.\end{aligned}$$

Exercise

Visit: <https://web.ma.utexas.edu/users/ysulyma/matrix/>

[//web.ma.utexas.edu/users/ysulyma/matrix/](https://web.ma.utexas.edu/users/ysulyma/matrix/)

(You can change the picture:

https://karlin.mff.cuni.cz/~kuncova/en/2122LS_FMat2/bilyctverec.jpg.)

Try the following matrices:

1. 1.1 $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$

1.2 $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$

1.3 $\begin{pmatrix} 0.71 & -0.71 \\ 0.71 & 0.71 \end{pmatrix}$

2. 2.1 $\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$

2.2 $\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$

2.3 $\begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$

3. 3.1 $\begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}$

3.2 $\begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix}$

3.3 $\begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}$

Exercise

And what about these matrices? What is the result of matrix multiplying?

$$1. \quad 1.1 \quad \begin{pmatrix} 0 & 1 \\ -2 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix} \cdot \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$$

$$1.2 \quad \begin{pmatrix} 0 & 2 \\ -1 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \cdot \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}$$

Cheat sheet for matrix transform:

https://en.wikipedia.org/wiki/File:2D_affine_transformation_matrix.svg

VI.2. Invertible matrices

Definition

Let $A \in M(n \times n)$. We say that A is an **invertible** matrix if there exists $B \in M(n \times n)$ such that

$$AB = BA = I.$$

Definition

We say that the matrix $B \in M(n \times n)$ is an **inverse** of a matrix $A \in M(n \times n)$ if $AB = BA = I$.

Remark

A matrix $A \in M(n \times n)$ is invertible if and only if it has an inverse.

Exercise

Let

$$\mathbf{A} = \begin{pmatrix} 0 & 4 \\ 2 & 0 \end{pmatrix}$$

Find \mathbf{A}^{-1}

A

$$\begin{pmatrix} 0 & 4 \\ 2 & 0 \end{pmatrix}$$

C

$$\begin{pmatrix} 0 & 1/4 \\ 1/2 & 0 \end{pmatrix}$$

B

$$\begin{pmatrix} 4 & 0 \\ 0 & 2 \end{pmatrix}$$

D

$$\begin{pmatrix} 0 & 1/2 \\ 1/4 & 0 \end{pmatrix}$$

Exercise

Let

$$\mathbf{A} = \begin{pmatrix} 0 & 4 \\ 2 & 0 \end{pmatrix}$$

Find \mathbf{A}^{-1}

A

$$\begin{pmatrix} 0 & 4 \\ 2 & 0 \end{pmatrix}$$

C

$$\begin{pmatrix} 0 & 1/4 \\ 1/2 & 0 \end{pmatrix}$$

B

$$\begin{pmatrix} 4 & 0 \\ 0 & 2 \end{pmatrix}$$

D

$$\begin{pmatrix} 0 & 1/2 \\ 1/4 & 0 \end{pmatrix}$$

D

Exercise

Find inverse for

$$\mathbf{A} = \begin{pmatrix} 1 & -3 \\ -2 & 0 \end{pmatrix}$$

Exercise

Find inverse for

$$\mathbf{A} = \begin{pmatrix} 1 & -3 \\ -2 & 0 \end{pmatrix}$$

$$\mathbf{A}^{-1} = \begin{pmatrix} 0 & -\frac{1}{2} \\ -\frac{1}{3} & -\frac{1}{6} \end{pmatrix}$$

Remark

- If $A \in M(n \times n)$ is invertible, then it has exactly one inverse, which is denoted by A^{-1} .

Remark

- If $\mathbf{A} \in M(n \times n)$ is invertible, then it has exactly one inverse, which is denoted by \mathbf{A}^{-1} .
- If some matrices $\mathbf{A}, \mathbf{B} \in M(n \times n)$ satisfy $\mathbf{AB} = \mathbf{I}$, then also $\mathbf{BA} = \mathbf{I}$.

Remark

- If $\mathbf{A} \in M(n \times n)$ is invertible, then it has exactly one inverse, which is denoted by \mathbf{A}^{-1} .
- If some matrices $\mathbf{A}, \mathbf{B} \in M(n \times n)$ satisfy $\mathbf{AB} = \mathbf{I}$, then also $\mathbf{BA} = \mathbf{I}$.

Theorem 4 (operations with invertible matrices)

Let $\mathbf{A}, \mathbf{B} \in M(n \times n)$ be invertible matrices. Then

- \mathbf{A}^{-1} is invertible and $(\mathbf{A}^{-1})^{-1} = \mathbf{A}$,
- \mathbf{A}^T is invertible and $(\mathbf{A}^T)^{-1} = (\mathbf{A}^{-1})^T$,
- \mathbf{AB} is invertible and $(\mathbf{AB})^{-1} = \mathbf{B}^{-1}\mathbf{A}^{-1}$.

Definition 1

The *Determinant* of the matrix \mathbf{A} of type $(1, 1)$ is equal to

$$\det \mathbf{A} = a_{1,1}.$$

The *Determinant* of the matrix \mathbf{A} of type $(2, 2)$ is equal to

$$\det \mathbf{A} = a_{1,1} \cdot a_{2,2} - a_{1,2} \cdot a_{2,1}.$$




Figure: <http://umv.science.upjs.sk/madaras/MZ1a/>

Exercise

Find the determinant of

$$\begin{pmatrix} 5 & 4 \\ 1 & 3 \end{pmatrix}$$

A 4

B 11

C 15

D 19

Exercise

Find the determinant of

$$\begin{pmatrix} 5 & 4 \\ 1 & 3 \end{pmatrix}$$

A 4

B 11

C 15

D 19

B

Definition 2 (Sarrus)

The *Determinant* of the matrix \mathbf{A} of type $(3, 3)$ is equal to

$$\det \mathbf{A} = a_{1,1}a_{2,2}a_{3,3} + a_{1,2}a_{2,3}a_{3,1} + a_{1,3}a_{2,1}a_{3,2} \\ - a_{1,3}a_{2,2}a_{3,1} - a_{1,2}a_{2,1}a_{3,3} - a_{1,1}a_{2,3}a_{3,2}.$$




Figure: https://de.wikipedia.org/wiki/Regel_von_Sarrus

Exercise

Find the determinant of

$$\begin{pmatrix} 5 & 2 & -1 \\ 0 & 3 & 4 \\ 0 & 0 & 1 \end{pmatrix}$$

A 0

B 6

C 15

D 22

Exercise

Find the determinant of

$$\begin{pmatrix} 5 & 2 & -1 \\ 0 & 3 & 4 \\ 0 & 0 & 1 \end{pmatrix}$$

A 0

B 6

C 15

D 22

C

Definition 3

Let \mathbf{A} be square matrix of the type (n, n) . Let $\mathbf{A}_{i,j}$ denote the matrix of type $(n-1, n-1)$, which is created from \mathbf{A} by omitting the i th row and j th column. Let $r \in \{1, \dots, n\}$. Then the determinant of \mathbf{A} is equal to

$$\det \mathbf{A} = a_{r,1}(-1)^{r+1} \det \mathbf{A}_{r,1} + a_{r,2}(-1)^{r+2} \det \mathbf{A}_{r,2} + \dots + a_{r,n}(-1)^{r+n} \det \mathbf{A}_{r,n}.$$




Figure:

<http://mathcentral.uregina.ca/OO/database/OO.09.06/h/suud1.html>

VI.3. Determinants

Definition

Let $A \in M(n \times n)$. The symbol A_{ij} denotes the $(n - 1)$ -by- $(n - 1)$ matrix which is created from A by omitting the i th row and the j th column.

VI.3. Determinants

Definition

Let $\mathbf{A} \in M(n \times n)$. The symbol \mathbf{A}_{ij} denotes the $(n - 1)$ -by- $(n - 1)$ matrix which is created from \mathbf{A} by omitting the i th row and the j th column.

$$\mathbf{A} = \begin{pmatrix} a_{1,1} & \cdots & a_{1,j-1} & a_{1,j} & a_{1,j+1} & \cdots & a_{1,n} \\ \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{i-1,1} & \cdots & a_{i-1,j-1} & a_{i-1,j} & a_{i-1,j+1} & \cdots & a_{i-1,n} \\ a_{i,1} & \cdots & a_{i,j-1} & a_{i,j} & a_{i,j+1} & \cdots & a_{i,n} \\ a_{i+1,1} & \cdots & a_{i+1,j-1} & a_{i+1,j} & a_{i+1,j+1} & \cdots & a_{i+1,n} \\ \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{n,1} & \cdots & a_{n,j-1} & a_{n,j} & a_{n,j+1} & \cdots & a_{n,n} \end{pmatrix}$$

VI.3. Determinants

Definition

Let $A \in M(n \times n)$. The symbol A_{ij} denotes the $(n - 1)$ -by- $(n - 1)$ matrix which is created from A by omitting the i th row and the j th column.

$$A = \begin{pmatrix} a_{1,1} & \cdots & a_{1,j-1} & a_{1,j} & a_{1,j+1} & \cdots & a_{1,n} \\ \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{i-1,1} & \cdots & a_{i-1,j-1} & a_{i-1,j} & a_{i-1,j+1} & \cdots & a_{i-1,n} \\ a_{i,1} & \cdots & a_{i,j-1} & a_{i,j} & a_{i,j+1} & \cdots & a_{i,n} \\ a_{i+1,1} & \cdots & a_{i+1,j-1} & a_{i+1,j} & a_{i+1,j+1} & \cdots & a_{i+1,n} \\ \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{n,1} & \cdots & a_{n,j-1} & a_{n,j} & a_{n,j+1} & \cdots & a_{n,n} \end{pmatrix}$$

VI.3. Determinants

Definition

Let $A \in M(n \times n)$. The symbol A_{ij} denotes the $(n - 1)$ -by- $(n - 1)$ matrix which is created from A by omitting the i th row and the j th column.

$$\begin{pmatrix} a_{1,1} & \cdots & a_{1,j-1} & & a_{1,j+1} & \cdots & a_{1,n} \\ \vdots & \ddots & \vdots & & \vdots & \ddots & \vdots \\ a_{i-1,1} & \cdots & a_{i-1,j-1} & & a_{i-1,j+1} & \cdots & a_{i-1,n} \\ & & & & & & \\ a_{i+1,1} & \cdots & a_{i+1,j-1} & & a_{i+1,j+1} & \cdots & a_{i+1,n} \\ \vdots & \ddots & \vdots & & \vdots & \ddots & \vdots \\ a_{n,1} & \cdots & a_{n,j-1} & & a_{n,j+1} & \cdots & a_{n,n} \end{pmatrix}$$

VI.3. Determinants

Definition

Let $A \in M(n \times n)$. The symbol A_{ij} denotes the $(n - 1)$ -by- $(n - 1)$ matrix which is created from A by omitting the i th row and the j th column.

$$A_{ij} = \begin{pmatrix} a_{1,1} & \cdots & a_{1,j-1} & a_{1,j+1} & \cdots & a_{1,n} \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ a_{i-1,1} & \cdots & a_{i-1,j-1} & a_{i-1,j+1} & \cdots & a_{i-1,n} \\ a_{i+1,1} & \cdots & a_{i+1,j-1} & a_{i+1,j+1} & \cdots & a_{i+1,n} \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ a_{n,1} & \cdots & a_{n,j-1} & a_{n,j+1} & \cdots & a_{n,n} \end{pmatrix}$$

Definition

Let $\mathbf{A} = (a_{ij})_{i,j=1..n}$. The **determinant** of the matrix \mathbf{A} is defined by

$$\det \mathbf{A} = \begin{cases} a_{11} & \text{if } n = 1, \\ \sum_{i=1}^n (-1)^{i+1} a_{i1} \det \mathbf{A}_{i1} & \text{if } n > 1. \end{cases}$$

For $\det \mathbf{A}$ we will also use the symbol

$$\begin{vmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \ddots & \vdots & \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{vmatrix}.$$

Theorem 5 (cofactor expansion)

Let $\mathbf{A} = (a_{ij})_{i,j=1..n}$, $k \in \{1, \dots, n\}$. Then

$$\det \mathbf{A} = \sum_{i=1}^n (-1)^{i+k} a_{ik} \det \mathbf{A}_{ik} \quad (\text{expansion along } k\text{th column}),$$

$$\det \mathbf{A} = \sum_{j=1}^n (-1)^{k+j} a_{kj} \det \mathbf{A}_{kj} \quad (\text{expansion along } k\text{th row}).$$

Exercise

Find

$$\begin{vmatrix} 1 & 2 & 0 & -1 \\ -2 & 3 & 1 & 0 \\ -4 & 2 & 0 & 2 \\ 1 & 2 & 3 & -3 \end{vmatrix}$$

Exercise

Find

$$\begin{vmatrix} 1 & 2 & 0 & -1 \\ -2 & 3 & 1 & 0 \\ -4 & 2 & 0 & 2 \\ 1 & 2 & 3 & -3 \end{vmatrix}$$

2

Lemma 6

Let $j, n \in \mathbb{N}$, $j \leq n$, and the matrices $\mathbf{A}, \mathbf{B}, \mathbf{C} \in M(n \times n)$ coincide at each row except for the j th row. Let the j th row of \mathbf{A} be equal to the sum of the j th rows of \mathbf{B} and \mathbf{C} . Then $\det \mathbf{A} = \det \mathbf{B} + \det \mathbf{C}$.

$$\begin{vmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & \ddots & \vdots \\ a_{j-1,1} & \cdots & a_{j-1,n} \\ u_1+v_1 & \cdots & u_n+v_n \\ a_{j+1,1} & \cdots & a_{j+1,n} \\ \vdots & \ddots & \vdots \\ a_{n1} & \cdots & a_{nn} \end{vmatrix} = \begin{vmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & \ddots & \vdots \\ a_{j-1,1} & \cdots & a_{j-1,n} \\ u_1 & \cdots & u_n \\ a_{j+1,1} & \cdots & a_{j+1,n} \\ \vdots & \ddots & \vdots \\ a_{n1} & \cdots & a_{nn} \end{vmatrix} + \begin{vmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & \ddots & \vdots \\ a_{j-1,1} & \cdots & a_{j-1,n} \\ v_1 & \cdots & v_n \\ a_{j+1,1} & \cdots & a_{j+1,n} \\ \vdots & \ddots & \vdots \\ a_{n1} & \cdots & a_{nn} \end{vmatrix}$$

Exercise

$$\begin{vmatrix} 1 & 0 & -1 \\ 2 & 1 & 3 \\ -2 & 4 & 0 \end{vmatrix} = \begin{vmatrix} 1 & 0 & -1 \\ 1 & -1 & 1 \\ -2 & 4 & 0 \end{vmatrix} + \begin{vmatrix} 1 & 0 & -1 \\ 1 & 2 & 2 \\ -2 & 4 & 0 \end{vmatrix}$$

Theorem 7 (determinant and transformations)

Let $\mathbf{A}, \mathbf{A}' \in M(n \times n)$.

- (i) *If the matrix \mathbf{A}' is created from the matrix \mathbf{A} by multiplying one row in \mathbf{A} by a real number μ , then $\det \mathbf{A}' = \mu \det \mathbf{A}$.*

Theorem 7 (determinant and transformations)

Let $\mathbf{A}, \mathbf{A}' \in M(n \times n)$.

- (i) *If the matrix \mathbf{A}' is created from the matrix \mathbf{A} by multiplying one row in \mathbf{A} by a real number μ , then $\det \mathbf{A}' = \mu \det \mathbf{A}$.*
- (ii) *If the matrix \mathbf{A}' is created from \mathbf{A} by interchanging two rows in \mathbf{A} (i.e. by applying the elementary row operation of the first type), then $\det \mathbf{A}' = -\det \mathbf{A}$.*

Theorem 7 (determinant and transformations)

Let $\mathbf{A}, \mathbf{A}' \in M(n \times n)$.

- (i) *If the matrix \mathbf{A}' is created from the matrix \mathbf{A} by multiplying one row in \mathbf{A} by a real number μ , then $\det \mathbf{A}' = \mu \det \mathbf{A}$.*
- (ii) *If the matrix \mathbf{A}' is created from \mathbf{A} by interchanging two rows in \mathbf{A} (i.e. by applying the elementary row operation of the first type), then $\det \mathbf{A}' = -\det \mathbf{A}$.*
- (iii) *If the matrix \mathbf{A}' is created from \mathbf{A} by adding a μ -multiple of a row in \mathbf{A} to another row in \mathbf{A} (i.e. by applying the elementary row operation of the third type), then $\det \mathbf{A}' = \det \mathbf{A}$.*

Theorem 7 (determinant and transformations)

Let $\mathbf{A}, \mathbf{A}' \in M(n \times n)$.

- (i) *If the matrix \mathbf{A}' is created from the matrix \mathbf{A} by multiplying one row in \mathbf{A} by a real number μ , then $\det \mathbf{A}' = \mu \det \mathbf{A}$.*
- (ii) *If the matrix \mathbf{A}' is created from \mathbf{A} by interchanging two rows in \mathbf{A} (i.e. by applying the elementary row operation of the first type), then $\det \mathbf{A}' = -\det \mathbf{A}$.*
- (iii) *If the matrix \mathbf{A}' is created from \mathbf{A} by adding a μ -multiple of a row in \mathbf{A} to another row in \mathbf{A} (i.e. by applying the elementary row operation of the third type), then $\det \mathbf{A}' = \det \mathbf{A}$.*
- (iv) *If \mathbf{A}' is created from \mathbf{A} by applying a transformation, then $\det \mathbf{A} \neq 0$ if and only if $\det \mathbf{A}' \neq 0$.*

Exercise

Write a summary of the previous theorem.

Exercise

Write a summary of the previous theorem.

Remark

The determinant of a matrix with a zero row is equal to zero.
The determinant of a matrix with two identical rows is also equal to zero.

Exercise

We have

$$\det \begin{pmatrix} -1 & 15 & 16 \\ 2 & 5 & 4 \\ 2 & 3 & 5 \end{pmatrix} = -107.$$

Find

$$\det \begin{pmatrix} 2 & 5 & 4 \\ 2 & 3 & 5 \\ -1 & 15 & 16 \end{pmatrix} ?$$

A -107

B 107

C something else

Exercise

We have

$$\det \begin{pmatrix} -1 & 15 & 16 \\ 2 & 5 & 4 \\ 2 & 3 & 5 \end{pmatrix} = -107.$$

Find

$$\det \begin{pmatrix} 2 & 5 & 4 \\ 2 & 3 & 5 \\ -1 & 15 & 16 \end{pmatrix} ?$$

A -107

B 107

C something else

A

Exercise

We have

$$\det \begin{pmatrix} -2 & 1 & 3 \\ 2 & 0 & 4 \\ 1 & 3 & 1 \end{pmatrix} = 44.$$

Find

$$\det \begin{pmatrix} -2 & 1 & 3 \\ 0 & 1 & 7 \\ 1 & 3 & 1 \end{pmatrix} ?$$

A 44

B -44

C 88

D something else

Exercise

We have

$$\det \begin{pmatrix} -2 & 1 & 3 \\ 2 & 0 & 4 \\ 1 & 3 & 1 \end{pmatrix} = 44.$$

Find

$$\det \begin{pmatrix} -2 & 1 & 3 \\ 0 & 1 & 7 \\ 1 & 3 & 1 \end{pmatrix} ?$$

A 44

B -44

C 88

D something else

A

Exercise

We have

$$\det \begin{pmatrix} -2 & 1 & 3 \\ 2 & 0 & 4 \\ 1 & 3 & 1 \end{pmatrix} = 44.$$

Find

$$\det \begin{pmatrix} -2 & 1 & 3 \\ 2 & 0 & 4 \\ 0 & 7 & 5 \end{pmatrix} ?$$

A 44

B -44

C 88

D 22

E something else

Exercise

We have

$$\det \begin{pmatrix} -2 & 1 & 3 \\ 2 & 0 & 4 \\ 1 & 3 & 1 \end{pmatrix} = 44.$$

Find

$$\det \begin{pmatrix} -2 & 1 & 3 \\ 2 & 0 & 4 \\ 0 & 7 & 5 \end{pmatrix} ?$$

A 44

B -44

C 88

D 22

E something else

C

Exercise

Let \mathbf{A} be a matrix of type (2×2) . Find $\det(5\mathbf{A})$.

A $5 \det \mathbf{A}$

B $10 \det \mathbf{A}$

C $25 \det \mathbf{A}$

D something else

Exercise

Let \mathbf{A} be a matrix of type (2×2) . Find $\det(5\mathbf{A})$.

A $5 \det \mathbf{A}$

B $10 \det \mathbf{A}$

C $25 \det \mathbf{A}$

D something else

C

Definition

Let $A = (a_{ij})_{i,j=1..n}$. We say that A is an **upper triangular matrix** if $a_{ij} = 0$ for $i > j$, $i, j \in \{1, \dots, n\}$.

Definition

Let $A = (a_{ij})_{i,j=1..n}$. We say that A is an **upper triangular matrix** if $a_{ij} = 0$ for $i > j$, $i, j \in \{1, \dots, n\}$. We say that A is a **lower triangular matrix** if $a_{ij} = 0$ for $i < j$, $i, j \in \{1, \dots, n\}$.

Example



$$\begin{pmatrix} 1 & 2 & 4 \\ 0 & -3 & 0 \\ 0 & 0 & 5 \end{pmatrix}$$



$$\begin{pmatrix} 1 & 0 & 0 \\ 2 & -4 & 0 \\ 3 & 3 & 3 \end{pmatrix}$$

Definition

Let $A = (a_{ij})_{i,j=1..n}$. We say that A is an **upper triangular matrix** if $a_{ij} = 0$ for $i > j$, $i, j \in \{1, \dots, n\}$. We say that A is a **lower triangular matrix** if $a_{ij} = 0$ for $i < j$, $i, j \in \{1, \dots, n\}$.

Example

$$\begin{pmatrix} 1 & 2 & 4 \\ 0 & -3 & 0 \\ 0 & 0 & 5 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 0 & 0 \\ 2 & -4 & 0 \\ 3 & 3 & 3 \end{pmatrix}$$

Theorem 8 (determinant of a triangular matrix)

Let $A = (a_{ij})_{i,j=1..n}$ be an upper or lower triangular matrix.
Then

$$\det A = a_{11} \cdot a_{22} \cdot \dots \cdot a_{nn}.$$

Theorem 9 (determinant and invertibility)

Let $\mathbf{A} \in M(n \times n)$. Then \mathbf{A} is invertible if and only if $\det \mathbf{A} \neq 0$.

Exercise

Which of the following matrices do NOT have inverse matrix?

A

$$\begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$$

B

$$\begin{pmatrix} 2 & 2 \\ 4 & 4 \end{pmatrix}$$

C

$$\begin{pmatrix} -1 & 0 \\ 0 & 3 \end{pmatrix}$$

D

$$\begin{pmatrix} 0 & 4 \\ 2 & 0 \end{pmatrix}$$

E All of them have inverse matrix.

Exercise

Which of the following matrices do NOT have inverse matrix?

A

$$\begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$$

B

$$\begin{pmatrix} 2 & 2 \\ 4 & 4 \end{pmatrix}$$

C

$$\begin{pmatrix} -1 & 0 \\ 0 & 3 \end{pmatrix}$$

D

$$\begin{pmatrix} 0 & 4 \\ 2 & 0 \end{pmatrix}$$

E All of them have inverse matrix.

B

Theorem 10 (determinant of a product)

Let $\mathbf{A}, \mathbf{B} \in M(n \times n)$. Then $\det \mathbf{AB} = \det \mathbf{A} \cdot \det \mathbf{B}$.

Remark

Let \mathbf{A} be invertible. Then $\det \mathbf{A}^{-1} = \frac{1}{\det \mathbf{A}}$

Exercise

Let $\det \mathbf{A} = 3$. Find $\det \mathbf{A}^{-1}$.

A 3

B $1/3$

C 9

D hard to say.

Exercise

Let $\det \mathbf{A} = 3$. Find $\det \mathbf{A}^{-1}$.

A 3

B $1/3$

C 9

D hard to say.

B

Theorem 11 (determinant of a transpose)

Let $\mathbf{A} \in M(n \times n)$. Then $\det \mathbf{A}^T = \det \mathbf{A}$.

Exercise

We have

$$\det \begin{pmatrix} -2 & 1 & 3 \\ 2 & 0 & 4 \\ 1 & 3 & 1 \end{pmatrix} = 44.$$

Find

$$\det \begin{pmatrix} -2 & 2 & 1 \\ 1 & 0 & 3 \\ 3 & 4 & 1 \end{pmatrix} ?$$

A 44

B 1/44

C 88

D 22

E -44

Exercise

We have

$$\det \begin{pmatrix} -2 & 1 & 3 \\ 2 & 0 & 4 \\ 1 & 3 & 1 \end{pmatrix} = 44.$$

Find

$$\det \begin{pmatrix} -2 & 2 & 1 \\ 1 & 0 & 3 \\ 3 & 4 & 1 \end{pmatrix} ?$$

A 44

B 1/44

C 88

D 22

E -44

A

Definition

Let $k, n \in \mathbb{N}$ and $\mathbf{v}^1, \dots, \mathbf{v}^k \in \mathbb{R}^n$. We say that a vector $\mathbf{u} \in \mathbb{R}^n$ is a **linear combination of the vectors $\mathbf{v}^1, \dots, \mathbf{v}^k$ with coefficients $\lambda_1, \dots, \lambda_k \in \mathbb{R}$** if

$$\mathbf{u} = \lambda_1 \mathbf{v}^1 + \dots + \lambda_k \mathbf{v}^k.$$

Definition

Let $k, n \in \mathbb{N}$ and $\mathbf{v}^1, \dots, \mathbf{v}^k \in \mathbb{R}^n$. We say that a vector $\mathbf{u} \in \mathbb{R}^n$ is a **linear combination of the vectors $\mathbf{v}^1, \dots, \mathbf{v}^k$ with coefficients $\lambda_1, \dots, \lambda_k \in \mathbb{R}$** if

$$\mathbf{u} = \lambda_1 \mathbf{v}^1 + \dots + \lambda_k \mathbf{v}^k.$$

By a **trivial linear combination** of vectors $\mathbf{v}^1, \dots, \mathbf{v}^k$ we mean the linear combination $0 \cdot \mathbf{v}^1 + \dots + 0 \cdot \mathbf{v}^k$. Linear combination which is not trivial is called **non-trivial**.

<https://www.geogebra.org/m/z8cZnvAm>

Exercise

Let $u = (1, 2, 4)$ and $v = (-2, 0, 5)$. Then $2u - 3v$ is

A $(-4, 4, 23)$

C $(8, 4, 23)$

B $(8, 4, -7)$

D $(7, 6, 2)$

Exercise

Let $u = (1, 2, 4)$ and $v = (-2, 0, 5)$. Then $2u - 3v$ is

A $(-4, 4, 23)$

C $(8, 4, 23)$

B $(8, 4, -7)$

D $(7, 6, 2)$

B

Exercise

Express $z = (-5, 3, 6)$ as the linear combination of $x = (1, -1, 4)$ and $y = (-3, 2, 6)$.

- A $-5x$
- B $-2x + y$
- C $x + 2y$
- D $2x + y$
- E hard to say

Exercise

Express $z = (-5, 3, 6)$ as the linear combination of $x = (1, -1, 4)$ and $y = (-3, 2, 6)$.

A $-5x$

B $-2x + y$

C $x + 2y$

D $2x + y$

E hard to say

E, impossible

Exercise

Express w as the linear combination of u and v .




Figure: <https://www.chegg.com/homework-help/questions-and-answers/write-vector-w-linear-combination-u-v-q55559120>

A $w = 2u + v$

B $w = u + v$

C $w = -u + v$

D $w = u - v$

E w cannot be written like that.

Exercise

Express w as the linear combination of u and v .




Figure: <https://www.chegg.com/homework-help/questions-and-answers/write-vector-w-linear-combination-u-v-q55559120>

A $w = 2u + v$

B $w = u + v$

C $w = -u + v$

D $w = u - v$

E w cannot be written like that.

A

Exercise

Which of the following vector can be written as the linear combination of vectors $(1, 0, 0)$, $(0, 1, 0)$, $(0, 0, 1)$?

A $(0, 2, 0)$

B $(-3, 0, 1)$

C $(0.4, 3.7, -1.5)$

Exercise

Which of the following vector can be written as the linear combination of vectors $(1, 0, 0)$, $(0, 1, 0)$, $(0, 0, 1)$?

A $(0, 2, 0)$

B $(-3, 0, 1)$

C $(0.4, 3.7, -1.5)$

A, B, C

Exercise

Which of the following vector can be written as the linear combination of vectors $(1, 0, 0)$, $(0, 1, 0)$, $(0, 0, 1)$?

A $(0, 2, 0)$

B $(-3, 0, 1)$

C $(0.4, 3.7, -1.5)$

A, B, C

Exercise

Describe the set of all linear combinations of vectors $(2, 4, 6)$ and $(-1, -2, -3)$?

A point

B line

C vector

D plane

E space

Exercise

Which of the following vector can be written as the linear combination of vectors $(1, 0, 0)$, $(0, 1, 0)$, $(0, 0, 1)$?

A $(0, 2, 0)$

B $(-3, 0, 1)$

C $(0.4, 3.7, -1.5)$

A, B, C

Exercise

Describe the set of all linear combinations of vectors $(2, 4, 6)$ and $(-1, -2, -3)$?

A point

B line

C vector

D plane

E space

B

Exercise

Describe the set of all linear combinations of vectors $(1, 2, 0)$ and $(-1, 1, 0)$?

A point B line C vector D plane E space

Exercise

Describe the set of all linear combinations of vectors $(1, 2, 0)$ and $(-1, 1, 0)$?

A point B line C vector D plane E space

D

Exercise

Describe the set of all linear combinations of vectors $(1, 2, 0)$, $(-1, 1, 0)$ and $(2, -2, 0)$?

A point B line C vector D plane E space

Exercise

Describe the set of all linear combinations of vectors $(1, 2, 0)$ and $(-1, 1, 0)$?

A point B line C vector D plane E space

D

Exercise

Describe the set of all linear combinations of vectors $(1, 2, 0)$, $(-1, 1, 0)$ and $(2, -2, 0)$?

A point B line C vector D plane E space

D

Definition

We say that vectors $\mathbf{v}^1, \dots, \mathbf{v}^k \in \mathbb{R}^n$ are **linearly dependent** if there exists their non-trivial linear combination which is equal to the zero vector.

Definition

We say that vectors $\mathbf{v}^1, \dots, \mathbf{v}^k \in \mathbb{R}^n$ are **linearly dependent** if there exists their non-trivial linear combination which is equal to the zero vector. We say that vectors $\mathbf{v}^1, \dots, \mathbf{v}^k \in \mathbb{R}^n$ are **linearly independent** if they are not linearly dependent, i.e. if whenever $\lambda_1, \dots, \lambda_k \in \mathbb{R}$ satisfy $\lambda_1 \mathbf{v}^1 + \dots + \lambda_k \mathbf{v}^k = \mathbf{o}$, then $\lambda_1 = \lambda_2 = \dots = \lambda_k = 0$.

Definition

We say that vectors $\mathbf{v}^1, \dots, \mathbf{v}^k \in \mathbb{R}^n$ are **linearly dependent** if there exists their non-trivial linear combination which is equal to the zero vector. We say that vectors $\mathbf{v}^1, \dots, \mathbf{v}^k \in \mathbb{R}^n$ are **linearly independent** if they are not linearly dependent, i.e. if whenever $\lambda_1, \dots, \lambda_k \in \mathbb{R}$ satisfy $\lambda_1 \mathbf{v}^1 + \dots + \lambda_k \mathbf{v}^k = \mathbf{o}$, then $\lambda_1 = \lambda_2 = \dots = \lambda_k = 0$.

Remark

Vectors $\mathbf{v}^1, \dots, \mathbf{v}^k$ are linearly dependent if and only if one of them can be expressed as a linear combination of the others.

Remark

Vectors $\mathbf{v}^1, \dots, \mathbf{v}^k$ are linearly dependent if and only if one of them can be expressed as a linear combination of the others.

Exercise

The vectors $(1, 0, 0)$, $(0, 0, 2)$, $(3, 0, 4)$ are

- A linearly dependent
- B linearly independent

Remark

Vectors $\mathbf{v}^1, \dots, \mathbf{v}^k$ are linearly dependent if and only if one of them can be expressed as a linear combination of the others.

Exercise

The vectors $(1, 0, 0)$, $(0, 0, 2)$, $(3, 0, 4)$ are

A linearly dependent

B linearly independent

A

Definition

Let $A \in M(m \times n)$. The **rank** of the matrix A is the maximal number of linearly independent row vectors of A , i.e. the rank is equal to $k \in \mathbb{N}$ if

- (i) there is k linearly independent row vectors of A and
- (ii) each l -tuple of row vectors of A , where $l > k$, is linearly dependent.

The rank of the zero matrix is zero. Rank of A is denoted by $\text{rank}(A)$.

Exercise

Find the rank of the matrix

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \\ 1 & 2 & 3 \end{pmatrix}$$

Exercise

Find the rank of the matrix

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \\ 1 & 2 & 3 \end{pmatrix}$$

3

Definition

We say that a matrix $A \in M(m \times n)$ is in a **row echelon form** if for each $i \in \{2, \dots, m\}$ the i th row of A is either a zero vector or it has more zeros at the beginning than the $(i - 1)$ th row.

Example

$$\begin{pmatrix} 1 & 4 & 1 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{pmatrix} \quad \begin{pmatrix} 1 & 4 & 1 \\ 0 & 0 & 2 \\ 0 & 0 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 4 & 1 \\ 0 & 5 & 2 \end{pmatrix}$$

Example (Not echelon form)

$$\begin{pmatrix} 2 & -4 & 1 \\ 0 & 0 & 3 \\ 0 & 2 & 0 \end{pmatrix} \quad \begin{pmatrix} -1 & 4 & 1 \\ 3 & 2 & 2 \\ 0 & 0 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 4 & 1 \\ 2 & 0 & 0 \end{pmatrix}$$

Remark

The rank of a row echelon matrix is equal to the number of its non-zero rows.

Example

$$\begin{pmatrix} 1 & 4 & 1 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{pmatrix} \begin{pmatrix} 1 & 4 & 1 \\ 0 & 0 & 2 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} 1 & 4 & 1 \\ 0 & 5 & 2 \end{pmatrix}$$

Definition

The **elementary row operations** on the matrix A are:

- (i) interchange of two rows,
- (ii) multiplication of a row by a non-zero real number,
- (iii) addition of a multiple of a row to another row.

Definition

A matrix **transformation** is a finite sequence of elementary row operations. If a matrix $B \in M(m \times n)$ results from the matrix $A \in M(m \times n)$ by applying a transformation T on the matrix A , then this fact is denoted by $A \xrightarrow{T} B$.

Theorem 12 (properties of matrix transformations)

(i) *Let $A \in M(m \times n)$. Then there exists a transformation transforming A to a row echelon matrix.*

Theorem 12 (properties of matrix transformations)

- (i) *Let $\mathbf{A} \in M(m \times n)$. Then there exists a transformation transforming \mathbf{A} to a row echelon matrix.*
- (ii) *Let T_1 be a transformation applicable to m -by- n matrices. Then there exists a transformation T_2 applicable to m -by- n matrices such that for any two matrices $\mathbf{A}, \mathbf{B} \in M(m \times n)$ we have $\mathbf{A} \xrightarrow{T_1} \mathbf{B}$ if and only if $\mathbf{B} \xrightarrow{T_2} \mathbf{A}$.*

Theorem 12 (properties of matrix transformations)

- (i) Let $\mathbf{A} \in M(m \times n)$. Then there exists a transformation transforming \mathbf{A} to a row echelon matrix.
- (ii) Let T_1 be a transformation applicable to m -by- n matrices. Then there exists a transformation T_2 applicable to m -by- n matrices such that for any two matrices $\mathbf{A}, \mathbf{B} \in M(m \times n)$ we have $\mathbf{A} \xrightarrow{T_1} \mathbf{B}$ if and only if $\mathbf{B} \xrightarrow{T_2} \mathbf{A}$.
- (iii) Let $\mathbf{A}, \mathbf{B} \in M(m \times n)$ and there exist a transformation T such that $\mathbf{A} \xrightarrow{T} \mathbf{B}$. Then $\text{rank}(\mathbf{A}) = \text{rank}(\mathbf{B})$.

Exercise

Let

$$\mathbf{A} = \begin{pmatrix} 5 & 4 & -8 & 1 \\ 1 & 3 & 4 & 8 \\ 0 & 2 & 1 & 3 \\ -1 & -2 & 4 & 1 \end{pmatrix}.$$

After the transformation we get

$$\begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

Find the rank of \mathbf{A} :

A 0

B 1

C 2

D 3

E 4

Exercise

Let

$$\mathbf{A} = \begin{pmatrix} 5 & 4 & -8 & 1 \\ 1 & 3 & 4 & 8 \\ 0 & 2 & 1 & 3 \\ -1 & -2 & 4 & 1 \end{pmatrix}.$$

After the transformation we get

$$\begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

Find the rank of \mathbf{A} :

A 0

B 1

C 2

D 3

E 4

D

Remark

Similarly as the elementary row operations one can define also elementary column operations. It can be shown that the elementary column operations do not change the rank of the matrix.

Remark


Similarly as the elementary row operations one can define also elementary column operations. It can be shown that the elementary column operations do not change the rank of the matrix.

Remark

It can be shown that $\text{rank}(\mathbf{A}) = \text{rank}(\mathbf{A}^T)$ for any $\mathbf{A} \in M(m \times n)$.

Exercise

We made a matrix from the vectors x , y , u , v and w . Find rank of this matrix.



<http://mathquest.carroll.edu/libraries/FHMW.student.edition.pdf>

A 1

B 2


C 3

D 4

E 5

Exercise

We made a matrix from the vectors x , y , u , v and w . Find rank of this matrix.



<http://mathquest.carroll.edu/libraries/FHMW.student.edition.pdf>

A 1

B 2

C 3

D 4

E 5

B

Theorem 13 (representation of a transformation)

Let T be a transformation on $m \times n$ matrices. Then there exists an invertible matrix $\mathbf{C}_T \in M(m \times m)$ satisfying:
whenever we apply the transformation T to a matrix $\mathbf{A} \in M(m \times n)$, we obtain the matrix $\mathbf{C}_T \mathbf{A}$.

Example

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 2 & 1 \end{pmatrix} \cdot \begin{pmatrix} 1 & 2 & 3 \\ -1 & -2 & -3 \\ 2 & 3 & 4 \end{pmatrix} = \begin{pmatrix} 1 & 2 & 3 \\ -1 & -2 & -3 \\ 0 & -1 & -2 \end{pmatrix}$$

Theorem 13 (representation of a transformation)

Let T be a transformation on $m \times n$ matrices. Then there exists an invertible matrix $C_T \in M(m \times m)$ satisfying: whenever we apply the transformation T to a matrix $A \in M(m \times n)$, we obtain the matrix $C_T A$.

Example

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 2 & 1 \end{pmatrix} \cdot \begin{pmatrix} 1 & 2 & 3 \\ -1 & -2 & -3 \\ 2 & 3 & 4 \end{pmatrix} = \begin{pmatrix} 1 & 2 & 3 \\ -1 & -2 & -3 \\ 0 & -1 & -2 \end{pmatrix}$$

Remark

Also the converse is true: For every invertible matrix C the mapping $A \mapsto CA$ is a transformation.

Lemma 14

Let $\mathbf{A} \in M(n \times n)$ and $\text{rank}(\mathbf{A}) = n$. Then there exists a transformation transforming \mathbf{A} to \mathbf{I} .

Lemma 14

Let $\mathbf{A} \in M(n \times n)$ and $\text{rank}(\mathbf{A}) = n$. Then there exists a transformation transforming \mathbf{A} to \mathbf{I} .

Theorem 15

Let $\mathbf{A} \in M(n \times n)$. Then \mathbf{A} is invertible if and only if $\text{rank}(\mathbf{A}) = n$.

Exercise

Find invertible matrices:

$$\mathbf{A} = \begin{pmatrix} 3 & 0 \\ 2 & 1 \end{pmatrix}$$

$$\mathbf{B} = \begin{pmatrix} 1 & 2 \\ -2 & -4 \end{pmatrix}$$

$$\mathbf{C} = \begin{pmatrix} 1 & 0 & 4 \\ 0 & 2 & 5 \\ 0 & 0 & -1 \end{pmatrix}$$

$$\mathbf{D} = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 4 & 1 \\ 3 & 6 & 4 \end{pmatrix}$$

Lemma 14

Let $\mathbf{A} \in M(n \times n)$ and $\text{rank}(\mathbf{A}) = n$. Then there exists a transformation transforming \mathbf{A} to \mathbf{I} .

Theorem 15

Let $\mathbf{A} \in M(n \times n)$. Then \mathbf{A} is invertible if and only if $\text{rank}(\mathbf{A}) = n$.

Exercise

Find invertible matrices:

$$\mathbf{A} = \begin{pmatrix} 3 & 0 \\ 2 & 1 \end{pmatrix}$$

$$\mathbf{B} = \begin{pmatrix} 1 & 2 \\ -2 & -4 \end{pmatrix}$$

$$\mathbf{C} = \begin{pmatrix} 1 & 0 & 4 \\ 0 & 2 & 5 \\ 0 & 0 & -1 \end{pmatrix}$$

$$\mathbf{D} = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 4 & 1 \\ 3 & 6 & 4 \end{pmatrix}$$

A, C

VI.4. Systems of linear equations

A system of m equations in n unknowns x_1, \dots, x_n :

$$a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = b_1,$$

$$a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = b_2,$$

$$\vdots$$

$$a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n = b_m,$$

(S)

where $a_{ij} \in \mathbb{R}$, $b_i \in \mathbb{R}$, $i = 1, \dots, m$, $j = 1, \dots, n$.

A system of m equations in n unknowns x_1, \dots, x_n :

$$\begin{aligned} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n &= b_1, \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n &= b_2, \\ &\vdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n &= b_m, \end{aligned} \tag{S}$$

where $a_{ij} \in \mathbb{R}$, $b_i \in \mathbb{R}$, $i = 1, \dots, m$, $j = 1, \dots, n$. The matrix form is

$$\mathbf{Ax} = \mathbf{b},$$

where $\mathbf{A} = \begin{pmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & \ddots & \vdots \\ a_{m1} & \cdots & a_{mn} \end{pmatrix} \in M(m \times n)$, is called the **coefficient matrix**, $\mathbf{b} = \begin{pmatrix} b_1 \\ \vdots \\ b_m \end{pmatrix} \in M(m \times 1)$ is called the **vector of the right-hand side** and $\mathbf{x} = \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix} \in M(n \times 1)$ is the **vector of unknowns**.

Definition

The matrix

$$(A|\mathbf{b}) = \left(\begin{array}{ccc|c} a_{11} & \dots & a_{1n} & b_1 \\ \vdots & \ddots & \vdots & \vdots \\ a_{m1} & \dots & a_{mn} & b_m \end{array} \right)$$

is called the **augmented matrix of the system (S)**.

$$6x + 2y = 100$$

$$4x + y = 60$$

$$\left(\begin{array}{cc|c} 6 & 2 & 100 \\ 4 & 1 & 60 \end{array} \right)$$

Proposition 16 (solutions of a transformed system)

Let $\mathbf{A} \in M(m \times n)$, $\mathbf{b} \in M(m \times 1)$ and let T be a transformation of matrices with m rows. Denote $\mathbf{A} \xrightarrow{T} \mathbf{A}'$, $\mathbf{b} \xrightarrow{T} \mathbf{b}'$. Then for any $\mathbf{y} \in M(n \times 1)$ we have $\mathbf{A}\mathbf{y} = \mathbf{b}$ if and only if $\mathbf{A}'\mathbf{y} = \mathbf{b}'$, i.e. the systems $\mathbf{A}\mathbf{x} = \mathbf{b}$ and $\mathbf{A}'\mathbf{x} = \mathbf{b}'$ have the same set of solutions.

Theorem 17 (Rouché-Fontené)

The system (S) has a solution if and only if its coefficient matrix has the same rank as its augmented matrix.

$$\left(\begin{array}{ccc|c} 1 & 2 & 3 & 4 \\ 0 & 1 & -2 & 2 \\ 0 & 0 & 4 & 1 \end{array} \right) \quad \left(\begin{array}{ccc|c} 1 & 2 & 3 & 4 \\ 0 & 1 & -2 & 2 \\ 0 & 0 & 0 & 0 \end{array} \right) \quad \left(\begin{array}{ccc|c} 1 & 2 & 3 & 4 \\ 0 & 1 & -2 & 2 \\ 0 & 0 & 0 & 1 \end{array} \right)$$

Theorem 18 (Cramer's rule)

Let $\mathbf{A} \in M(n \times n)$ be an invertible matrix, $\mathbf{b} \in M(n \times 1)$, $\mathbf{x} \in M(n \times 1)$, and $\mathbf{Ax} = \mathbf{b}$. Then

$$x_j = \frac{\begin{vmatrix} a_{11} & \dots & a_{1,j-1} & b_1 & a_{1,j+1} & \dots & a_{1n} \\ \vdots & & & \vdots & & & \vdots \\ a_{n1} & \dots & a_{n,j-1} & b_n & a_{n,j+1} & \dots & a_{nn} \end{vmatrix}}{\det \mathbf{A}}$$

for $j = 1, \dots, n$.

Exercise

$$\left(\begin{array}{ccc|c} 1 & 1 & -1 & 6 \\ 3 & -2 & 1 & -5 \\ 1 & 3 & -2 & 14 \end{array} \right)$$

Theorem 18 (Cramer's rule)

Let $\mathbf{A} \in M(n \times n)$ be an invertible matrix, $\mathbf{b} \in M(n \times 1)$, $\mathbf{x} \in M(n \times 1)$, and $\mathbf{Ax} = \mathbf{b}$. Then

$$x_j = \frac{\begin{vmatrix} a_{11} & \dots & a_{1,j-1} & b_1 & a_{1,j+1} & \dots & a_{1n} \\ \vdots & & \vdots & \vdots & \vdots & & \vdots \\ a_{n1} & \dots & a_{n,j-1} & b_n & a_{n,j+1} & \dots & a_{nn} \end{vmatrix}}{\det \mathbf{A}}$$

for $j = 1, \dots, n$.

Exercise

$$\left(\begin{array}{ccc|c} 1 & 1 & -1 & 6 \\ 3 & -2 & 1 & -5 \\ 1 & 3 & -2 & 14 \end{array} \right)$$

$$x = 1, y = 3, z = -2$$

VI.5. Definiteness of matrices

Definition

We say that a **symmetric** matrix $A \in M(n \times n)$ is

- **positive definite** (PD), if $\mathbf{u}^T A \mathbf{u} > 0$ for all $\mathbf{u} \in \mathbb{R}^n$, $\mathbf{u} \neq \mathbf{o}$,
- **negative definite** (ND), if $\mathbf{u}^T A \mathbf{u} < 0$ for all $\mathbf{u} \in \mathbb{R}^n$, $\mathbf{u} \neq \mathbf{o}$,
- **positive semidefinite** (PSD), if $\mathbf{u}^T A \mathbf{u} \geq 0$ for all $\mathbf{u} \in \mathbb{R}^n$,
- **negative semidefinite** (NSD), if $\mathbf{u}^T A \mathbf{u} \leq 0$ for all $\mathbf{u} \in \mathbb{R}^n$,
- **indefinite** (ID), if there exist $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$ such that $\mathbf{u}^T A \mathbf{u} > 0$ and $\mathbf{v}^T A \mathbf{v} < 0$.

$$\begin{aligned} (x \ y) \cdot \begin{pmatrix} 7 & 1 \\ 1 & 2 \end{pmatrix} \cdot \begin{pmatrix} x \\ y \end{pmatrix} &= 7x^2 + 4xy + y^2 = 3x^2 + 4x^2 + 4xy + y^2 \\ &= 3x^2 + (y + 2x)^2 > 0 \end{aligned}$$

Proposition 19 (definiteness of diagonal matrices)

Let $A \in M(n \times n)$ be **diagonal** (i.e. $a_{ij} = 0$ whenever $i \neq j$).
Then

- A is PD if and only if $a_{ii} > 0$ for all $i = 1, 2, \dots, n$,
- A is ND if and only if $a_{ii} < 0$ for all $i = 1, 2, \dots, n$,
- A is PSD if and only if $a_{ii} \geq 0$ for all $i = 1, 2, \dots, n$,
- A is NSD if and only if $a_{ii} \leq 0$ for all $i = 1, 2, \dots, n$,
- A is ID if and only if there exist $i, j \in \{1, 2, \dots, n\}$ such that $a_{ii} > 0$ and $a_{jj} < 0$.

Exercise

Decide about definiteness of the following matrices:

$$\begin{pmatrix} -2 & 0 \\ 0 & -5 \end{pmatrix} \quad \begin{pmatrix} 2 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 8 \end{pmatrix} \quad \begin{pmatrix} 2 & 0 & 0 & 0 \\ 0 & -3 & 0 & 0 \\ 0 & 0 & 4 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

Proposition 20 (necessary conditions for definiteness)

Let $A \in M(n \times n)$ be a symmetric matrix. Then

- If A is PD, then $a_{ii} > 0$ for all $i = 1, 2, \dots, n$,
- If A is ND, then $a_{ii} < 0$ for all $i = 1, 2, \dots, n$,
- If A is PSD, then $a_{ii} \geq 0$ for all $i = 1, 2, \dots, n$,
- If A is NSD, then $a_{ii} \leq 0$ for all $i = 1, 2, \dots, n$,
- If there exist $i, j \in \{1, 2, \dots, n\}$ such that $a_{ii} > 0$ and $a_{jj} < 0$, then A is ID.

Exercise

Which of this matrices can NOT be negative semidefinite?

A

$$\begin{pmatrix} 5 & 1 & -4 \\ 1 & 9 & 2 \\ -4 & 2 & -5 \end{pmatrix}$$

B

$$\begin{pmatrix} -1 & 3 & 8 \\ 3 & -2 & 1 \\ 8 & 1 & -2 \end{pmatrix}$$

C

$$\begin{pmatrix} -1 & 2 & -11 \\ 2 & 0 & 6 \\ -11 & 6 & -5 \end{pmatrix}$$

Proposition 20 (necessary conditions for definiteness)

Let $A \in M(n \times n)$ be a symmetric matrix. Then

- If A is PD, then $a_{ii} > 0$ for all $i = 1, 2, \dots, n$,
- If A is ND, then $a_{ii} < 0$ for all $i = 1, 2, \dots, n$,
- If A is PSD, then $a_{ii} \geq 0$ for all $i = 1, 2, \dots, n$,
- If A is NSD, then $a_{ii} \leq 0$ for all $i = 1, 2, \dots, n$,
- If there exist $i, j \in \{1, 2, \dots, n\}$ such that $a_{ii} > 0$ and $a_{jj} < 0$, then A is ID.

Exercise

Which of this matrices can NOT be negative semidefinite?

A

$$\begin{pmatrix} 5 & 1 & -4 \\ 1 & 9 & 2 \\ -4 & 2 & -5 \end{pmatrix}$$

B

$$\begin{pmatrix} -1 & 3 & 8 \\ 3 & -2 & 1 \\ 8 & 1 & -2 \end{pmatrix}$$

C

$$\begin{pmatrix} -1 & 2 & -11 \\ 2 & 0 & 6 \\ -11 & 6 & -5 \end{pmatrix}$$

A

Theorem 21 (Sylvester's criterion)

Let $\mathbf{A} = (a_{ij}) \in M(n \times n)$ be a symmetric matrix. Then \mathbf{A} is

- positive definite if and only if

$$\begin{vmatrix} a_{11} & \dots & a_{1k} \\ \vdots & & \vdots \\ a_{k1} & \dots & a_{kk} \end{vmatrix} > 0 \quad \text{for all } k = 1, \dots, n,$$

- negative definite if and only if

$$(-1)^k \begin{vmatrix} a_{11} & \dots & a_{1k} \\ \vdots & & \vdots \\ a_{k1} & \dots & a_{kk} \end{vmatrix} > 0 \quad \text{for all } k = 1, \dots, n,$$