

Group cohomology and wallpaper groups III

Michal Hrbek

Fall school of Department of Algebra

November 23, 2014

Recapitulation

What do we have so far:

What do we have so far:

- Each wallpaper group G is an extension of \mathbb{Z}^2 by a point group G_0 .

Recapitulation

What do we have so far:

- Each wallpaper group G is an extension of \mathbb{Z}^2 by a point group G_0 . That is, G has a normal subgroup T isomorphic to \mathbb{Z}^2 , such that $G/T \simeq G_0$.

Recapitulation

What do we have so far:

- Each wallpaper group G is an extension of \mathbb{Z}^2 by a point group G_0 . That is, G has a normal subgroup T isomorphic to \mathbb{Z}^2 , such that $G/T \simeq G_0$.
- The point group G_0 is finite and uniquely determined by G . Moreover, G_0 is always one of the following groups:

$$C_1, C_2, C_3, C_4, C_6, D_1, D_2, D_3, D_4, D_6.$$

Recapitulation

What do we have so far:

- Each wallpaper group G is an extension of \mathbb{Z}^2 by a point group G_0 . That is, G has a normal subgroup T isomorphic to \mathbb{Z}^2 , such that $G/T \simeq G_0$.
- The point group G_0 is finite and uniquely determined by G . Moreover, G_0 is always one of the following groups:

$$C_1, C_2, C_3, C_4, C_6, D_1, D_2, D_3, D_4, D_6.$$

- With any extension of T by G_0 comes naturally an action φ of G_0 on T defined as follows: $\varphi_g(t) = x_g t x_g^{-1}$, where x_g is an element of G which is sent to g in the projection $G \rightarrow G_0$.

Recapitulation

Any extension

$$0 \rightarrow T \rightarrow G \rightarrow G_0 \rightarrow 1$$

is up to equivalence uniquely determined by the following data:

Recapitulation

Any extension

$$0 \rightarrow T \rightarrow G \rightarrow G_0 \rightarrow 1$$

is up to equivalence uniquely determined by the following data:

- ① Choice of the point group $G_0 \in \{C_i, D_i \mid i = 1, 2, 3, 4, 6\}$.

Recapitulation

Any extension

$$0 \rightarrow T \rightarrow G \rightarrow G_0 \rightarrow 1$$

is up to equivalence uniquely determined by the following data:

- ① Choice of the point group $G_0 \in \{C_i, D_i \mid i = 1, 2, 3, 4, 6\}$.
- ② Choice of the action φ of G_0 on T .

Recapitulation

Any extension

$$0 \rightarrow T \rightarrow G \rightarrow G_0 \rightarrow 1$$

is up to equivalence uniquely determined by the following data:

- ① Choice of the point group $G_0 \in \{C_i, D_i \mid i = 1, 2, 3, 4, 6\}$.
- ② Choice of the action φ of G_0 on T .
- ③ Choice of an element of the second cohomology group $H_\varphi^2(G_0, T)$.

Any extension

$$0 \rightarrow T \rightarrow G \rightarrow G_0 \rightarrow 1$$

is up to equivalence uniquely determined by the following data:

- ① Choice of the point group $G_0 \in \{C_i, D_i \mid i = 1, 2, 3, 4, 6\}$.
- ② Choice of the action φ of G_0 on T .
- ③ Choice of an element of the second cohomology group $H_\varphi^2(G_0, T)$.

E.g., the zero element of $H_\varphi^2(G_0, T)$ corresponds to the trivial extension of T by G_0 with action φ , that is, the semidirect product $T \rtimes_\varphi G_0$.

Recapitulation

- There are 13 distinct (non-conjugate) actions of point groups on \mathbb{Z}^2 .

Recapitulation

- There are 13 distinct (non-conjugate) actions of point groups on \mathbb{Z}^2 .
- That is, 13 inequivalent representations $G_0 \rightarrow GL(\mathbb{Z}^2)$, letting G_0 run over all possible point groups.

Recapitulation

- There are 13 distinct (non-conjugate) actions of point groups on \mathbb{Z}^2 .
- That is, 13 inequivalent representations $G_0 \rightarrow GL(\mathbb{Z}^2)$, letting G_0 run over all possible point groups.
- There is exactly one representation of each cyclic group C_i , and of dihedral groups D_4 and D_6 .

Recapitulation

- There are 13 distinct (non-conjugate) actions of point groups on \mathbb{Z}^2 .
- That is, 13 inequivalent representations $G_0 \rightarrow GL(\mathbb{Z}^2)$, letting G_0 run over all possible point groups.
- There is exactly one representation of each cyclic group C_i , and of dihedral groups D_4 and D_6 .
- There two representations of each of D_1, D_2, D_3 .

Recapitulation

- There are 13 distinct (non-conjugate) actions of point groups on \mathbb{Z}^2 .
- That is, 13 inequivalent representations $G_0 \rightarrow GL(\mathbb{Z}^2)$, letting G_0 run over all possible point groups.
- There is exactly one representation of each cyclic group C_i , and of dihedral groups D_4 and D_6 .
- There two representations of each of D_1, D_2, D_3 .
- Let us distinguish the representations as $D_{1,p}, D_{1,c}, D_{2,p}, D_{2,c}, D_{3,l}, D_{3,s}$.

Recapitulation

Together we have 18 equivalence classes of extensions:

G_0	$H^2(G_0, T)$	No. of extensions
C_1	0	1
C_2	0	1
C_3	0	1
C_4	0	1
C_6	0	1
$D_{1,p}$	\mathbb{Z}_2	2
$D_{1,c}$	0	1
$D_{2,p}$	$\mathbb{Z}_2 \times \mathbb{Z}_2$	4
$D_{2,c}$	0	1
$D_{3,l}$	0	1
$D_{3,s}$	0	1
D_4	\mathbb{Z}_2	2
D_6	0	1

How do the extensions look? Are they all non-isomorphic?

Lemma

Two extensions with different point groups and/or actions are non-isomorphic.

Lemma

Two extensions with different point groups and/or actions are non-isomorphic.

Proof.

-



Lemma

Two extensions with different point groups and/or actions are non-isomorphic.

Proof.

-



It follows that we have at least 13 non-isomorphic wallpaper groups, namely the semidirect products $T \rtimes_{\varphi} G_0$ for each point group G_0 and each action $\varphi : G_0 \rightarrow GL(\mathbb{Z}^2)$.

Lemma

Two extensions with different point groups and/or actions are non-isomorphic.

Proof.

-



It follows that we have at least 13 non-isomorphic wallpaper groups, namely the semidirect products $T \rtimes_{\varphi} G_0$ for each point group G_0 and each action $\varphi : G_0 \rightarrow GL(\mathbb{Z}^2)$.

Let us see what the corresponding patterns look like.


p1 - Just translations

G_0	C_1
G	$\mathbb{Z}^2 \rtimes C_1 = \mathbb{Z}_2$




pm - 1 reflection

G_0	D_1
G	$\mathbb{Z}^2 \rtimes D_{1,p}$



cm - 1 reflection, 1 glide reflection

G_0	D_1
G	$\mathbb{Z}^2 \rtimes D_{1,c}$




p2 - 180° rotation

G_0	C_2
G	$\mathbb{Z}^2 \rtimes C_2$



pmm - Two reflections and a 180° rotation only in the axes intersection

G_0	D_2
G	$\mathbb{Z}^2 \rtimes D_{2,p}$




cmm - Two reflections and a 180° rotation NOT in the axes intersection

G_0	D_2
G	$\mathbb{Z}^2 \rtimes D_{2,c}$




p3 - 120° rotation

G_0	C_3
G	$\mathbb{Z}^2 \rtimes C_3$



p3m1 - 120° rotation around the intersection of 3 reflection axes

G_0	D_3
G	$\mathbb{Z}^2 \rtimes D_{3,l}$



p31m - 120° rotation around a point, which is NOT an intersection of axes

G_0	D_3
G	$\mathbb{Z}^2 \rtimes D_{3,s}$




p4 - 90° rotation

G_0	C_4
G	$\mathbb{Z}^2 \rtimes C_4$



p4m - 90° rotation and 4 reflection axes

G_0	D_4
G	$\mathbb{Z}^2 \rtimes D_4$




p6 - 60° rotation

G_0	C_6
G	$\mathbb{Z}^2 \rtimes C_6$



p6m - 60° rotation and 6 reflection axes

G_0	D_6
G	$\mathbb{Z}^2 \rtimes D_6$



Let our point group be $D_{1,p}$.

Let our point group be $D_{1,p}$. I.e., $G_0 = D_1 = C_2 = \{1, g\}$ with action given by the matrix

$$\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

Non-split extensions

Let our point group be $D_{1,p}$. I.e., $G_0 = D_1 = C_2 = \{1, g\}$ with action given by the matrix

$$\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

As $H^2(D_{1,p}, T) = \mathbb{Z}_2$, we know that there is one non-split extension. What is it?

- Fix basis $\{t_1, t_2\}$ such that $g(t_1) = t_1$ and $g(t_2) = -t_2$.

- Fix basis $\{t_1, t_2\}$ such that $g(t_1) = t_1$ and $g(t_2) = -t_2$.
- Recall: As G is a subgroup of $\text{Iso}(\mathbb{R}^2) = \mathbb{R}^2 \rtimes \text{Aut}(\mathbb{R}^2)$, each element of G can be written as (g, t_g) , where $g \in G_0$, and $t_g \in \mathbb{R}^2$.

- Fix basis $\{t_1, t_2\}$ such that $g(t_1) = t_1$ and $g(t_2) = -t_2$.
- Recall: As G is a subgroup of $\text{Iso}(\mathbb{R}^2) = \mathbb{R}^2 \rtimes \text{Aut}(\mathbb{R}^2)$, each element of G can be written as (g, t_g) , where $g \in G_0$, and $t_g \in \mathbb{R}^2$.
- $t_g \in \mathbb{R}^2$ be such that $(g, t_g) \in G$.

- Fix basis $\{t_1, t_2\}$ such that $g(t_1) = t_1$ and $g(t_2) = -t_2$.
- Recall: As G is a subgroup of $\text{Iso}(\mathbb{R}^2) = \mathbb{R}^2 \rtimes \text{Aut}(\mathbb{R}^2)$, each element of G can be written as (g, t_g) , where $g \in G_0$, and $t_g \in \mathbb{R}^2$.
- $t_g \in \mathbb{R}^2$ be such that $(g, t_g) \in G$.
- Compute the 2-cocycle $c(g, g)$:

$$\begin{aligned} c(g, g) &= (g, t_g)(g, t_g)(1, t_1)^{-1} = (g^2, t_g + g(t_g))(1, t_1)^{-1} = \\ &= t_g + g(t_g) - t_1. \end{aligned}$$

- Fix basis $\{t_1, t_2\}$ such that $g(t_1) = t_1$ and $g(t_2) = -t_2$.
- Recall: As G is a subgroup of $\text{Iso}(\mathbb{R}^2) = \mathbb{R}^2 \rtimes \text{Aut}(\mathbb{R}^2)$, each element of G can be written as (g, t_g) , where $g \in G_0$, and $t_g \in \mathbb{R}^2$.
- $t_g \in \mathbb{R}^2$ be such that $(g, t_g) \in G$.
- Compute the 2-cocycle $c(g, g)$:

$$\begin{aligned} c(g, g) &= (g, t_g)(g, t_g)(1, t_1)^{-1} = (g^2, t_g + g(t_g))(1, t_1)^{-1} = \\ &= t_g + g(t_g) - t_1. \end{aligned}$$

- Let $t_g = \alpha t_1 + \beta t_2$. As $c(g, g), t_1 \in T$, we have that $t_g + g(t_g) \in T$.

- Fix basis $\{t_1, t_2\}$ such that $g(t_1) = t_1$ and $g(t_2) = -t_2$.
- Recall: As G is a subgroup of $\text{Iso}(\mathbb{R}^2) = \mathbb{R}^2 \rtimes \text{Aut}(\mathbb{R}^2)$, each element of G can be written as (g, t_g) , where $g \in G_0$, and $t_g \in \mathbb{R}^2$.
- $t_g \in \mathbb{R}^2$ be such that $(g, t_g) \in G$.
- Compute the 2-cocycle $c(g, g)$:

$$\begin{aligned} c(g, g) &= (g, t_g)(g, t_g)(1, t_1)^{-1} = (g^2, t_g + g(t_g))(1, t_1)^{-1} = \\ &= t_g + g(t_g) - t_1. \end{aligned}$$

- Let $t_g = \alpha t_1 + \beta t_2$. As $c(g, g), t_1 \in T$, we have that $t_g + g(t_g) \in T$.
- Hence, $(2t_1, 0) \in T$, and thus α is modulo T either 0 or $\frac{1}{2}$.

- Fix basis $\{t_1, t_2\}$ such that $g(t_1) = t_1$ and $g(t_2) = -t_2$.
- Recall: As G is a subgroup of $\text{Iso}(\mathbb{R}^2) = \mathbb{R}^2 \rtimes \text{Aut}(\mathbb{R}^2)$, each element of G can be written as (g, t_g) , where $g \in G_0$, and $t_g \in \mathbb{R}^2$.
- $t_g \in \mathbb{R}^2$ be such that $(g, t_g) \in G$.
- Compute the 2-cocycle $c(g, g)$:

$$\begin{aligned} c(g, g) &= (g, t_g)(g, t_g)(1, t_1)^{-1} = (g^2, t_g + g(t_g))(1, t_1)^{-1} = \\ &= t_g + g(t_g) - t_1. \end{aligned}$$

- Let $t_g = \alpha t_1 + \beta t_2$. As $c(g, g), t_1 \in T$, we have that $t_g + g(t_g) \in T$.
- Hence, $(2t_1, 0) \in T$, and thus α is modulo T either 0 or $\frac{1}{2}$.
- Value of β is irrelevant as $t_1 = 0$ results in $c(g, g)$ being a 2-boundary.

- Choice of $\alpha = 0$ results in the split extension, that is, the semidirect product we know.

- Choice of $\alpha = 0$ results in the split extension, that is, the semidirect product we know.
- Choice $\alpha = \frac{1}{2}$:

- Choice of $\alpha = 0$ results in the split extension, that is, the semidirect product we know.
- Choice $\alpha = \frac{1}{2}$:

G is then generated by t_1, t_2 and $(g, \frac{1}{2}t_1)$. We claim that G is not isomorphic to $T \ltimes D_1$.

- Choice of $\alpha = 0$ results in the split extension, that is, the semidirect product we know.
- Choice $\alpha = \frac{1}{2}$:

G is then generated by t_1, t_2 and $(g, \frac{1}{2}t_1)$. We claim that G is not isomorphic to $T \ltimes D_1$.

- $T \ltimes D_1$ contains D_1 as a subgroup.

- Choice of $\alpha = 0$ results in the split extension, that is, the semidirect product we know.
- Choice $\alpha = \frac{1}{2}$:

G is then generated by t_1, t_2 and $(g, \frac{1}{2}t_1)$. We claim that G is not isomorphic to $T \ltimes D_1$.

- $T \ltimes D_1$ contains D_1 as a subgroup.
- For any $t \in T$, $(g, \frac{1}{2}t_1 + t)^2 = (1, t_1 + t + g(t))$ is not zero.

- Choice of $\alpha = 0$ results in the split extension, that is, the semidirect product we know.
- Choice $\alpha = \frac{1}{2}$:

G is then generated by t_1, t_2 and $(g, \frac{1}{2}t_1)$. We claim that G is not isomorphic to $T \ltimes D_1$.

- $T \ltimes D_1$ contains D_1 as a subgroup.
- For any $t \in T$, $(g, \frac{1}{2}t_1 + t)^2 = (1, t_1 + t + g(t))$ is not zero.
- Therefore, G does not contain any non-trivial element of finite order.

pg - 1 glide reflection

G_0	D_1
G	$\text{Span}(t_1, t_2, (g, \frac{1}{2}t_1)) \subseteq \text{Iso}(\mathbb{R}^2)$



Further non-split extensions

Point group $G_0 = D_4$ with the unique (standard) action.

Further non-split extensions

Point group $G_0 = D_4$ with the unique (standard) action.

- D_4 is generated by rotation r and reflection f .

Further non-split extensions

Point group $G_0 = D_4$ with the unique (standard) action.

- D_4 is generated by rotation r and reflection f .
- Basis t_1, t_2 of T such that $r(t_1) = t_2, r(t_2) = -t_1$ and $f(t_1) = -t_1, f(t_2) = -t_2$.

Further non-split extensions

Point group $G_0 = D_4$ with the unique (standard) action.

- D_4 is generated by rotation r and reflection f .
- Basis t_1, t_2 of T such that $r(t_1) = t_2, r(t_2) = -t_1$ and $f(t_1) = -t_1, f(t_2) = -t_2$.
- Fact: $H^2(C_i, T) = 0$. Therefore, $(r, 0) \in G$.

Further non-split extensions

Point group $G_0 = D_4$ with the unique (standard) action.

- D_4 is generated by rotation r and reflection f .
- Basis t_1, t_2 of T such that $r(t_1) = t_2, r(t_2) = -t_1$ and $f(t_1) = -t_1, f(t_2) = -t_2$.
- Fact: $H^2(C_i, T) = 0$. Therefore, $(r, 0) \in G$.
- Let $t_f \in \mathbb{R}^2$ be such that $(f, t_f) \in G$.

Further non-split extensions

Point group $G_0 = D_4$ with the unique (standard) action.

- D_4 is generated by rotation r and reflection f .
- Basis t_1, t_2 of T such that $r(t_1) = t_2, r(t_2) = -t_1$ and $f(t_1) = -t_1, f(t_2) = -t_2$.
- Fact: $H^2(C_i, T) = 0$. Therefore, $(r, 0) \in G$.
- Let $t_f \in \mathbb{R}^2$ be such that $(f, t_f) \in G$.
- Compute $(r, 0)(f, t_f) = (rf, r(t_f))$

Further non-split extensions

Point group $G_0 = D_4$ with the unique (standard) action.

- D_4 is generated by rotation r and reflection f .
- Basis t_1, t_2 of T such that $r(t_1) = t_2, r(t_2) = -t_1$ and $f(t_1) = -t_1, f(t_2) = -t_2$.
- Fact: $H^2(C_i, T) = 0$. Therefore, $(r, 0) \in G$.
- Let $t_f \in \mathbb{R}^2$ be such that $(f, t_f) \in G$.
- Compute $(r, 0)(f, t_f) = (rf, r(t_f))$
- Compute the 2-cocycle: $c(rf, r) = (rf, r(t_f))(r, 0)(f, t_f)$, as $rfr = f$.

Further non-split extensions

Point group $G_0 = D_4$ with the unique (standard) action.

- D_4 is generated by rotation r and reflection f .
- Basis t_1, t_2 of T such that $r(t_1) = t_2, r(t_2) = -t_1$ and $f(t_1) = -t_1, f(t_2) = -t_2$.
- Fact: $H^2(C_i, T) = 0$. Therefore, $(r, 0) \in G$.
- Let $t_f \in \mathbb{R}^2$ be such that $(f, t_f) \in G$.
- Compute $(r, 0)(f, t_f) = (rf, r(t_f))$
- Compute the 2-cocycle: $c(rf, r) = (rf, r(t_f))(r, 0)(f, t_f)$, as $rfr = f$.
- This equals to:
 $(rfr, r(t_f))(f, t_f) = (1, f(t_f) + r(t_f)) = r(t_f) - t_f$, an element of T .

- Let $t_f = \alpha t_1 + \beta t_2$. We have
$$r(t_f) - t_f = (-\alpha - \beta)t_1 + (\alpha - \beta)t_2.$$

- Let $t_f = \alpha t_1 + \beta t_2$. We have
$$r(t_f) - t_f = (-\alpha - \beta)t_1 + (\alpha - \beta)t_2.$$
- Modulo T , this yields two options: $\alpha = \beta = 0$ and $\alpha = \beta = \frac{1}{2}$.

- Let $t_f = \alpha t_1 + \beta t_2$. We have


$$r(t_f) - t_f = (-\alpha - \beta)t_1 + (\alpha - \beta)t_2.$$
- Modulo T , this yields two options: $\alpha = \beta = 0$ and
 $\alpha = \beta = \frac{1}{2}.$
- First option produces the semidirect product $T \rtimes D_4$, the second is a group generated by $t_1, t_2, r, (f, \frac{1}{2}(t_1 + t_2)).$

- Let $t_f = \alpha t_1 + \beta t_2$. We have

$$r(t_f) - t_f = (-\alpha - \beta)t_1 + (\alpha - \beta)t_2.$$
- Modulo T , this yields two options: $\alpha = \beta = 0$ and $\alpha = \beta = \frac{1}{2}$.
- First option produces the semidirect product $T \rtimes D_4$, the second is a group generated by $t_1, t_2, r, (f, \frac{1}{2}(t_1 + t_2))$.
- These two are non-isomorphic, as the non-split extension does not contain a copy of D_4 .

p4g - 90° rotation and 2 reflection axes and 2 glide reflections

G_0	D_4
G	$\text{Span}(t_1, t_2, r, (f, \frac{1}{2}(t_1 + t_2))) \subseteq \text{Iso}(\mathbb{R}^2)$



Almost done

Point group $D_{2,p}$, i.e. $G_0 = D_2$ with action given by matrices:

$$\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

Almost done

Point group $D_{2,p}$, i.e. $G_0 = D_2$ with action given by matrices:

$$\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

- The first matrix corresponds to 180° rotation r , the second to reflection f .

Almost done

Point group $D_{2,p}$, i.e. $G_0 = D_2$ with action given by matrices:

$$\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

- The first matrix corresponds to 180° rotation r , the second to reflection f .
- As in the last case, we get
$$c(rf, f) = (rf, r(t_f))(r, 0)(f, t_f) = r(t_f) - t_f \in T.$$

Almost done

Point group $D_{2,p}$, i.e. $G_0 = D_2$ with action given by matrices:

$$\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

- The first matrix corresponds to 180° rotation r , the second to reflection f .
- As in the last case, we get
$$c(rf, f) = (rf, r(t_f))(r, 0)(f, t_f) = r(t_f) - t_f \in T.$$
- Let t_1, t_2 is a basis of T satisfying $r(t_i) = -t_i$ and $f(t_1) = t_1, f(t_2) = -t_2$, and $t_f = \alpha t_1 + \beta t_2$.

Almost done

Point group $D_{2,p}$, i.e. $G_0 = D_2$ with action given by matrices:

$$\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

- The first matrix corresponds to 180° rotation r , the second to reflection f .
- As in the last case, we get $c(rf, f) = (rf, r(t_f))(r, 0)(f, t_f) = r(t_f) - t_f \in T$.
- Let t_1, t_2 is a basis of T satisfying $r(t_i) = -t_i$ and $f(t_1) = t_1, f(t_2) = -t_2$, and $t_f = \alpha t_1 + \beta t_2$.
- We obtain $r(t_f) - t_f = -2t_f \in T$. Modulo T , we have 4 options: $t_f = 0, t_f = \frac{1}{2}t_1, t_f = \frac{1}{2}t_2, t_f = \frac{1}{2}(t_1 + t_2)$.

Almost done

Point group $D_{2,p}$, i.e. $G_0 = D_2$ with action given by matrices:

$$\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

- The first matrix corresponds to 180° rotation r , the second to reflection f .
- As in the last case, we get $c(rf, f) = (rf, r(t_f))(r, 0)(f, t_f) = r(t_f) - t_f \in T$.
- Let t_1, t_2 is a basis of T satisfying $r(t_i) = -t_i$ and $f(t_1) = t_1, f(t_2) = -t_2$, and $t_f = \alpha t_1 + \beta t_2$.
- We obtain $r(t_f) - t_f = -2t_f \in T$. Modulo T , we have 4 options: $t_f = 0, t_f = \frac{1}{2}t_1, t_f = \frac{1}{2}t_2, t_f = \frac{1}{2}(t_1 + t_2)$.
- Of the 4 obtained extensions, the two middle ones are obviously isomorphic.

Almost done


Point group $D_{2,p}$, i.e. $G_0 = D_2$ with action given by matrices:

$$\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

- The first matrix corresponds to 180° rotation r , the second to reflection f .
- As in the last case, we get $c(rf, f) = (rf, r(t_f))(r, 0)(f, t_f) = r(t_f) - t_f \in T$.
- Let t_1, t_2 is a basis of T satisfying $r(t_i) = -t_i$ and $f(t_1) = t_1, f(t_2) = -t_2$, and $t_f = \alpha t_1 + \beta t_2$.
- We obtain $r(t_f) - t_f = -2t_f \in T$. Modulo T , we have 4 options: $t_f = 0, t_f = \frac{1}{2}t_1, t_f = \frac{1}{2}t_2, t_f = \frac{1}{2}(t_1 + t_2)$.
- Of the 4 obtained extensions, the two middle ones are obviously isomorphic.
- The three remaining extensions are pairwise non-isomorphic.

pmg - 180° rotation, one reflection, one glide reflection

G_0	D_2
G	$\text{Span}(t_1, t_2, r, (f, \frac{1}{2}t_1)) \subseteq \text{Iso}(\mathbb{R}^2)$



Spiegelung (S) * Symmetrie in Bezug auf vertikale Linie (90°)

Quelle: m. IX - 6.3

pgg - 180° rotation, no reflections, two glide reflections

G_0	D_2
G	$\text{Span}(t_1, t_2, r, (f, \frac{1}{2}(t_1 + t_2))) \subseteq \text{Iso}(\mathbb{R}^2)$



Result: 17 wallpaper patterns!

Thank you for bearing with me! :-)