

# Group cohomology and wallpaper groups - Part II

Pavel Čoupek

MFF UK

23. 11. 2014

# Key problem

Let  $G$  be a wallpaper group.

# Key problem

Let  $G$  be a wallpaper group. We have the following data:

# Key problem

Let  $G$  be a wallpaper group. We have the following data:

- ▶  $T \simeq \mathbb{Z}^2$  ... lattice of translations in  $G$ ,

# Key problem

Let  $G$  be a wallpaper group. We have the following data:

- ▶  $T \simeq \mathbb{Z}^2$  ... lattice of translations in  $G$ ,
- ▶  $G_0$  ... the *point group*, acting on  $T$ ,

# Key problem

Let  $G$  be a wallpaper group. We have the following data:

- ▶  $T \simeq \mathbb{Z}^2$  ... lattice of translations in  $G$ ,
- ▶  $G_0$  ... the *point group*, acting on  $T$ ,
- ▶  $G_0 \simeq G/T$ .

# Key problem

Let  $G$  be a wallpaper group. We have the following data:

- ▶  $T \simeq \mathbb{Z}^2$  ... lattice of translations in  $G$ ,
- ▶  $G_0$  ... the *point group*, acting on  $T$ ,
- ▶  $G_0 \simeq G/T$ .

## Question

How can be the group  $G$  recovered from the data?

# Key problem

Let  $G$  be a wallpaper group. We have the following data:

- ▶  $T \simeq \mathbb{Z}^2$  ... lattice of translations in  $G$ ,
- ▶  $G_0$  ... the *point group*, acting on  $T$ ,
- ▶  $G_0 \simeq G/T$ .

## Question

How can be the group  $G$  recovered from the data?

**Answer:** Via the *second cohomology group*  $H^2(G_0, T)$ .



## Definition

- ▶ A sequence of groups and group homomorphisms

$$\dots \xrightarrow{f^{n-2}} A^{n-1} \xrightarrow{f^{n-1}} A^n \xrightarrow{f^n} A^{n+1} \xrightarrow{f^{n+1}} \dots$$

is called an *exact sequence* if for every integer  $n$ ,  
 $\text{Ker } f^{n+1} = \text{Im } f^n$ .

## Definition

- ▶ A sequence of groups and group homomorphisms

$$\dots \xrightarrow{f^{n-2}} A^{n-1} \xrightarrow{f^{n-1}} A^n \xrightarrow{f^n} A^{n+1} \xrightarrow{f^{n+1}} \dots$$

is called an *exact sequence* if for every integer  $n$ ,  
 $\text{Ker } f^{n+1} = \text{Im } f^n$ .

- ▶ An exact sequence of the form

$$1 \longrightarrow A \longrightarrow B \longrightarrow C \longrightarrow 1$$

is called a *short exact sequence*, or an *extension of  $A$  by  $C$* .

## Definition

- ▶ A sequence of groups and group homomorphisms

$$\dots \xrightarrow{f^{n-2}} A^{n-1} \xrightarrow{f^{n-1}} A^n \xrightarrow{f^n} A^{n+1} \xrightarrow{f^{n+1}} \dots$$

is called an *exact sequence* if for every integer  $n$ ,  
 $\text{Ker } f^{n+1} = \text{Im } f^n$ .

- ▶ An exact sequence of the form

$$\begin{array}{ccccccccccccccc} \dots & \longrightarrow & 1 & \longrightarrow & 1 & \longrightarrow & A & \longrightarrow & B & \longrightarrow & C & \longrightarrow & 1 & \longrightarrow & 1 & \longrightarrow & \dots \\ & & & & \parallel & & \parallel & & \parallel & & \parallel & & \parallel & & & & \\ & & \dots & & A^{-2} & & A^{-1} & & A^0 & & A^1 & & A^2 & & \dots & & \end{array}$$

is called a *short exact sequence*, or an *extension of  $A$  by  $C$* .

(picture)

## Definition

Two extensions of  $A$  by  $C$

$$1 \longrightarrow A \xrightarrow{\alpha_i} B_i \xrightarrow{\beta_i} C \longrightarrow 1, \quad i = 1, 2$$

are *equivalent* if there exists a group isomorphism  $g : B_1 \longrightarrow B_2$  such that the diagram

$$\begin{array}{ccccccccc} 1 & \longrightarrow & A & \xrightarrow{\alpha_1} & B_1 & \xrightarrow{\beta_1} & C & \longrightarrow & 1 \\ & & \parallel & & \downarrow g & & \parallel & & \\ 1 & \longrightarrow & A & \xrightarrow{\alpha_2} & B_2 & \xrightarrow{\beta_2} & C & \longrightarrow & 1 \end{array}$$

is commutative, that is,  $g \circ \alpha_1 = \alpha_2$  and  $\beta_2 \circ g = \beta_1$ .

## Example

Consider two extensions

$$1 \longrightarrow \mathbb{Z}_3 \longrightarrow \mathbb{Z}_9 \longrightarrow \mathbb{Z}_3 \longrightarrow 1$$

$$1 \longrightarrow \mathbb{Z}_3 \longrightarrow \mathbb{Z}_9 \longrightarrow \mathbb{Z}_3 \longrightarrow 1$$

## Example

Consider two extensions

$$1 \longrightarrow \mathbb{Z}_3 \xrightarrow{a \mapsto 3a} \mathbb{Z}_9 \longrightarrow \mathbb{Z}_3 \longrightarrow 1$$

$$1 \longrightarrow \mathbb{Z}_3 \xrightarrow{a \mapsto 3a} \mathbb{Z}_9 \longrightarrow \mathbb{Z}_3 \longrightarrow 1$$

## Example

Consider two extensions

$$1 \longrightarrow \mathbb{Z}_3 \xrightarrow{a \mapsto 3a} \mathbb{Z}_9 \xrightarrow{b \mapsto b \% 3} \mathbb{Z}_3 \longrightarrow 1$$

$$1 \longrightarrow \mathbb{Z}_3 \xrightarrow{a \mapsto 3a} \mathbb{Z}_9 \xrightarrow{b \mapsto 2b \% 3} \mathbb{Z}_3 \longrightarrow 1$$



**Claim:** *These are not equivalent.*

$$\begin{array}{ccccccc}
 1 & \longrightarrow & \mathbb{Z}_3 & \xrightarrow{a \mapsto 3a} & \mathbb{Z}_9 & \xrightarrow{b \mapsto b \% 3} & \mathbb{Z}_3 \longrightarrow 1 \\
 & & \parallel & & \downarrow \text{?} \exists g & & \parallel \\
 1 & \longrightarrow & \mathbb{Z}_3 & \xrightarrow{a \mapsto 3a} & \mathbb{Z}_9 & \xrightarrow{b \mapsto 2b \% 3} & \mathbb{Z}_3 \longrightarrow 1
 \end{array}$$

$\circlearrowleft_1$        $\circlearrowleft_2$

**Claim:** *These are not equivalent.*

$$\begin{array}{ccccccc}
 1 & \longrightarrow & \mathbb{Z}_3 & \xrightarrow{a \mapsto 3a} & \mathbb{Z}_9 & \xrightarrow{b \mapsto b \% 3} & \mathbb{Z}_3 \longrightarrow 1 \\
 & & \parallel & \circlearrowleft_1 & \downarrow \text{?} \exists g \circlearrowleft_2 & & \parallel \\
 1 & \longrightarrow & \mathbb{Z}_3 & \xrightarrow{a \mapsto 3a} & \mathbb{Z}_9 & \xrightarrow{b \mapsto 2b \% 3} & \mathbb{Z}_3 \longrightarrow 1
 \end{array}$$

$$g : a \mapsto ka \text{ for some } k \in \{1, 2, 4, 5, 7, 8\}$$

**Claim:** *These are not equivalent.*

$$\begin{array}{ccccccc}
 1 & \longrightarrow & \mathbb{Z}_3 & \xrightarrow{a \mapsto 3a} & \mathbb{Z}_9 & \xrightarrow{b \mapsto b\%3} & \mathbb{Z}_3 \longrightarrow 1 \\
 & & \parallel & \circlearrowleft_1 & \downarrow \text{?} \exists g \circlearrowleft_2 & & \parallel \\
 1 & \longrightarrow & \mathbb{Z}_3 & \xrightarrow{a \mapsto 3a} & \mathbb{Z}_9 & \xrightarrow{b \mapsto 2b\%3} & \mathbb{Z}_3 \longrightarrow 1
 \end{array}$$

$$g : a \mapsto ka \text{ for some } k \in \{1, 2, 4, 5, 7, 8\}$$

$$1. \quad g(3a) = 3a, \quad a = 1, 2, 3$$

$$\Rightarrow g = [b \mapsto kb] \text{ for } k = 1, 4, 7$$

**Claim:** *These are not equivalent.*

$$\begin{array}{ccccccc}
 1 & \longrightarrow & \mathbb{Z}_3 & \xrightarrow{a \mapsto 3a} & \mathbb{Z}_9 & \xrightarrow{b \mapsto b \% 3} & \mathbb{Z}_3 \longrightarrow 1 \\
 & & \parallel & \circlearrowleft_1 & \downarrow \text{?} \exists g \circlearrowleft_2 & & \parallel \\
 1 & \longrightarrow & \mathbb{Z}_3 & \xrightarrow{a \mapsto 3a} & \mathbb{Z}_9 & \xrightarrow{b \mapsto 2b \% 3} & \mathbb{Z}_3 \longrightarrow 1
 \end{array}$$

$$g : a \mapsto ka \text{ for some } k \in \{1, 2, 4, 5, 7, 8\}$$

$$1. \quad g(3a) = 3a, \quad a = 1, 2, 3$$

$$\Rightarrow g = [b \mapsto kb] \text{ for } k = 1, 4, 7$$

$$2. \quad 2g(a) \% 3 = a \% 3, \quad a = 1, 2, 3$$

$$\Rightarrow g = [b \mapsto kb] \text{ for } k = 2, 5, 8$$

## Proposition

*Consider a group extension*

$$1 \longrightarrow T \xrightarrow{\subseteq} G \xrightarrow{\pi} G_0 \longrightarrow 1$$

*with  $T$  Abelian. For each  $g \in G_0$ , choose  $x_g \in G$  such that  $\pi(x_g) = g$ . Then*

$$g * t := x_g t x_g^{-1}, \quad g \in G_0, t \in T$$

*is a correctly and uniquely defined action of  $G_0$  on  $T$ .*

# Proof

# Proof

- ▶ **“uniquely defined”:**

# Proof

- ▶ **“uniquely defined”:**

$$\pi(x_g) = \pi(x'_g) = g \in G_0$$



# Proof

- ▶ **“uniquely defined”:**

$$\pi(x_g) = \pi(x'_g) = g \in G_0 \rightsquigarrow x'_g = x_g t', \quad t' \in T$$

# Proof

- ▶ “uniquely defined”:

$$\pi(x_g) = \pi(x'_g) = g \in G_0 \rightsquigarrow x'_g = x_g t', \quad t' \in T$$
$$t \in T \dots$$

# Proof

- ▶ “uniquely defined”:

$$\pi(x_g) = \pi(x'_g) = g \in G_0 \rightsquigarrow x'_g = x_g t', \quad t' \in T$$

$$t \in T \dots x'_g t (x'_g)^{-1}$$

# Proof

► “uniquely defined”:

$$\pi(x_g) = \pi(x'_g) = g \in G_0 \rightsquigarrow x'_g = x_g t', \quad t' \in T$$

$$t \in T \dots x'_g t (x'_g)^{-1} = x_g t' t t'^{-1} x_g^{-1}$$

# Proof

- ▶ “uniquely defined”:

$$\pi(x_g) = \pi(x'_g) = g \in G_0 \rightsquigarrow x'_g = x_g t', \quad t' \in T$$

$$t \in T \dots x'_g t (x'_g)^{-1} = x_g t' t t'^{-1} x_g^{-1} = x_g t x_g^{-1}$$

# Proof

- ▶ “uniquely defined”:

$$\pi(x_g) = \pi(x'_g) = g \in G_0 \rightsquigarrow x'_g = x_g t', \quad t' \in T$$

$$t \in T \dots x'_g t (x'_g)^{-1} = x_g t' t t'^{-1} x_g^{-1} = x_g t x_g^{-1}$$

- ▶ “action”:

# Proof

- ▶ “uniquely defined”:

$$\pi(x_g) = \pi(x'_g) = g \in G_0 \rightsquigarrow x'_g = x_g t', \quad t' \in T$$

$$t \in T \dots x'_g t (x'_g)^{-1} = x_g t' t t'^{-1} x_g^{-1} = x_g t x_g^{-1}$$

- ▶ “action”:

$$g * (h * t)$$

# Proof

- ▶ “uniquely defined”:

$$\pi(x_g) = \pi(x'_g) = g \in G_0 \rightsquigarrow x'_g = x_g t', \quad t' \in T$$

$$t \in T \dots x'_g t (x'_g)^{-1} = x_g t' t t'^{-1} x_g^{-1} = x_g t x_g^{-1}$$

- ▶ “action”:

$$g * (h * t) = x_g (x_h t x_h^{-1}) x_g^{-1}$$



# Proof

- ▶ “uniquely defined”:

$$\pi(x_g) = \pi(x'_g) = g \in G_0 \rightsquigarrow x'_g = x_g t', \quad t' \in T$$

$$t \in T \dots x'_g t (x'_g)^{-1} = x_g t' t t'^{-1} x_g^{-1} = x_g t x_g^{-1}$$

- ▶ “action”:

$$g * (h * t) = x_g (x_h t x_h^{-1}) x_g^{-1} = x_{gh} t x_{gh}^{-1}$$

# Proof

- ▶ “uniquely defined”:

$$\pi(x_g) = \pi(x'_g) = g \in G_0 \rightsquigarrow x'_g = x_g t', \quad t' \in T$$

$$t \in T \dots x'_g t (x'_g)^{-1} = x_g t' t t'^{-1} x_g^{-1} = x_g t x_g^{-1}$$

- ▶ “action”:

$$g * (h * t) = x_g (x_h t x_h^{-1}) x_g^{-1} = x_{gh} t x_{gh}^{-1} = (gh) * t$$

# Proof

► “uniquely defined”:

$$\pi(x_g) = \pi(x'_g) = g \in G_0 \rightsquigarrow x'_g = x_g t', \quad t' \in T$$

$$t \in T \dots x'_g t (x'_g)^{-1} = x_g t' t t'^{-1} x_g^{-1} = x_g t x_g^{-1}$$

► “action”:

$$g * (h * t) = x_g (x_h t x_h^{-1}) x_g^{-1} = x_{gh} t x_{gh}^{-1} = (gh) * t$$

But  $\pi(x_g x_h) = \pi(x_g) \pi(x_h) = gh$ , hence  $x_g x_h$  is correct choice of preimage of  $gh$ .

# Proof

- ▶ **“uniquely defined”**:

$$\pi(x_g) = \pi(x'_g) = g \in G_0 \rightsquigarrow x'_g = x_g t', \quad t' \in T$$

$$t \in T \dots x'_g t (x'_g)^{-1} = x_g t' t t'^{-1} x_g^{-1} = x_g t x_g^{-1}$$

- ▶ **“action”**:

$$g * (h * t) = x_g (x_h t x_h^{-1}) x_g^{-1} = x_{gh} t x_{gh}^{-1} = (gh) * t$$

But  $\pi(x_g x_h) = \pi(x_g) \pi(x_h) = gh$ , hence  $x_g x_h$  is correct choice of preimage of  $gh$ .

- ▶ **“compatibility”**:

# Proof

- **“uniquely defined”**:

$$\pi(x_g) = \pi(x'_g) = g \in G_0 \rightsquigarrow x'_g = x_g t', \quad t' \in T$$

$$t \in T \dots x'_g t (x'_g)^{-1} = x_g t' t t'^{-1} x_g^{-1} = x_g t x_g^{-1}$$

- **“action”**:

$$g * (h * t) = x_g (x_h t x_h^{-1}) x_g^{-1} = x_{gh} t x_{gh}^{-1} = (gh) * t$$

But  $\pi(x_g x_h) = \pi(x_g) \pi(x_h) = gh$ , hence  $x_g x_h$  is correct choice of preimage of  $gh$ .

- **“compatibility”**:  $((g * -)$  is automorphism of  $T$  for every  $g)$

# Proof

- ▶ **“uniquely defined”**:

$$\pi(x_g) = \pi(x'_g) = g \in G_0 \rightsquigarrow x'_g = x_g t', \quad t' \in T$$

$$t \in T \dots x'_g t (x'_g)^{-1} = x_g t' t t'^{-1} x_g^{-1} = x_g t x_g^{-1}$$

- ▶ **“action”**:

$$g * (h * t) = x_g (x_h t x_h^{-1}) x_g^{-1} = x_{gh} t x_{gh}^{-1} = (gh) * t$$

But  $\pi(x_g x_h) = \pi(x_g) \pi(x_h) = gh$ , hence  $x_g x_h$  is correct choice of preimage of  $gh$ .

- ▶ **“compatibility”**:  $((g * -)$  is automorphism of  $T$  for every  $g)$

$$(x_g t x_g^{-1})(x_g t' x_g^{-1})$$

# Proof

- ▶ **“uniquely defined”**:

$$\pi(x_g) = \pi(x'_g) = g \in G_0 \rightsquigarrow x'_g = x_g t', \quad t' \in T$$

$$t \in T \dots x'_g t (x'_g)^{-1} = x_g t' t t'^{-1} x_g^{-1} = x_g t x_g^{-1}$$

- ▶ **“action”**:

$$g * (h * t) = x_g (x_h t x_h^{-1}) x_g^{-1} = x_{gh} t x_{gh}^{-1} = (gh) * t$$

But  $\pi(x_g x_h) = \pi(x_g) \pi(x_h) = gh$ , hence  $x_g x_h$  is correct choice of preimage of  $gh$ .

- ▶ **“compatibility”**:  $((g * -)$  is automorphism of  $T$  for every  $g)$

$$(x_g t x_g^{-1})(x_g t' x_g^{-1}) = x_g t (x_g x_g^{-1}) t' x_g^{-1}$$

# Proof

- ▶ **“uniquely defined”**:

$$\pi(x_g) = \pi(x'_g) = g \in G_0 \rightsquigarrow x'_g = x_g t', \quad t' \in T$$

$$t \in T \dots x'_g t (x'_g)^{-1} = x_g t' t t'^{-1} x_g^{-1} = x_g t x_g^{-1}$$

- ▶ **“action”**:

$$g * (h * t) = x_g (x_h t x_h^{-1}) x_g^{-1} = x_{gh} t x_{gh}^{-1} = (gh) * t$$

But  $\pi(x_g x_h) = \pi(x_g) \pi(x_h) = gh$ , hence  $x_g x_h$  is correct choice of preimage of  $gh$ .

- ▶ **“compatibility”**:  $((g * -)$  is automorphism of  $T$  for every  $g)$

$$(x_g t x_g^{-1})(x_g t' x_g^{-1}) = x_g t (x_g x_g^{-1}) t' x_g^{-1} = x_g t t' x_g^{-1}$$



## Remarks

1. Assume  $T$  is the lattice of wallpaper group  $G$  with point group  $G_0$ .

## Remarks

1. Assume  $T$  is the lattice of wallpaper group  $G$  with point group  $G_0$ .  $\leadsto$  We get an extension:

$$1 \longrightarrow T \longrightarrow G \longrightarrow G_0 \longrightarrow 1$$

## Remarks

1. Assume  $T$  is the lattice of wallpaper group  $G$  with point group  $G_0$ .  $\rightsquigarrow$  We get an extension:

$$1 \longrightarrow T \longrightarrow G \longrightarrow G_0 \longrightarrow 1$$

Then the given action of  $G_0$  on  $T$  corresponds to the one induced by the extension.

## Remarks

1. Assume  $T$  is the lattice of wallpaper group  $G$  with point group  $G_0$ .  $\rightsquigarrow$  We get an extension:

$$1 \longrightarrow T \longrightarrow G \longrightarrow G_0 \longrightarrow 1$$

$A$

Then the given action of  $G_0$  on  $T$  corresponds to the one induced by the extension.

## Remarks

1. Assume  $T$  is the lattice of wallpaper group  $G$  with point group  $G_0$ .  $\rightsquigarrow$  We get an extension:

$$1 \longrightarrow T \longrightarrow G \longrightarrow G_0 \longrightarrow 1$$

$$(A, v) \longleftarrow A$$

Then the given action of  $G_0$  on  $T$  corresponds to the one induced by the extension.

## Remarks

1. Assume  $T$  is the lattice of wallpaper group  $G$  with point group  $G_0$ .  $\rightsquigarrow$  We get an extension:

$$1 \longrightarrow T \longrightarrow G \longrightarrow G_0 \longrightarrow 1$$

$$(A, v) \longleftarrow A$$

$$(\text{Id}, w)$$

Then the given action of  $G_0$  on  $T$  corresponds to the one induced by the extension.

## Remarks

1. Assume  $T$  is the lattice of wallpaper group  $G$  with point group  $G_0$ .  $\rightsquigarrow$  We get an extension:

$$1 \longrightarrow T \longrightarrow G \longrightarrow G_0 \longrightarrow 1$$

$$(A, v) \longleftarrow A$$

$$(A, v)(\text{Id}, w)(A, v)^{-1}$$

Then the given action of  $G_0$  on  $T$  corresponds to the one induced by the extension.

## Remarks

1. Assume  $T$  is the lattice of wallpaper group  $G$  with point group  $G_0$ .  $\rightsquigarrow$  We get an extension:

$$1 \longrightarrow T \longrightarrow G \longrightarrow G_0 \longrightarrow 1$$

$$(A, v) \longleftarrow A$$

$$(A, v)(\text{Id}, w)(A, v)^{-1} = (\text{Id}, Aw)$$

Then the given action of  $G_0$  on  $T$  corresponds to the one induced by the extension.



## Remarks

### 2. In the extension

$$1 \longrightarrow T \longrightarrow G \longrightarrow G_0 \longrightarrow 1$$

is  $G$  of the form  $G = T \rtimes G_0$

## Remarks

### 2. In the extension

$$1 \longrightarrow T \longrightarrow G \longrightarrow G_0 \longrightarrow 1$$

is  $G$  of the form  $G = T \rtimes G_0$  if and only if the map  $g \mapsto x_g$  can be chosen as homomorphism.

## Remarks

### 2. In the extension

$$1 \longrightarrow T \longrightarrow G \longrightarrow G_0 \longrightarrow 1$$

is  $G$  of the form  $G = T \rtimes G_0$  if and only if the map  $g \mapsto x_g$  can be chosen as homomorphism.

(Then  $G = T \rtimes_{\varphi} G_0$ , where  $\varphi : g \mapsto (g * -)$ .)

## Remarks

### 2. In the extension

$$1 \longrightarrow T \longrightarrow G \longrightarrow G_0 \longrightarrow 1$$

is  $G$  of the form  $G = T \rtimes G_0$  if and only if the map  $g \mapsto x_g$  can be chosen as homomorphism.

(Then  $G = T \rtimes_{\varphi} G_0$ , where  $\varphi : g \mapsto (g * -)$ .)

In particular, this “trivial” extension always exists.

## Definition

- ▶ A sequence of Abelian groups

$$(\mathcal{A}) \quad \dots \xrightarrow{d^{n-2}} A^{n-1} \xrightarrow{d^{n-1}} A^n \xrightarrow{d^n} A^{n+1} \xrightarrow{d^{n+1}} \dots$$

is called a *chain complex* if for every  $n \in \mathbb{Z}$ ,  
 $\text{Im } d^n \subseteq \text{Ker } d^{n+1}$ , or, equivalently,  $d^{n+1} \circ d^n = 0$ .

## Definition

- ▶ A sequence of Abelian groups

$$(\mathcal{A}) \quad \dots \xrightarrow{d^{n-2}} A^{n-1} \xrightarrow{d^{n-1}} A^n \xrightarrow{d^n} A^{n+1} \xrightarrow{d^{n+1}} \dots$$

is called a *chain complex* if for every  $n \in \mathbb{Z}$ ,  
 $\text{Im } d^n \subseteq \text{Ker } d^{n+1}$ , or, equivalently,  $d^{n+1} \circ d^n = 0$ .

- ▶ For a chain complex  $(\mathcal{A})$  and  $n \in \mathbb{Z}$ , we define the *n-th cohomology group of  $(\mathcal{A})$*  by

$$\mathbb{H}^n(\mathcal{A}) = \text{Ker } d^n / \text{Im } d^{n-1}.$$

(picture)

## Proposition ( $\pm$ Snake lemma)

Let  $0 \rightarrow \mathcal{A} \rightarrow \mathcal{B} \rightarrow \mathcal{C} \rightarrow 0$  be a short exact sequence of chain complexes - that is, a commutative diagram

$$\begin{array}{ccccccc}
 & 0 & & 0 & & 0 & & 0 \\
 & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & A^{n-2} & \longrightarrow & A^{n-1} & \longrightarrow & A^n & \longrightarrow & A^{n+1} & \longrightarrow & \cdots & (A) \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & B^{n-2} & \longrightarrow & B^{n-1} & \longrightarrow & B^n & \longrightarrow & B^{n+1} & \longrightarrow & \cdots & (B) \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & C^{n-2} & \longrightarrow & C^{n-1} & \longrightarrow & C^n & \longrightarrow & C^{n+1} & \longrightarrow & \cdots & (C) \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0 & & 0
 \end{array}$$

such that the rows are complexes  $\mathcal{A}, \mathcal{B}, \mathcal{C}$  and the columns are exact. Then there is an exact sequence of the following form:

$$\cdots \longrightarrow \mathbb{H}^{n-1}(\mathcal{C}) \longrightarrow \mathbb{H}^n(\mathcal{A}) \longrightarrow \mathbb{H}^n(\mathcal{B}) \longrightarrow \mathbb{H}^n(\mathcal{C}) \longrightarrow \mathbb{H}^{n+1}(\mathcal{A}) \longrightarrow \cdots$$



# Idea of proof

$$\begin{array}{ccccccc}
 & 0 & & 0 & & 0 & & 0 \\
 & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & A^{n-1} & \longrightarrow & A^n & \longrightarrow & A^{n+1} & \longrightarrow & A^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \alpha_n & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & B^{n-1} & \longrightarrow & B^n & \longrightarrow & B^{n+1} & \longrightarrow & B^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \beta_n & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & C^{n-1} & \longrightarrow & C^n & \longrightarrow & C^{n+1} & \longrightarrow & C^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0 & & 0
 \end{array}$$

$$\cdots \longrightarrow \mathbb{H}^n(\mathcal{A}) \longrightarrow \mathbb{H}^n(\mathcal{B}) \longrightarrow \mathbb{H}^n(\mathcal{C}) \longrightarrow \mathbb{H}^{n+1}(\mathcal{A}) \longrightarrow \cdots$$

# Idea of proof

$$\begin{array}{ccccccc}
 & 0 & & 0 & & 0 & & 0 \\
 & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & A^{n-1} & \longrightarrow & A^n & \longrightarrow & A^{n+1} & \longrightarrow & A^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \alpha_n & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & B^{n-1} & \longrightarrow & B^n & \longrightarrow & B^{n+1} & \longrightarrow & B^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \beta_n & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & C^{n-1} & \longrightarrow & C^n & \longrightarrow & C^{n+1} & \longrightarrow & C^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0 & & 0
 \end{array}$$
  

$$\cdots \longrightarrow \mathbb{H}^n(\mathcal{A}) \xrightarrow{\quad \text{red arrow} \quad} \mathbb{H}^n(\mathcal{B}) \longrightarrow \mathbb{H}^n(\mathcal{C}) \longrightarrow \mathbb{H}^{n+1}(\mathcal{A}) \longrightarrow \cdots$$

# Idea of proof

$$\begin{array}{ccccccc}
 & 0 & & 0 & & 0 & & 0 \\
 & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & A^{n-1} & \longrightarrow & A^n & \longrightarrow & A^{n+1} & \longrightarrow & A^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \alpha_n & & \downarrow & & \downarrow & & \\
 \cdots & \longrightarrow & B^{n-1} & \longrightarrow & B^n & \longrightarrow & B^{n+1} & \longrightarrow & B^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \beta_n & & \downarrow & & \downarrow & & \\
 \cdots & \longrightarrow & C^{n-1} & \longrightarrow & C^n & \longrightarrow & C^{n+1} & \longrightarrow & C^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \\
 & & 0 & & 0 & & 0 & & & & \\
 & & \overline{a} & & & & & & & & \\
 \cdots & \longrightarrow & \mathbb{H}^n(\mathcal{A}) & \xrightarrow{\quad} & \mathbb{H}^n(\mathcal{B}) & \longrightarrow & \mathbb{H}^n(\mathcal{C}) & \longrightarrow & \mathbb{H}^{n+1}(\mathcal{A}) & \longrightarrow & \cdots
 \end{array}$$

# Idea of proof

$$\begin{array}{ccccccc}
 & 0 & & 0 & & 0 & & 0 \\
 & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & A^{n-1} & \longrightarrow & A^n & \longrightarrow & A^{n+1} & \longrightarrow & A^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \alpha_n & & \downarrow & & \downarrow & & \\
 \cdots & \longrightarrow & B^{n-1} & \longrightarrow & B^n & \longrightarrow & B^{n+1} & \longrightarrow & B^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \beta_n & & \downarrow & & \downarrow & & \\
 \cdots & \longrightarrow & C^{n-1} & \longrightarrow & C^n & \longrightarrow & C^{n+1} & \longrightarrow & C^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \\
 & & 0 & & 0 & & 0 & & & & \\
 & & & & & & & & & & \\
 & & \bar{a} & & & & & & & & \\
 \cdots & \longrightarrow & \mathbb{H}^n(\mathcal{A}) & \xrightarrow{\quad} & \mathbb{H}^n(\mathcal{B}) & \longrightarrow & \mathbb{H}^n(\mathcal{C}) & \longrightarrow & \mathbb{H}^{n+1}(\mathcal{A}) & \longrightarrow & \cdots
 \end{array}$$

# Idea of proof

$$\begin{array}{ccccccc}
 & 0 & & 0 & & 0 & & 0 \\
 & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & A^{n-1} & \longrightarrow & A^n & \longrightarrow & A^{n+1} & \longrightarrow & A^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \alpha_n^a & & \downarrow & & \downarrow & & \\
 \cdots & \longrightarrow & B^{n-1} & \longrightarrow & B^n & \longrightarrow & B^{n+1} & \longrightarrow & B^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \alpha_n(a) \beta_n & & \downarrow & & \downarrow & & \\
 \cdots & \longrightarrow & C^{n-1} & \longrightarrow & C^n & \longrightarrow & C^{n+1} & \longrightarrow & C^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \\
 & & 0 & & 0 & & 0 & & & & \\
 & & & & & & & & & & \\
 & & \bar{a} & & & & & & & & \\
 \cdots & \longrightarrow & \mathbb{H}^n(\mathcal{A}) & \xrightarrow{\quad} & \mathbb{H}^n(\mathcal{B}) & \longrightarrow & \mathbb{H}^n(\mathcal{C}) & \longrightarrow & \mathbb{H}^{n+1}(\mathcal{A}) & \longrightarrow & \cdots
 \end{array}$$

# Idea of proof

$$\begin{array}{ccccccc}
 & 0 & & 0 & & 0 & & 0 \\
 & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & A^{n-1} & \longrightarrow & A^n & \longrightarrow & A^{n+1} & \longrightarrow & A^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \alpha_n^a & & \downarrow & & \downarrow & & \\
 \cdots & \longrightarrow & B^{n-1} & \longrightarrow & B^n & \longrightarrow & B^{n+1} & \longrightarrow & B^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \alpha_n(a) & & \downarrow & & \downarrow & & \\
 \cdots & \longrightarrow & C^{n-1} & \longrightarrow & C^n & \longrightarrow & C^{n+1} & \longrightarrow & C^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \\
 & & 0 & & 0 & & 0 & & & & \\
 & & & & & & & & & & \\
 & & \bar{a} & \longmapsto & \overline{\alpha_n(a)} & & & & & & \\
 \cdots & \longrightarrow & \mathbb{H}^n(\mathcal{A}) & \longrightarrow & \mathbb{H}^n(\mathcal{B}) & \longrightarrow & \mathbb{H}^n(\mathcal{C}) & \longrightarrow & \mathbb{H}^{n+1}(\mathcal{A}) & \longrightarrow & \cdots
 \end{array}$$

# Idea of proof

$$\begin{array}{ccccccc}
 & 0 & & 0 & & 0 & & 0 \\
 & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & A^{n-1} & \longrightarrow & A^n & \longrightarrow & A^{n+1} & \longrightarrow & A^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \alpha_n & & \downarrow & & \downarrow & & \\
 \cdots & \longrightarrow & B^{n-1} & \longrightarrow & B^n & \longrightarrow & B^{n+1} & \longrightarrow & B^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \beta_n & & \downarrow & & \downarrow & & \\
 \cdots & \longrightarrow & C^{n-1} & \longrightarrow & C^n & \longrightarrow & C^{n+1} & \longrightarrow & C^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \\
 & & 0 & & 0 & & 0 & & & & 
 \end{array}$$

$$\cdots \longrightarrow \mathbb{H}^n(\mathcal{A}) \longrightarrow \mathbb{H}^n(\mathcal{B}) \xrightarrow{\quad} \mathbb{H}^n(\mathcal{C}) \longrightarrow \mathbb{H}^{n+1}(\mathcal{A}) \longrightarrow \cdots$$

# Idea of proof

$$\begin{array}{ccccccc}
 & 0 & & 0 & & 0 & & 0 \\
 & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & A^{n-1} & \longrightarrow & A^n & \longrightarrow & A^{n+1} & \longrightarrow & A^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \alpha_n & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & B^{n-1} & \longrightarrow & B^n & \longrightarrow & B^{n+1} & \longrightarrow & B^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \beta_n & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & C^{n-1} & \longrightarrow & C^n & \longrightarrow & C^{n+1} & \longrightarrow & C^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \beta_n(b) & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0 & & 0
 \end{array}$$

$\bar{b} \longmapsto \overline{\beta_n(b)}$   
 $\cdots \longrightarrow \mathbb{H}^n(\mathcal{A}) \longrightarrow \mathbb{H}^n(\mathcal{B}) \xrightarrow{\quad} \mathbb{H}^n(\mathcal{C}) \longrightarrow \mathbb{H}^{n+1}(\mathcal{A}) \longrightarrow \cdots$



# Idea of proof

$$\begin{array}{ccccccc}
 & 0 & & 0 & & 0 & & 0 \\
 & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & A^{n-1} & \longrightarrow & A^n & \longrightarrow & A^{n+1} & \longrightarrow & A^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \alpha_n & & \downarrow & & \downarrow & & \\
 \cdots & \longrightarrow & B^{n-1} & \longrightarrow & B^n & \longrightarrow & B^{n+1} & \longrightarrow & B^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \beta_n & & \downarrow & & \downarrow & & \\
 \cdots & \longrightarrow & C^{n-1} & \longrightarrow & C^n & \longrightarrow & C^{n+1} & \longrightarrow & C^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \\
 & & 0 & & 0 & & 0 & & & & \\
 \\
 \cdots & \longrightarrow & \mathbb{H}^n(\mathcal{A}) & \longrightarrow & \mathbb{H}^n(\mathcal{B}) & \longrightarrow & \mathbb{H}^n(\mathcal{C}) & \xrightarrow{\quad} & \mathbb{H}^{n+1}(\mathcal{A}) & \longrightarrow & \cdots
 \end{array}$$

# Idea of proof

$$\begin{array}{ccccccc}
 & 0 & & 0 & & 0 & & 0 \\
 & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & A^{n-1} & \longrightarrow & A^n & \longrightarrow & A^{n+1} & \longrightarrow & A^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \alpha_n & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & B^{n-1} & \longrightarrow & B^n & \longrightarrow & B^{n+1} & \longrightarrow & B^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \beta_n & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & C^{n-1} & \longrightarrow & C^n & \longrightarrow & C^{n+1} & \longrightarrow & C^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0 & & 0
 \end{array}$$
  

$$\cdots \longrightarrow \mathbb{H}^n(\mathcal{A}) \longrightarrow \mathbb{H}^n(\mathcal{B}) \longrightarrow \overset{\bar{c}}{\mathbb{H}^n(\mathcal{C})} \longrightarrow \mathbb{H}^{n+1}(\mathcal{A}) \longrightarrow \cdots$$

# Idea of proof

$$\begin{array}{ccccccc}
 & 0 & & 0 & & 0 & & 0 \\
 & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & A^{n-1} & \longrightarrow & A^n & \longrightarrow & A^{n+1} & \longrightarrow & A^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \alpha_n & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & B^{n-1} & \longrightarrow & B^n & \longrightarrow & B^{n+1} & \longrightarrow & B^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \beta_n & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & C^{n-1} & \longrightarrow & C^n & \longrightarrow & C^{n+1} & \longrightarrow & C^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0 & & 0 \\
 & & & & \textcolor{red}{c} & & & & \\
 & & & & & & \textcolor{red}{\bar{c}} & & \\
 \cdots & \longrightarrow & \mathbb{H}^n(\mathcal{A}) & \longrightarrow & \mathbb{H}^n(\mathcal{B}) & \longrightarrow & \mathbb{H}^n(\mathcal{C}) & \textcolor{red}{\longrightarrow} & \mathbb{H}^{n+1}(\mathcal{A}) & \longrightarrow & \cdots
 \end{array}$$

# Idea of proof

$$\begin{array}{ccccccc}
 & 0 & & 0 & & 0 & & 0 \\
 & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & A^{n-1} & \longrightarrow & A^n & \longrightarrow & A^{n+1} & \longrightarrow & A^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \alpha_n & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & B^{n-1} & \xrightarrow{c'} & B^n & \longrightarrow & B^{n+1} & \longrightarrow & B^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \beta_n & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & C^{n-1} & \xrightarrow{c} & C^n & \longrightarrow & C^{n+1} & \longrightarrow & C^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0 & & 0
 \end{array}$$

$\cdots \longrightarrow \mathbb{H}^n(\mathcal{A}) \longrightarrow \mathbb{H}^n(\mathcal{B}) \xrightarrow{\bar{c}} \mathbb{H}^{n+1}(\mathcal{A}) \longrightarrow \cdots$

# Idea of proof

$$\begin{array}{ccccccc}
 & 0 & & 0 & & 0 & & 0 \\
 & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & A^{n-1} & \longrightarrow & A^n & \longrightarrow & A^{n+1} & \longrightarrow & A^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \alpha_n & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & B^{n-1} & \xrightarrow{c'} & B^n & \longrightarrow & B^{n+1} & \longrightarrow & B^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \beta_n & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & C^{n-1} & \xrightarrow{c} & C^n & \longrightarrow & C^{n+1} & \longrightarrow & C^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0 & & 0
 \end{array}$$

$\cdots \longrightarrow \mathbb{H}^n(\mathcal{A}) \longrightarrow \mathbb{H}^n(\mathcal{B}) \xrightarrow{\bar{c}} \mathbb{H}^{n+1}(\mathcal{A}) \longrightarrow \cdots$

# Idea of proof

$$\begin{array}{ccccccc}
 & 0 & & 0 & & 0 & & 0 \\
 & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & A^{n-1} & \longrightarrow & A^n & \longrightarrow & A^{n+1} & \longrightarrow & A^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \alpha_n & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & B^{n-1} & \xrightarrow{c'} & B^n & \xrightarrow{d^n(c')} & B^{n+1} & \longrightarrow & B^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \beta_n & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & C^{n-1} & \xrightarrow{c} & C^n & \longrightarrow & C^{n+1} & \longrightarrow & C^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0 & & 0
 \end{array}$$

$\cdots \longrightarrow \mathbb{H}^n(\mathcal{A}) \longrightarrow \mathbb{H}^n(\mathcal{B}) \xrightarrow{\bar{c}} \mathbb{H}^{n+1}(\mathcal{A}) \longrightarrow \cdots$

# Idea of proof

$$\begin{array}{ccccccc}
 & 0 & & 0 & & 0 & & 0 \\
 & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & A^{n-1} & \longrightarrow & A^n & \longrightarrow & A^{n+1} & \longrightarrow & A^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \alpha_n & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & B^{n-1} & \xrightarrow{c'} & B^n & \xrightarrow{d^n(c')} & B^{n+1} & \longrightarrow & B^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \beta_n & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & C^{n-1} & \xrightarrow{c} & C^n & \longrightarrow & C^{n+1} & \longrightarrow & C^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0 & & 0
 \end{array}$$

$\cdots \longrightarrow \mathbb{H}^n(\mathcal{A}) \longrightarrow \mathbb{H}^n(\mathcal{B}) \xrightarrow{\bar{c}} \mathbb{H}^n(\mathcal{C}) \longrightarrow \mathbb{H}^{n+1}(\mathcal{A}) \longrightarrow \cdots$

# Idea of proof

$$\begin{array}{ccccccc}
 & 0 & & 0 & & 0 & & 0 \\
 & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & A^{n-1} & \longrightarrow & A^n & \longrightarrow & A^{n+1} & \longrightarrow & A^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \alpha_n & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & B^{n-1} & \xrightarrow{c'} & B^n & \xrightarrow{d^n(c')} & B^{n+1} & \longrightarrow & B^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \beta_n & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & C^{n-1} & \xrightarrow{c} & C^n & \longrightarrow & C^{n+1} & \longrightarrow & C^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0 & & 0
 \end{array}$$

$\cdots \longrightarrow \mathbb{H}^n(\mathcal{A}) \longrightarrow \mathbb{H}^n(\mathcal{B}) \xrightarrow{\bar{c}} \mathbb{H}^n(\mathcal{C}) \longrightarrow \mathbb{H}^{n+1}(\mathcal{A}) \longrightarrow \cdots$



# Idea of proof

$$\begin{array}{ccccccc}
 & 0 & & 0 & & 0 & & 0 \\
 & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & A^{n-1} & \longrightarrow & A^n & \longrightarrow & A^{n+1} & \longrightarrow & A^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \alpha_n & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & B^{n-1} & \xrightarrow{c'} & B^n & \xrightarrow{d^n(c')} & B^{n+1} & \longrightarrow & B^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \beta_n & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & C^{n-1} & \xrightarrow{c} & C^n & \xrightarrow{??} & C^{n+1} & \longrightarrow & C^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0 & & 0
 \end{array}$$

$\cdots \longrightarrow \mathbb{H}^n(\mathcal{A}) \longrightarrow \mathbb{H}^n(\mathcal{B}) \xrightarrow{\bar{c}} \mathbb{H}^n(\mathcal{C}) \longrightarrow \mathbb{H}^{n+1}(\mathcal{A}) \longrightarrow \cdots$

# Idea of proof

$$\begin{array}{ccccccc}
 & 0 & & 0 & & 0 & & 0 \\
 & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & A^{n-1} & \longrightarrow & A^n & \longrightarrow & A^{n+1} & \longrightarrow & A^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \alpha_n & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & B^{n-1} & \xrightarrow{c'} & B^n & \longrightarrow & B^{n+1} & \xrightarrow{d^n(c')} & B^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \beta_n & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & C^{n-1} & \longrightarrow & C^n & \longrightarrow & C^{n+1} & \longrightarrow & C^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0 & & 0
 \end{array}$$

$c$  (red) is below  $C^n$ ,  $??$  (blue) is below  $C^{n+1}$ . A dashed blue arrow points from  $C^n$  to  $C^{n+1}$ .

$$\cdots \longrightarrow \mathbb{H}^n(\mathcal{A}) \longrightarrow \mathbb{H}^n(\mathcal{B}) \xrightarrow{\bar{c}} \mathbb{H}^n(\mathcal{C}) \longrightarrow \mathbb{H}^{n+1}(\mathcal{A}) \longrightarrow \cdots$$



# Idea of proof

$$\begin{array}{ccccccc}
 & 0 & & 0 & & 0 & & 0 \\
 & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & A^{n-1} & \longrightarrow & A^n & \longrightarrow & A^{n+1} & \longrightarrow & A^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \alpha_n & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & B^{n-1} & \xrightarrow{c'} & B^n & \xrightarrow{d^n(c')} & B^{n+1} & \longrightarrow & B^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow \beta_n & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & C^{n-1} & \xrightarrow{c} & C^n & \longrightarrow & C^{n+1} & \longrightarrow & C^{n+2} & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0 & & 0
 \end{array}$$

$\cdots \longrightarrow \mathbb{H}^n(\mathcal{A}) \longrightarrow \mathbb{H}^n(\mathcal{B}) \xrightarrow{\bar{c}} \mathbb{H}^n(\mathcal{C}) \longrightarrow \mathbb{H}^{n+1}(\mathcal{A}) \longrightarrow \cdots$

# Idea of proof

$$\begin{array}{ccccccc}
 & 0 & & 0 & & 0 & & 0 \\
 & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & A^{n-1} & \longrightarrow & A^n & \longrightarrow & A^{n+1} & \xrightarrow{c''} A^{n+2} \longrightarrow \cdots \\
 & & \downarrow & & \downarrow \alpha_n & & \downarrow & \\
 \cdots & \longrightarrow & B^{n-1} & \xrightarrow{c'} & B^n & \xrightarrow{\quad} & B^{n+1} & \xrightarrow{d^n(c')} B^{n+2} \longrightarrow \cdots \\
 & & \downarrow & & \downarrow \beta_n & & \downarrow & \\
 \cdots & \longrightarrow & C^{n-1} & \xrightarrow{c} & C^n & \longrightarrow & C^{n+1} & \longrightarrow C^{n+2} \longrightarrow \cdots \\
 & & \downarrow & & \downarrow & & \downarrow & \\
 & & 0 & & 0 & & 0 & \\
 & & & & & & \bar{c} & \\
 \cdots & \longrightarrow & \mathbb{H}^n(\mathcal{A}) & \longrightarrow & \mathbb{H}^n(\mathcal{B}) & \longrightarrow & \mathbb{H}^n(\mathcal{C}) & \xrightarrow{\quad} \mathbb{H}^{n+1}(\mathcal{A}) \longrightarrow \cdots
 \end{array}$$

# Idea of proof

$$\begin{array}{ccccccc}
 & 0 & & 0 & & 0 & & 0 \\
 & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & A^{n-1} & \longrightarrow & A^n & \longrightarrow & A^{n+1} & \xrightarrow{c''} A^{n+2} \longrightarrow \cdots \\
 & & \downarrow & & \downarrow \alpha_n & & \downarrow & \\
 \cdots & \longrightarrow & B^{n-1} & \xrightarrow{c'} & B^n & \xrightarrow{\quad} & B^{n+1} & \xrightarrow{d^n(c')} B^{n+2} \longrightarrow \cdots \\
 & & \downarrow & & \downarrow \beta_n & & \downarrow & \\
 \cdots & \longrightarrow & C^{n-1} & \xrightarrow{c} & C^n & \longrightarrow & C^{n+1} & \longrightarrow C^{n+2} \longrightarrow \cdots \\
 & & \downarrow & & \downarrow & & \downarrow & \\
 & & 0 & & 0 & & 0 & \\
 & & & & & & & \\
 \cdots & \longrightarrow & \mathbb{H}^n(\mathcal{A}) & \longrightarrow & \mathbb{H}^n(\mathcal{B}) & \xrightarrow{\bar{c}} & \mathbb{H}^n(\mathcal{C}) & \xrightarrow{\quad} \mathbb{H}^{n+1}(\mathcal{A}) \longrightarrow \cdots
 \end{array}$$

The diagram illustrates the idea of proof for the relationship between group cohomology and group extensions. It shows a commutative diagram involving three rows of chain complexes and a bottom row of cohomology groups.

- The top row is a chain complex:  $\cdots \longrightarrow A^{n-1} \longrightarrow A^n \longrightarrow A^{n+1} \xrightarrow{c''} A^{n+2} \longrightarrow \cdots$ . Above each term is a zero object with a downward arrow.
- The middle row is a chain complex:  $\cdots \longrightarrow B^{n-1} \xrightarrow{c'} B^n \longrightarrow B^{n+1} \xrightarrow{d^n(c')} B^{n+2} \longrightarrow \cdots$ . Vertical arrows connect  $A^{n-1}$  to  $B^{n-1}$ ,  $A^n$  to  $B^n$  (labeled  $\alpha_n$ ), and  $A^{n+1}$  to  $B^{n+1}$ .
- The bottom row of the diagram is a chain complex:  $\cdots \longrightarrow C^{n-1} \xrightarrow{c} C^n \longrightarrow C^{n+1} \longrightarrow C^{n+2} \longrightarrow \cdots$ . Vertical arrows connect  $B^{n-1}$  to  $C^{n-1}$ ,  $B^n$  to  $C^n$  (labeled  $\beta_n$ ), and  $B^{n+1}$  to  $C^{n+1}$ .
- Below the chain complexes are zero objects:  $0$  below  $C^{n-1}$ ,  $0$  below  $C^n$ , and  $0$  below  $C^{n+1}$ .
- The bottom row of the diagram is a sequence of cohomology groups:  $\cdots \longrightarrow \mathbb{H}^n(\mathcal{A}) \longrightarrow \mathbb{H}^n(\mathcal{B}) \xrightarrow{\bar{c}} \mathbb{H}^n(\mathcal{C}) \xrightarrow{\quad} \mathbb{H}^{n+1}(\mathcal{A}) \longrightarrow \cdots$ . Red arrows indicate the mapping from  $\mathbb{H}^n(\mathcal{C})$  to  $\mathbb{H}^{n+1}(\mathcal{A})$ , with labels  $\bar{c}$  and  $\overline{c''}$ .

## Definition

Let  $T$  be an Abelian group and  $G_0$  a group with action on  $T$ .

For a positive integer  $n$ , denote  $C^n(G_0, T)$  the group of all maps

$f : (G_0)^n \rightarrow T$ , and define group homomorphisms

$d^n : C^n(G_0, T) \rightarrow C^{n+1}(G_0, T)$  by

$$\begin{aligned} [d^n(f)](g_1, \dots, g_{n+1}) = & g_1 * f(g_2, \dots, g_{n+1}) \\ & + \sum_{i=1}^n (-1)^i f(g_1, g_2, \dots, (g_i g_{i+1}), \dots, g_{n+1}) \\ & + (-1)^{n+1} f(g_1, g_2, \dots, g_n). \end{aligned}$$

Further set  $C^0(G_0, T) = T$  and

$d^0(g) : t \mapsto g * t - t, \quad g \in G_0, t \in T.$

## Proposition

*For every  $n$ ,  $d^{n+1} \circ d^n = 0$ . That is,  $C^n(G_0, T)$  together with  $d^n, n \in \mathbb{N}_0$ , form a chain complex.*



## Proposition

*For every  $n$ ,  $d^{n+1} \circ d^n = 0$ . That is,  $C^n(G_0, T)$  together with  $d^n, n \in \mathbb{N}_0$ , form a chain complex.*

(Proof is omitted.)

## Proposition

*For every  $n$ ,  $d^{n+1} \circ d^n = 0$ . That is,  $C^n(G_0, T)$  together with  $d^n, n \in \mathbb{N}_0$ , form a chain complex.*

(Proof is omitted.)

## Definition

Given a non-negative integer  $n$ , we define

$$H^n(G_0, T) = \mathbb{H}^n(C^\bullet(G_0, T)).$$

We call  $H^n(G_0, T)$  the  $n$ -th cohomology group of  $G_0$  with coefficients in  $T$ .

**Warning:** For the next few slides,  $T$  is written multiplicatively!

**Warning:** For the next few slides,  $T$  is written multiplicatively!  
In particular, the group  $H^2(G_0, T)$  consists of so-called

**Warning:** For the next few slides,  $T$  is written multiplicatively!  
In particular, the group  $H^2(G_0, T)$  consists of so-called

*2-cocycles*: Maps  $c : G_0 \times G_0 \rightarrow T$  satisfying

$$c(g, h) \cdot c(gh, k) = (g * c(h, k)) \cdot c(g, hk), \quad g, h, k \in G_0$$

**Warning:** For the next few slides,  $T$  is written multiplicatively!  
In particular, the group  $H^2(G_0, T)$  consists of so-called

*2-cocycles:* Maps  $c : G_0 \times G_0 \rightarrow T$  satisfying

$$c(g, h) \cdot c(gh, k) = (g * c(h, k)) \cdot c(g, hk), \quad g, h, k \in G_0$$

modulo

*2-coboundaries:* Maps  $b : G_0 \times G_0 \rightarrow T$  satisfying

$$b(g, h) = (g * f(h)) \cdot (f(gh))^{-1} \cdot f(g)$$

for some map  $f : G_0 \rightarrow T$ .

## Theorem

*There is a 1-1 correspondence between*

## Theorem

*There is a 1-1 correspondence between*

- ▶ *equivalence classes of extensions of  $T$  by  $G_0$  with the given action, and*



## Theorem

*There is a 1-1 correspondence between*

- ▶ *equivalence classes of extensions of  $T$  by  $G_0$  with the given action, and*
- ▶ *the elements of  $H^2(G_0, T)$ .*

# Proof $(\text{extensions of } T \text{ by } G_0 \iff H^2(G_0, T))$

## 1. Going forward:

# Proof $(\text{extensions of } T \text{ by } G_0 \iff H^2(G_0, T))$

1. **Going forward:** Consider an extension

$1 \rightarrow T \rightarrow G \xrightarrow{\pi} G_0 \rightarrow 1$ , fix representatives  $x_g \in \pi^{-1}(g)$ .

# Proof (extensions of $T$ by $G_0 \iff H^2(G_0, T)$ )

1. **Going forward:** Consider an extension  
 $1 \rightarrow T \rightarrow G \xrightarrow{\pi} G_0 \rightarrow 1$ , fix representatives  $x_g \in \pi^{-1}(g)$ .

Then

$$c(g, h) := x_g x_h x_{gh}^{-1}$$

is a 2-cocycle:

# Proof (extensions of $T$ by $G_0 \iff H^2(G_0, T)$ )

1. **Going forward:** Consider an extension

$1 \rightarrow T \rightarrow G \xrightarrow{\pi} G_0 \rightarrow 1$ , fix representatives  $x_g \in \pi^{-1}(g)$ .

Then

$$c(g, h) := x_g x_h x_{gh}^{-1}$$

is a 2-cocycle:

$$(g * c(h, k)) \cdot c(g, hk)$$

# Proof (extensions of $T$ by $G_0 \iff H^2(G_0, T)$ )

## 1. Going forward: Consider an extension

$1 \rightarrow T \rightarrow G \xrightarrow{\pi} G_0 \rightarrow 1$ , fix representatives  $x_g \in \pi^{-1}(g)$ .

Then

$$c(g, h) := x_g x_h x_{gh}^{-1}$$

is a 2-cocycle:

$$(g * c(h, k)) \cdot c(g, hk) = x_g (x_h x_k x_{hk}^{-1}) x_g^{-1} \cdot (x_g x_{hk} x_{ghk}^{-1})$$

# Proof (extensions of $T$ by $G_0 \iff H^2(G_0, T)$ )

## 1. Going forward: Consider an extension

$1 \rightarrow T \rightarrow G \xrightarrow{\pi} G_0 \rightarrow 1$ , fix representatives  $x_g \in \pi^{-1}(g)$ .

Then

$$c(g, h) := x_g x_h x_{gh}^{-1}$$

is a 2-cocycle:

$$\begin{aligned} (g * c(h, k)) \cdot c(g, hk) &= x_g (x_h x_k x_{hk}^{-1}) x_g^{-1} \cdot (x_g x_{hk} x_{ghk}^{-1}) \\ &= x_g x_h x_k x_{ghk}^{-1} \end{aligned}$$

# Proof (extensions of $T$ by $G_0 \iff H^2(G_0, T)$ )

## 1. Going forward: Consider an extension

$1 \rightarrow T \rightarrow G \xrightarrow{\pi} G_0 \rightarrow 1$ , fix representatives  $x_g \in \pi^{-1}(g)$ .

Then

$$c(g, h) := x_g x_h x_{gh}^{-1}$$

is a 2-cocycle:

$$\begin{aligned} (g * c(h, k)) \cdot c(g, hk) &= x_g (x_h x_k x_{hk}^{-1}) x_g^{-1} \cdot (x_g x_{hk} x_{ghk}^{-1}) \\ &= x_g x_h x_k x_{ghk}^{-1} = x_g x_h (x_{gh}^{-1} x_{gh}) x_k x_{ghk}^{-1} \end{aligned}$$



# Proof (extensions of $T$ by $G_0 \iff H^2(G_0, T)$ )

## 1. Going forward: Consider an extension

$1 \rightarrow T \rightarrow G \xrightarrow{\pi} G_0 \rightarrow 1$ , fix representatives  $x_g \in \pi^{-1}(g)$ .

Then

$$c(g, h) := x_g x_h x_{gh}^{-1}$$

is a 2-cocycle:

$$\begin{aligned} (g * c(h, k)) \cdot c(g, hk) &= x_g (x_h x_k x_{hk}^{-1}) x_g^{-1} \cdot (x_g x_{hk} x_{ghk}^{-1}) \\ &= x_g x_h x_k x_{ghk}^{-1} = x_g x_h (x_{gh}^{-1} x_{gh}) x_k x_{ghk}^{-1} \\ &= (x_g x_h x_{gh}^{-1}) (x_{gh} x_k x_{ghk}^{-1}) \end{aligned}$$

# Proof (extensions of $T$ by $G_0 \iff H^2(G_0, T)$ )

## 1. Going forward: Consider an extension

$1 \rightarrow T \rightarrow G \xrightarrow{\pi} G_0 \rightarrow 1$ , fix representatives  $x_g \in \pi^{-1}(g)$ .

Then

$$c(g, h) := x_g x_h x_{gh}^{-1}$$

is a 2-cocycle:

$$\begin{aligned} (g * c(h, k)) \cdot c(g, hk) &= x_g (x_h x_k x_{hk}^{-1}) x_g^{-1} \cdot (x_g x_{hk} x_{ghk}^{-1}) \\ &= x_g x_h x_k x_{ghk}^{-1} = x_g x_h (x_{gh}^{-1} x_{gh}) x_k x_{ghk}^{-1} \\ &= (x_g x_h x_{gh}^{-1}) (x_{gh} x_k x_{ghk}^{-1}) = c(g, h) \cdot c(gh, k) \end{aligned}$$

Proof  $(\text{extensions of } T \text{ by } G_0 \iff H^2(G_0, T))$

## 2. Going back:

Proof  $(\text{extensions of } T \text{ by } G_0 \iff H^2(G_0, T))$

2. **Going back:** Consider a 2-cocycle  $c(g, h)$ .

# Proof $(\text{extensions of } T \text{ by } G_0 \iff H^2(G_0, T))$

2. **Going back:** Consider a 2-cocycle  $c(g, h)$ .

On  $G := T \times G_0$ , define group operation by

$$(t, g) \cdot (s, h) := (t \cdot (g * s) \cdot c(g, h), gh).$$

# Proof (extensions of $T$ by $G_0 \iff H^2(G_0, T)$ )

2. **Going back:** Consider a 2-cocycle  $c(g, h)$ .

On  $G := T \times G_0$ , define group operation by

$$(t, g) \cdot (s, h) := (t \cdot (g * s) \cdot c(g, h), gh).$$

Then it can be verified that

- ▶  $(G, \cdot)$  is a group,

# Proof (extensions of $T$ by $G_0 \iff H^2(G_0, T)$ )

2. **Going back:** Consider a 2-cocycle  $c(g, h)$ .

On  $G := T \times G_0$ , define group operation by

$$(t, g) \cdot (s, h) := (t \cdot (g * s) \cdot c(g, h), gh).$$

Then it can be verified that

- ▶  $(G, \cdot)$  is a group,
- ▶  $T \simeq T \times \{1\} \trianglelefteq G$ ,

# Proof (extensions of $T$ by $G_0 \iff H^2(G_0, T)$ )

2. **Going back:** Consider a 2-cocycle  $c(g, h)$ .

On  $G := T \times G_0$ , define group operation by

$$(t, g) \cdot (s, h) := (t \cdot (g * s) \cdot c(g, h), gh).$$

Then it can be verified that

- ▶  $(G, \cdot)$  is a group,
- ▶  $T \simeq T \times \{1\} \trianglelefteq G$ ,
- ▶  $G/T \simeq G_0$ .



Proof  $(\text{extensions of } T \text{ by } G_0 \iff H^2(G_0, T))$

**What about 2-coboundaries?**

# Proof $(\text{extensions of } T \text{ by } G_0 \iff H^2(G_0, T))$

## What about 2-coboundaries?

Choose representatives  $\{x'_g\}_{g \in G_0}$  instead of  $\{x_g\}_{g \in G_0}$  to obtain a 2-cocycle  $c'(g, h)$ .

# Proof $(\text{extensions of } T \text{ by } G_0 \iff H^2(G_0, T))$

## What about 2-coboundaries?

Choose representatives  $\{x'_g\}_{g \in G_0}$  instead of  $\{x_g\}_{g \in G_0}$  to obtain a 2-cocycle  $c'(g, h)$ .

Then  $x'_g = t_g x_g$  for every  $g \in G_0$  and some  $t_g \in T$ .

# Proof (extensions of $T$ by $G_0 \iff H^2(G_0, T)$ )

## What about 2-coboundaries?

Choose representatives  $\{x'_g\}_{g \in G_0}$  instead of  $\{x_g\}_{g \in G_0}$  to obtain a 2-cocycle  $c'(g, h)$ .

Then  $x'_g = t_g x_g$  for every  $g \in G_0$  and some  $t_g \in T$ . It follows that

$$c'(g, h) = x'_g x'_h (x'_{gh})^{-1}$$

# Proof (extensions of $T$ by $G_0 \iff H^2(G_0, T)$ )

## What about 2-coboundaries?

Choose representatives  $\{x'_g\}_{g \in G_0}$  instead of  $\{x_g\}_{g \in G_0}$  to obtain a 2-cocycle  $c'(g, h)$ .

Then  $x'_g = t_g x_g$  for every  $g \in G_0$  and some  $t_g \in T$ . It follows that

$$c'(g, h) = x'_g x'_h (x'_{gh})^{-1} = (t_g x_g)(t_h x_h)(t_{gh} x_{gh})^{-1}$$

# Proof (extensions of $T$ by $G_0 \iff H^2(G_0, T)$ )

## What about 2-coboundaries?

Choose representatives  $\{x'_g\}_{g \in G_0}$  instead of  $\{x_g\}_{g \in G_0}$  to obtain a 2-cocycle  $c'(g, h)$ .

Then  $x'_g = t_g x_g$  for every  $g \in G_0$  and some  $t_g \in T$ . It follows that

$$\begin{aligned} c'(g, h) &= x'_g x'_h (x'_{gh})^{-1} = (t_g x_g)(t_h x_h)(t_{gh} x_{gh})^{-1} \\ &= t_g x_g t_h (x_g^{-1} x_g) x_h x_{gh}^{-1} t_{gh}^{-1} \end{aligned}$$

# Proof (extensions of $T$ by $G_0 \iff H^2(G_0, T)$ )

## What about 2-coboundaries?

Choose representatives  $\{x'_g\}_{g \in G_0}$  instead of  $\{x_g\}_{g \in G_0}$  to obtain a 2-cocycle  $c'(g, h)$ .

Then  $x'_g = t_g x_g$  for every  $g \in G_0$  and some  $t_g \in T$ . It follows that

$$\begin{aligned} c'(g, h) &= x'_g x'_h (x'_{gh})^{-1} = (t_g x_g)(t_h x_h)(t_{gh} x_{gh})^{-1} \\ &= t_g x_g t_h (x_g^{-1} x_g) x_h x_{gh}^{-1} t_{gh}^{-1} = t_g (x_g t_h x_g^{-1}) x_g x_h x_{gh}^{-1} t_{gh}^{-1} \end{aligned}$$

# Proof (extensions of $T$ by $G_0 \rightleftarrows H^2(G_0, T)$ )

## What about 2-coboundaries?

Choose representatives  $\{x'_g\}_{g \in G_0}$  instead of  $\{x_g\}_{g \in G_0}$  to obtain a 2-cocycle  $c'(g, h)$ .

Then  $x'_g = t_g x_g$  for every  $g \in G_0$  and some  $t_g \in T$ . It follows that

$$\begin{aligned} c'(g, h) &= x'_g x'_h (x'_{gh})^{-1} = (t_g x_g)(t_h x_h)(t_{gh} x_{gh})^{-1} \\ &= t_g x_g t_h (x_g^{-1} x_g) x_h x_{gh}^{-1} t_{gh}^{-1} = t_g (x_g t_h x_g^{-1}) x_g x_h x_{gh}^{-1} t_{gh}^{-1} \\ &= t_g (g * t_h) c(g, h) t_{gh}^{-1} \end{aligned}$$



# Proof (extensions of $T$ by $G_0 \iff H^2(G_0, T)$ )

## What about 2-coboundaries?

Choose representatives  $\{x'_g\}_{g \in G_0}$  instead of  $\{x_g\}_{g \in G_0}$  to obtain a 2-cocycle  $c'(g, h)$ .

Then  $x'_g = t_g x_g$  for every  $g \in G_0$  and some  $t_g \in T$ . It follows that

$$\begin{aligned} c'(g, h) &= x'_g x'_h (x'_{gh})^{-1} = (t_g x_g)(t_h x_h)(t_{gh} x_{gh})^{-1} \\ &= t_g x_g t_h (x_g^{-1} x_g) x_h x_{gh}^{-1} t_{gh}^{-1} = t_g (x_g t_h x_g^{-1}) x_g x_h x_{gh}^{-1} t_{gh}^{-1} \\ &= t_g (g * t_h) c(g, h) t_{gh}^{-1} = ((g * t_h) t_{gh}^{-1} t_g) c(g, h) \end{aligned}$$

# Proof (extensions of $T$ by $G_0 \iff H^2(G_0, T)$ )

## What about 2-coboundaries?

Choose representatives  $\{x'_g\}_{g \in G_0}$  instead of  $\{x_g\}_{g \in G_0}$  to obtain a 2-cocycle  $c'(g, h)$ .

Then  $x'_g = t_g x_g$  for every  $g \in G_0$  and some  $t_g \in T$ . It follows that

$$\begin{aligned} c'(g, h) &= x'_g x'_h (x'_{gh})^{-1} = (t_g x_g)(t_h x_h)(t_{gh} x_{gh})^{-1} \\ &= t_g x_g t_h (x_g^{-1} x_g) x_h x_{gh}^{-1} t_{gh}^{-1} = t_g (x_g t_h x_g^{-1}) x_g x_h x_{gh}^{-1} t_{gh}^{-1} \\ &= t_g (g * t_h) c(g, h) t_{gh}^{-1} = ((g * t_h) t_{gh}^{-1} t_g) c(g, h) \\ &= b(g, h) c(g, h), \end{aligned}$$

where  $b(g, h) = [d^1(f)](g, h)$  is a 2-coboundary obtained from the map  $f(g) = t_g$ .

# Proof (extensions of $T$ by $G_0 \iff H^2(G_0, T)$ )

## What about 2-coboundaries?

Choose representatives  $\{x'_g\}_{g \in G_0}$  instead of  $\{x_g\}_{g \in G_0}$  to obtain a 2-cocycle  $c'(g, h)$ .

Then  $x'_g = t_g x_g$  for every  $g \in G_0$  and some  $t_g \in T$ . It follows that

$$\begin{aligned} c'(g, h) &= x'_g x'_h (x'_{gh})^{-1} = (t_g x_g)(t_h x_h)(t_{gh} x_{gh})^{-1} \\ &= t_g x_g t_h (x_g^{-1} x_g) x_h x_{gh}^{-1} t_{gh}^{-1} = t_g (x_g t_h x_g^{-1}) x_g x_h x_{gh}^{-1} t_{gh}^{-1} \\ &= t_g (g * t_h) c(g, h) t_{gh}^{-1} = ((g * t_h) t_{gh}^{-1} t_g) c(g, h) \\ &= b(g, h) c(g, h), \end{aligned}$$

where  $b(g, h) = [d^1(f)](g, h)$  is a 2-coboundary obtained from the map  $f(g) = t_g$ .

Thus, the 2-cocycle obtained is unique modulo 2-coboundaries.

## Example

Consider the point group  $G_0 = D_{1,p} = \left\langle \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \right\rangle$  acting naturally on  $T = \mathbb{Z}^2$ . Then

$$H^2(G_0, T) \simeq \mathbb{Z}_2.$$

That is, there are two non-equivalent group extension of  $T$  by  $G_0$  inducing the given action.

## Proposition ( $\pm$ Snake lemma)

Let  $0 \rightarrow \mathcal{A} \rightarrow \mathcal{B} \rightarrow \mathcal{C} \rightarrow 0$  be a short exact sequence of chain complexes - that is, a commutative diagram

$$\begin{array}{ccccccccc}
 & & 0 & & 0 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \\
 \cdots & \longrightarrow & A^{n-2} & \longrightarrow & A^{n-1} & \longrightarrow & A^n & \longrightarrow & A^{n+1} & \longrightarrow & \cdots & (A) \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \\
 \cdots & \longrightarrow & B^{n-2} & \longrightarrow & B^{n-1} & \longrightarrow & B^n & \longrightarrow & B^{n+1} & \longrightarrow & \cdots & (B) \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \\
 \cdots & \longrightarrow & C^{n-2} & \longrightarrow & C^{n-1} & \longrightarrow & C^n & \longrightarrow & C^{n+1} & \longrightarrow & \cdots & (C) \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \\
 & & 0 & & 0 & & 0 & & 0 & & 
 \end{array}$$

such that the rows are complexes  $\mathcal{A}, \mathcal{B}, \mathcal{C}$  and the columns are exact. Then there is an exact sequence of the following form:

$$\cdots \longrightarrow \mathbb{H}^{n-1}(\mathcal{C}) \longrightarrow \mathbb{H}^n(\mathcal{A}) \longrightarrow \mathbb{H}^n(\mathcal{B}) \longrightarrow \mathbb{H}^n(\mathcal{C}) \longrightarrow \mathbb{H}^{n+1}(\mathcal{A}) \longrightarrow \cdots$$

**Fact:** There is a short exact sequence of chain complexes

$$\begin{array}{ccccccc}
 & 0 & & 0 & & 0 & & 0 & & 0 & & 0 \\
 & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & \mathbb{Z} & \longrightarrow & \mathbb{Z}^2 & \xrightarrow{\begin{pmatrix} 0 & 0 \\ 0 & -2 \end{pmatrix}} & \mathbb{Z}^2 & \xrightarrow{\begin{pmatrix} 2 & 0 \\ 0 & 0 \end{pmatrix}} & \mathbb{Z}^2 & \xrightarrow{\begin{pmatrix} 0 & 0 \\ 0 & -2 \end{pmatrix}} & \mathbb{Z}^2 & \xrightarrow{\begin{pmatrix} 2 & 0 \\ 0 & 0 \end{pmatrix}} & \dots
 \end{array} \quad (\mathcal{A})$$

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \mathbb{Z} & \longrightarrow & C^0(G_0, T) & \longrightarrow & C^1(G_0, T) & \longrightarrow & C^2(G_0, T) & \longrightarrow & C^3(G_0, T) & \longrightarrow & \dots
 \end{array} \quad (\mathcal{B})$$

$$\begin{array}{ccccccc}
 0 & \longrightarrow & 0 & \longrightarrow & C^0 & \longrightarrow & C^1 & \longrightarrow & C^2 & \longrightarrow & C^3 & \longrightarrow & \dots \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \\
 & & 0 & & 0 & & 0 & & 0 & & 0 & & 
 \end{array} \quad (\mathcal{C})$$

where the sequence  $(\mathcal{C})$  is also exact.

Thus, we obtain a long exact sequence

$$\begin{aligned} 0 &\longrightarrow \mathbb{H}^1(\mathcal{A}) \longrightarrow H^1(G_0, T) \longrightarrow 0 \longrightarrow \mathbb{H}^2(\mathcal{A}) \longrightarrow H^2(G_0, T) \longrightarrow 0 \\ \mathbb{H}^0(\mathcal{C}) &\longrightarrow \mathbb{H}^1(\mathcal{A}) \longrightarrow \mathbb{H}^1(\mathcal{B}) \longrightarrow \mathbb{H}^1(\mathcal{C}) \longrightarrow \mathbb{H}^2(\mathcal{A}) \longrightarrow \mathbb{H}^2(\mathcal{C}) \longrightarrow \mathbb{H}^3(\mathcal{C}) \end{aligned}$$

Thus, we obtain a long exact sequence

$$\begin{aligned} 0 \longrightarrow \mathbb{H}^1(\mathcal{A}) \longrightarrow H^1(G_0, T) \longrightarrow 0 \longrightarrow \mathbb{H}^2(\mathcal{A}) \longrightarrow H^2(G_0, T) \longrightarrow 0 \\ \mathbb{H}^0(\mathcal{C}) \longrightarrow \mathbb{H}^1(\mathcal{A}) \longrightarrow \mathbb{H}^1(\mathcal{B}) \longrightarrow \mathbb{H}^1(\mathcal{C}) \longrightarrow \mathbb{H}^2(\mathcal{A}) \longrightarrow \mathbb{H}^2(\mathcal{C}) \longrightarrow \mathbb{H}^3(\mathcal{C}) \end{aligned}$$

In particular,  $H^2(G_0, T) \simeq \mathbb{H}^2(\mathcal{A})$ .



Thus, we obtain a long exact sequence

$$0 \longrightarrow \mathbb{H}^1(\mathcal{A}) \longrightarrow H^1(G_0, T) \longrightarrow 0 \longrightarrow \mathbb{H}^2(\mathcal{A}) \longrightarrow H^2(G_0, T) \longrightarrow 0$$

$$\mathbb{H}^0(\mathcal{C}) \longrightarrow \mathbb{H}^1(\mathcal{A}) \longrightarrow \mathbb{H}^1(\mathcal{B}) \longrightarrow \mathbb{H}^1(\mathcal{C}) \longrightarrow \mathbb{H}^2(\mathcal{A}) \longrightarrow \mathbb{H}^2(\mathcal{C}) \longrightarrow \mathbb{H}^3(\mathcal{C})$$

In particular,  $H^2(G_0, T) \simeq \mathbb{H}^2(\mathcal{A})$ .

Thus,

$$H^2(G_0, T) \simeq \left\{ \begin{pmatrix} 0 \\ z \end{pmatrix} \mid z \in \mathbb{Z} \right\} / \left\{ \begin{pmatrix} 0 \\ -2z \end{pmatrix} \mid z \in \mathbb{Z} \right\} \simeq \mathbb{Z}_2.$$

We conclude this part by listing the second cohomology groups of all possible choices of the point group and the lattice:

$G_0$	$H^2(G_0, T)$	No. of extensions
$C_1$	0	1
$C_2$	0	1
$C_3$	0	1
$C_4$	0	1
$C_6$	0	1
$D_{1,p}$	$\mathbb{Z}_2$	2
$D_{1,c}$	0	1
$D_{2,p}$	$\mathbb{Z}_2 \times \mathbb{Z}_2$	4
$D_{2,c}$	0	1
$D_{3,l}$	0	1
$D_{3,s}$	0	1
$D_4$	$\mathbb{Z}_2$	2
$D_6$	0	1

Thank you for your attention!