

Persistent Homology II

Michal Hrbek

Charles University - Faculty of Mathematics and Physics

November 24, 2016

Outline: (note the rapidly decreasing amount of rigour)

- 1 Persistent modules and Carlsson's decomposition theorem (some real mathematics),
- 2 Methodology of using persistent homology to discover topological features of point clouds (a legit method of data mining),
- 3 The experiment of Carlsson et al. - topological features of the space of 3×3 patches of natural images (some proper shamanism).

Let k be a field and $(P, <)$ a totally ordered set with a dense countable subset (in our applications, typically either \mathbb{N} or \mathbb{R} with the usual order).

Persistent modules

Let k be a field and $(P, <)$ a totally ordered set with a dense countable subset (in our applications, typically either \mathbb{N} or \mathbb{R} with the usual order).

A $(P-)$ persistent module M is a functor from P to the category of k -vector spaces. (Here, poset P is viewed naturally as a category.)

Let k be a field and $(P, <)$ a totally ordered set with a dense countable subset (in our applications, typically either \mathbb{N} or \mathbb{R} with the usual order).

A $(P-)$ persistent module M is a functor from P to the category of k -vector spaces. (Here, poset P is viewed naturally as a category.)

Explicitly: Data for M consist of vector spaces $V_p, p \in P$ and k -linear maps $f_{pq}^M : V_p \rightarrow V_q$ for all $p \leq q$, together with conditions:

- $f_{pp}^M = \text{id}_{V_p}$ for all $p \in P$,
- $f_{qr}^M \circ f_{pq} = f_{pq}$ for all $p \leq q \leq r \in P$.

A subset I of $(P, <)$ is an interval if $p \leq r \leq q \in P$, where $p, q \in I$, implies that $r \in I$.

A subset I of $(P, <)$ is an interval if $p \leq r \leq q \in P$, where $p, q \in I$, implies that $r \in I$.

We can assign a P -persistent module M_I to I in the following way:

- $M_I(p) = \begin{cases} k, & p \in I \\ 0, & p \notin I \end{cases}$,
- $M_I(p < q) = \begin{cases} \text{Id}_k, & p, q \in I \\ 0, & \text{otherwise } \notin I \end{cases}$

A P -persistent module M is pointwise finite, if $M(p)$ is finite dimensional for each $p \in P$.

A P -persistent module M is pointwise finite, if $M(p)$ is finite dimensional for each $p \in P$.

Theorem (Carlsson, Crawley-Boevey)

Under certain mild conditions for finiteness of M (satisfied e.g. when M is pointwise finite), any P -persistent module M is isomorphic direct sum of interval modules.

A P -persistent module M is pointwise finite, if $M(p)$ is finite dimensional for each $p \in P$.

Theorem (Carlsson, Crawley-Boevey)

Under certain mild conditions for finiteness of M (satisfied e.g. when M is pointwise finite), any P -persistent module M is isomorphic direct sum of interval modules.

What does isomorphic mean?

Let M, N be P -persistent modules. A homomorphism $f : M \rightarrow N$ of P -persistent modules is a collection of k -linear maps $\varphi_p : M(p) \rightarrow N(p)$ such that the following diagram commutes for any $p < q \in P$:

$$\begin{array}{ccc} M(p) & \xrightarrow{f_{pq}^M} & M(q) \\ \varphi_p \downarrow & & \downarrow \varphi_q \\ N(p) & \xrightarrow{f_{pq}^N} & N(q) \end{array}$$

Let M, N be P -persistent modules. A homomorphism $f : M \rightarrow N$ of P -persistent modules is a collection of k -linear maps $\varphi_p : M(p) \rightarrow N(p)$ such that the following diagram commutes for any $p < q \in P$:

$$\begin{array}{ccc} M(p) & \xrightarrow{f_{pq}^M} & M(q) \\ \varphi_p \downarrow & & \downarrow \varphi_q \\ N(p) & \xrightarrow{f_{pq}^N} & N(q) \end{array}$$

In fancy words, $\varphi : M \rightarrow N$ is a natural transformation of functors N and M .

Let M, N be P -persistent modules. A homomorphism $f : M \rightarrow N$ of P -persistent modules is a collection of k -linear maps $\varphi_p : M(p) \rightarrow N(p)$ such that the following diagram commutes for any $p < q \in P$:

$$\begin{array}{ccc} M(p) & \xrightarrow{f_{pq}^M} & M(q) \\ \varphi_p \downarrow & & \downarrow \varphi_q \\ N(p) & \xrightarrow{f_{pq}^N} & N(q) \end{array}$$

In fancy words, $\varphi : M \rightarrow N$ is a natural transformation of functors N and M .

We say that φ is an isomorphism of persistent modules if $\varphi(p)$ is an isomorphism for each $p \in P$.

A P -persistent module M is pointwise finite, if $M(p)$ is finite dimensional for each $p \in P$.

Theorem (Carlsson, Crawley-Boevey)

Under certain mild conditions for finiteness of M (satisfied e.g. when M is pointwise finite), any P -persistent module M is isomorphic direct sum of interval modules.

What is a direct sum?

Let $M_i, i \in I$ be a collection of P -persistent modules. Then we define the direct sum $\bigoplus_{i \in I} M_i$ to be the P -persistent module defined as follows:

- $\bigoplus_{i \in I} M_i(p) = \bigoplus_{i \in I} M_i(p),$
- $\bigoplus_{i \in I} M_i(p < q) = \bigoplus_{i \in I} M_i(p < q).$

Let X be a point cloud - a finite subset of \mathbb{R}^d .

Let X be a point cloud - a finite subset of \mathbb{R}^d .

- Choose a method of turning X into a simplicial complex - Čech complex, Vietoris-Rips, witness...

Let X be a point cloud - a finite subset of \mathbb{R}^d .

- Choose a method of turning X into a simplicial complex - Čech complex, Vietoris-Rips, witness...
- Compute the simplicial complex $C_\epsilon(X)$ for each parameter $\epsilon > 0$ (well, not really of course)


Let X be a point cloud - a finite subset of \mathbb{R}^d .

- Choose a method of turning X into a simplicial complex - Čech complex, Vietoris-Rips, witness...
- Compute the simplicial complex $C_\epsilon(X)$ for each parameter $\epsilon > 0$ (well, not really of course)
- X is finite, so there is only finitely many steps, in which the complex changes.



Let X be a point cloud - a finite subset of \mathbb{R}^d .

- Choose a method of turning X into a simplicial complex - Čech complex, Vietoris-Rips, witness...
- Compute the simplicial complex $C_\epsilon(X)$ for each parameter $\epsilon > 0$ (well, not really of course)
- X is finite, so there is only finitely many steps, in which the complex changes.
- Let $0 = \epsilon_0 < \epsilon_1 < \dots < \epsilon_n < \epsilon_{n+1} = \infty$ be parameters such that $C_{\epsilon_i}(X) = C_\delta(X)$ for any $\epsilon_i < \delta < \epsilon_{i+1}$ and $i < n + 1$.


$$\epsilon_i < \epsilon_j$$



$$\epsilon_i < \epsilon_j$$



$$C_{\epsilon_i}(X) \hookrightarrow C_{\epsilon_j}(X)$$



- Rename $C_i = C_{\epsilon_i}(X)$.

- Rename $C_i = C_{\epsilon_i}(X)$.

- Rename $C_i = C_{\epsilon_i}(X)$.
- We obtain an increasing sequence of complexes

$$C_0 \hookrightarrow C_1 \hookrightarrow \cdots \hookrightarrow C_{n+1}.$$

- Rename $C_i = C_{\epsilon_i}(X)$.
- We obtain an increasing sequence of complexes

$$C_0 \hookrightarrow C_1 \hookrightarrow \cdots \hookrightarrow C_{n+1}.$$

- By functoriality, this filtration yields maps on k -th homologies for each $k \geq 0$:

$$H_k(C_0) \rightarrow H_k(C_1) \rightarrow \cdots \rightarrow H_k(C_{n+1}).$$

- Rename $C_i = C_{\epsilon_i}(X)$.
- We obtain an increasing sequence of complexes

$$C_0 \hookrightarrow C_1 \hookrightarrow \cdots \hookrightarrow C_{n+1}.$$

- By functoriality, this filtration yields maps on k -th homologies for each $k \geq 0$:

$$H_k(C_0) \rightarrow H_k(C_1) \rightarrow \cdots \rightarrow H_k(C_{n+1}).$$

- Thus, we get for each $k \geq 0$ an \mathbb{N} -persistent module H_k .

- By Carlsson's Theorem, H_k decomposes into a direct sum of interval modules $\bigoplus_{i=1}^l M(I_i)$.


- By Carlsson's Theorem, H_k decomposes into a direct sum of interval modules $\bigoplus_{i=1}^l M(I_i)$.
- The intervals I_i , a priori intervals of \mathbb{N} , are interpreted as intervals of \mathbb{R} with borders from $\{\epsilon_0, \epsilon_1, \dots, \epsilon_{n+1}\}$.

- By Carlsson's Theorem, H_k decomposes into a direct sum of interval modules $\bigoplus_{i=1}^l M(I_i)$.
- The intervals I_i , a priori intervals of \mathbb{N} , are interpreted as intervals of \mathbb{R} with borders from $\{\epsilon_0, \epsilon_1, \dots, \epsilon_{n+1}\}$.
- Each interval $I_i = (\epsilon_i, \epsilon_j)$ corresponds to a “birth” of a non-trivial homology cycle (think of a “hole”) at parameter ϵ_i , and its “death” at parameter ϵ_j .

Complete topological information obtained by this method in each homological dimension is given by a multiset of intervals. We call this multiset a barcode.

Barcodes

Complete topological information obtained by this method in each homological dimension is given by a multiset of intervals. We call this multiset a barcode.



3x3 patches of natural images

In what follows, “they”=Carlsson et al.:

3x3 patches of natural images

In what follows, “they”=Carlsson et al.:

- Input data: Large collection of grayscale natural images.

3x3 patches of natural images

In what follows, “they”=Carlsson et al.:

- Input data: Large collection of grayscale natural images.
- From each image, they choose 5000 random 3x3-pixel patches. This gives a point cloud in \mathbb{R}^9 .

3x3 patches of natural images

In what follows, “they”=Carlsson et al.:

- Input data: Large collection of grayscale natural images.
- From each image, they choose 5000 random 3x3-pixel patches. This gives a point cloud in \mathbb{R}^9 .
- Compute a “D-norm” of each patch measuring its contrast (certain quadratic form of logs of intensities of pixels).

3x3 patches of natural images

In what follows, “they”=Carlsson et al.:

- Input data: Large collection of grayscale natural images.
- From each image, they choose 5000 random 3x3-pixel patches. This gives a point cloud in \mathbb{R}^9 .
- Compute a “D-norm” of each patch measuring its contrast (certain quadratic form of logs of intensities of pixels).
- Keep for each image only those patches being in the top 20% with respect to D-norm.

- Mean center the data - subtract a mean of intensity from each pixel. That is, two patches are now considered identical if their “brightness” differ by a constant. This puts all the patches into a 8-dimensional subspace of \mathbb{R}^9 .

- Mean center the data - subtract a mean of intensity from each pixel. That is, two patches are now considered identical if their “brightness” differ by a constant. This puts all the patches into a 8-dimensional subspace of \mathbb{R}^9 .
- Normalize with respect to the D-norm (normalize the contrast). That is, two patches are now considered identical if they only differ by “turning up contrast”. This puts the data on a 7-dimensional ellipsoid inside \mathbb{R}^8 .

- Mean center the data - subtract a mean of intensity from each pixel. That is, two patches are now considered identical if their “brightness” differ by a constant. This puts all the patches into a 8-dimensional subspace of \mathbb{R}^9 .
- Normalize with respect to the D-norm (normalize the contrast). That is, two patches are now considered identical if they only differ by “turning up contrast”. This puts the data on a 7-dimensional ellipsoid inside \mathbb{R}^8 .
- Do a coordinate change in order to make this ellipsoid into a 7-sphere S^7 . Our data now is a point cloud \mathcal{M} of approx. 4 millions points on S^7 .

How does \mathcal{M} sit in S^7 ?

- Further heuristic: $\delta_k(x)$ is the distance of x from the k -th closest point of \mathcal{M} to x .

How does \mathcal{M} sit in S^7 ?

- Further heuristic: $\delta_k(x)$ is the distance of x from the k -th closest point of \mathcal{M} to x .
- $\mathcal{M}[k, T] = \{x \in \mathcal{M} \mid \delta_k(x) \text{ is among the } T\% \text{ lowest values of } \delta_k \text{ on } \mathcal{M}\}.$


How does \mathcal{M} sit in S^7 ?

- Further heuristic: $\delta_k(x)$ is the distance of x from the k -th closest point of \mathcal{M} to x .
- $\mathcal{M}[k, T] = \{x \in \mathcal{M} \mid \delta_k(x) \text{ is among the } T\% \text{ lowest values of } \delta_k \text{ on } \mathcal{M}\}$.
- For computational reasons, take samples of size approx. 10^4 from \mathcal{M} , and construct simplicial complexes using the witness method.

How does \mathcal{M} sit in S^7 ?

- Further heuristic: $\delta_k(x)$ is the distance of x from the k -th closest point of \mathcal{M} to x .
- $\mathcal{M}[k, T] = \{x \in \mathcal{M} \mid \delta_k(x) \text{ is among the } T\% \text{ lowest values of } \delta_k \text{ on } \mathcal{M}\}$.
- For computational reasons, take samples of size approx. 10^4 from \mathcal{M} , and construct simplicial complexes using the witness method.
- Compute the persistent homology barcodes with coefficients in \mathbb{Z}_2 using the PLEX package.

Barcode in homological dimension 0 of $\mathcal{M}[15, 30]$:



Interpretation: \mathcal{M} is connected (has one component).

Barcodes

Barcode in homological dimension 1 of $\mathcal{M}[300, 30]$ (very crude approximation):



Interpretation: \mathcal{M} in this detail has “1 hole” - simplest explanation is, that it looks like a circle.


Barcodes

Barcode in homological dimension 1 of $\mathcal{M}[300, 30]$ (very crude approximation):



Interpretation: \mathcal{M} in this detail has “1 hole” - simplest explanation is, that it looks like a circle.

The picture below gives such an explanation.



Barcode in homological dimension 1 of $\mathcal{M}[15, 30]$:



Barcode in homological dimension 1 of $\mathcal{M}[15, 30]$:




Interpretation: \mathcal{M} actually has more 1-holes - the first Betti number seems to 5.

Barcode in homological dimension 1 of $\mathcal{M}[15, 30]$:



Interpretation: \mathcal{M} actually has more 1-holes - the first Betti number seems to 5.


Carlsson et al. propose the following “3-circle” model:



The green and the red circle both intersect the black one in two points. The red and the green circle are disjoint.

$$\begin{aligned}\text{First Betti number} &= \# \text{arcs} - \# \text{vertices} + \# \text{components} \\ &= 8 - 4 + 1 = 5\end{aligned}$$


Again, a sort of explanation:



Primary circle corresponds to “linear patches”. Two secondary circles correspond to “horizontally/vertically aligned patches”. Take polynomial in 2 variables, and evaluate on set $\{-1, 0, 1\} \times \{-1, 0, 1\}$. Linear, quadratic patches...

Klein bottle

Klein bottle is a 2-surface defined as a topological quotient of a square:



Klein bottles does not embed into \mathbb{R}^3 , but it does embed into \mathbb{R}^4 .

The 3-circle model embeds into the Klein bottle:




- What now? Meddle with the parameters until we find the Klein bottle.

- What now? Meddle with the parameters until we find the Klein bottle.
- The Betti numbers with coefficients in \mathbb{Z}_2 of the Klein bottle are $\beta_0 = 1, \beta_1 = 2, \beta_2 = 1, \beta_i = 0$ for $i > 2$.

- What now? Meddle with the parameters until we find the Klein bottle.
- The Betti numbers with coefficients in \mathbb{Z}_2 of the Klein bottle are $\beta_0 = 1, \beta_1 = 2, \beta_2 = 1, \beta_i = 0$ for $i > 2$.
- While changing parameter k , we unfortunately always get zero second Betti number.

- What now? Meddle with the parameters until we find the Klein bottle.
- The Betti numbers with coefficients in \mathbb{Z}_2 of the Klein bottle are $\beta_0 = 1, \beta_1 = 2, \beta_2 = 1, \beta_i = 0$ for $i > 2$.
- While changing parameter k , we unfortunately always get zero second Betti number.
- More heuristic to the rescue... Authors argue that their method has preference for linear and vertically/horizontally aligned patches. They add a certain set of points \mathcal{Q} from \mathcal{M} not in $\mathcal{M}[100, 30]$ of pure quadratic character.

Barcodes for $\mathcal{M}[100, 30] \cup \mathcal{Q}$:



Interpretation: Point cloud $\mathcal{M}[100, 30] \cup \mathcal{Q}$ “looks like” something with Betti numbers $\beta_0 = 1, \beta_1 = 2, \beta_2 = 1$.

More handwaving: There are two 2-surfaces with Betti numbers mod 2 being $\beta_0 = 1, \beta_1 = 2, \beta_2 = 1$ - Klein bottle and the torus. Their homology groups differ mod 3 though, and the authors claim to have done the corresponding persistent homology computation, and that it is in the favor of the Klein bottle...

