

# Growth rates of solvable algebras

Algebras & Clones, July 2014, Prague

Keith. A Kearnes, Emil W. Kiss, Ágnes Szendrei

[kearnes@euclid.colorado.edu](mailto:kearnes@euclid.colorado.edu)

[ewkiss@cs.elte.hu](mailto:ewkiss@cs.elte.hu)

[szendrei@euclid.colorado.edu](mailto:szendrei@euclid.colorado.edu)

University of Colorado Boulder, USA

Eötvös University, Budapest, Hungary

University of Colorado Boulder, USA

# Growth rate

The growth rate of a finite algebra  $\mathbf{A}$  is the function  $d_{\mathbf{A}}(n) =$  the least size of a generating set for  $\mathbf{A}^n$ .

# Growth rate

The growth rate of a finite algebra  $\mathbf{A}$  is the function  $d_{\mathbf{A}}(n) =$  the least size of a generating set for  $\mathbf{A}^n$ .

## Examples

$\mathbf{A}$  is a module over a ring. Then  $d_{\mathbf{A}}(n) = \Theta(n)$  (linear).

# Growth rate

The growth rate of a finite algebra  $\mathbf{A}$  is the function  $d_{\mathbf{A}}(n) =$  the least size of a generating set for  $\mathbf{A}^n$ .

## Examples

$\mathbf{A}$  is a module over a ring. Then  $d_{\mathbf{A}}(n) = \Theta(n)$  (linear).

**Reason:** The size of a basis in a vector space  $\mathbf{F}^n$  is  $n$ .

# Growth rate

The growth rate of a finite algebra  $\mathbf{A}$  is the function  $d_{\mathbf{A}}(n) =$  the least size of a generating set for  $\mathbf{A}^n$ .

## Examples

$\mathbf{A}$  is a module over a ring. Then  $d_{\mathbf{A}}(n) = \Theta(n)$  (linear).

**Reason:** The size of a basis in a vector space  $\mathbf{F}^n$  is  $n$ .

$\mathbf{A}$  is a Boolean algebra. Then  $d_{\mathbf{A}}(n) = \Theta(\log(n))$  (logarithmic).

# Growth rate

The growth rate of a finite algebra  $\mathbf{A}$  is the function  $d_{\mathbf{A}}(n) =$  the least size of a generating set for  $\mathbf{A}^n$ .

## Examples

$\mathbf{A}$  is a module over a ring. Then  $d_{\mathbf{A}}(n) = \Theta(n)$  (linear).

**Reason:** The size of a basis in a vector space  $\mathbf{F}^n$  is  $n$ .

$\mathbf{A}$  is a Boolean algebra. Then  $d_{\mathbf{A}}(n) = \Theta(\log(n))$  (logarithmic).  
The same holds if  $\mathbf{A}$  is a simple nonabelian group.

# Growth rate

The growth rate of a finite algebra  $\mathbf{A}$  is the function  $d_{\mathbf{A}}(n) =$  the least size of a generating set for  $\mathbf{A}^n$ .

## Examples

$\mathbf{A}$  is a module over a ring. Then  $d_{\mathbf{A}}(n) = \Theta(n)$  (linear).

**Reason:** The size of a basis in a vector space  $\mathbf{F}^n$  is  $n$ .

$\mathbf{A}$  is a Boolean algebra. Then  $d_{\mathbf{A}}(n) = \Theta(\log(n))$  (logarithmic).  
The same holds if  $\mathbf{A}$  is a simple nonabelian group.

**Reason:** all finitary functions on  $\mathbf{A}$  are polynomials.

# Growth rate

The growth rate of a finite algebra  $\mathbf{A}$  is the function  $d_{\mathbf{A}}(n) =$  the least size of a generating set for  $\mathbf{A}^n$ .

## Examples

$\mathbf{A}$  is a module over a ring. Then  $d_{\mathbf{A}}(n) = \Theta(n)$  (linear).

**Reason:** The size of a basis in a vector space  $\mathbf{F}^n$  is  $n$ .

$\mathbf{A}$  is a Boolean algebra. Then  $d_{\mathbf{A}}(n) = \Theta(\log(n))$  (logarithmic).  
The same holds if  $\mathbf{A}$  is a simple nonabelian group.

**Reason:** all finitary functions on  $\mathbf{A}$  are polynomials.

$\mathbf{A}$  is a unary algebra. Then  $d_{\mathbf{A}}(n) = 2^{\Theta(n)}$  (exponential).



# Growth rate

The growth rate of a finite algebra  $\mathbf{A}$  is the function  $d_{\mathbf{A}}(n) =$  the least size of a generating set for  $\mathbf{A}^n$ .

## Examples

$\mathbf{A}$  is a module over a ring. Then  $d_{\mathbf{A}}(n) = \Theta(n)$  (linear).

**Reason:** The size of a basis in a vector space  $\mathbf{F}^n$  is  $n$ .

$\mathbf{A}$  is a Boolean algebra. Then  $d_{\mathbf{A}}(n) = \Theta(\log(n))$  (logarithmic).  
The same holds if  $\mathbf{A}$  is a simple nonabelian group.

**Reason:** all finitary functions on  $\mathbf{A}$  are polynomials.

$\mathbf{A}$  is a unary algebra. Then  $d_{\mathbf{A}}(n) = 2^{\Theta(n)}$  (exponential).

**Reason:** The free algebras over  $\mathbf{A}$  have polynomially bounded size.

# Wiegold dichotomy

## Theorem (J. Wiegold, 1974)

$\mathbf{G}$  is a finite group. If  $\mathbf{G}$  has a nontrivial abelian factor group, then  $d_{\mathbf{G}}$  is linear.

# Wiegold dichotomy

## Theorem (J. Wiegold, 1974)

$\mathbf{G}$  is a finite group. If  $\mathbf{G}$  has a nontrivial abelian factor group, then  $d_{\mathbf{G}}$  is linear. Otherwise (that is, if  $\mathbf{G}$  is perfect)  $d_{\mathbf{G}}$  is logarithmic.

# Wiegold dichotomy

## Theorem (J. Wiegold, 1974)

$\mathbf{G}$  is a finite group. If  $\mathbf{G}$  has a nontrivial abelian factor group, then  $d_{\mathbf{G}}$  is linear. Otherwise (that is, if  $\mathbf{G}$  is perfect)  $d_{\mathbf{G}}$  is logarithmic.

## Remarks

- If  $\mathbf{B}$  is a homomorphic image of  $\mathbf{A}$ , then  $d_{\mathbf{B}}(n) \leq d_{\mathbf{A}}(n)$ .

# Wiegold dichotomy

## Theorem (J. Wiegold, 1974)

$\mathbf{G}$  is a finite group. If  $\mathbf{G}$  has a nontrivial abelian factor group, then  $d_{\mathbf{G}}$  is linear. Otherwise (that is, if  $\mathbf{G}$  is perfect)  $d_{\mathbf{G}}$  is logarithmic.

## Remarks

- If  $\mathbf{B}$  is a homomorphic image of  $\mathbf{A}$ , then  $d_{\mathbf{B}}(n) \leq d_{\mathbf{A}}(n)$ .  
So if  $\mathbf{G}$  has an abelian factor, then  $d_{\mathbf{G}}$  is at least linear.

# Wiegold dichotomy

## Theorem (J. Wiegold, 1974)

$\mathbf{G}$  is a finite group. If  $\mathbf{G}$  has a nontrivial abelian factor group, then  $d_{\mathbf{G}}$  is linear. Otherwise (that is, if  $\mathbf{G}$  is perfect)  $d_{\mathbf{G}}$  is logarithmic.

## Remarks

- If  $\mathbf{B}$  is a homomorphic image of  $\mathbf{A}$ , then  $d_{\mathbf{B}}(n) \leq d_{\mathbf{A}}(n)$ .  
So if  $\mathbf{G}$  has an abelian factor, then  $d_{\mathbf{G}}$  is at least linear.
- If  $\mathbf{B}$  is an expansion of  $\mathbf{A}$ , then  $d_{\mathbf{B}}(n) \leq d_{\mathbf{A}}(n)$ .

# Wiegold dichotomy

## Theorem (J. Wiegold, 1974)

$\mathbf{G}$  is a finite group. If  $\mathbf{G}$  has a nontrivial abelian factor group, then  $d_{\mathbf{G}}$  is linear. Otherwise (that is, if  $\mathbf{G}$  is perfect)  $d_{\mathbf{G}}$  is logarithmic.

## Remarks

- If  $\mathbf{B}$  is a homomorphic image of  $\mathbf{A}$ , then  $d_{\mathbf{B}}(n) \leq d_{\mathbf{A}}(n)$ .  
So if  $\mathbf{G}$  has an abelian factor, then  $d_{\mathbf{G}}$  is at least linear.
- If  $\mathbf{B}$  is an expansion of  $\mathbf{A}$ , then  $d_{\mathbf{B}}(n) \leq d_{\mathbf{A}}(n)$ .  
The richer the structure, the smaller the growth rate.

# Wiegold dichotomy

## Theorem (J. Wiegold, 1974)

$\mathbf{G}$  is a finite group. If  $\mathbf{G}$  has a nontrivial abelian factor group, then  $d_{\mathbf{G}}$  is linear. Otherwise (that is, if  $\mathbf{G}$  is perfect)  $d_{\mathbf{G}}$  is logarithmic.

## Remarks

- If  $\mathbf{B}$  is a homomorphic image of  $\mathbf{A}$ , then  $d_{\mathbf{B}}(n) \leq d_{\mathbf{A}}(n)$ .  
So if  $\mathbf{G}$  has an abelian factor, then  $d_{\mathbf{G}}$  is at least linear.
- If  $\mathbf{B}$  is an expansion of  $\mathbf{A}$ , then  $d_{\mathbf{B}}(n) \leq d_{\mathbf{A}}(n)$ .  
The richer the structure, the smaller the growth rate.

Wiegold-dichotomy holds for **Maltsev algebras** (see later).



# Wiegold dichotomy

## Theorem (J. Wiegold, 1974)

$\mathbf{G}$  is a finite group. If  $\mathbf{G}$  has a nontrivial abelian factor group, then  $d_{\mathbf{G}}$  is linear. Otherwise (that is, if  $\mathbf{G}$  is perfect)  $d_{\mathbf{G}}$  is logarithmic.

## Remarks

- If  $\mathbf{B}$  is a homomorphic image of  $\mathbf{A}$ , then  $d_{\mathbf{B}}(n) \leq d_{\mathbf{A}}(n)$ .  
So if  $\mathbf{G}$  has an abelian factor, then  $d_{\mathbf{G}}$  is at least linear.
- If  $\mathbf{B}$  is an expansion of  $\mathbf{A}$ , then  $d_{\mathbf{B}}(n) \leq d_{\mathbf{A}}(n)$ .  
The richer the structure, the smaller the growth rate.

Wiegold-dichotomy holds for **Maltsev algebras** (see later).

## Motivating problem

What are the possible growth rates of finite algebras?

# Pointed cube terms

**Example:** Maltsev-term ( $x_1 x_2^{-1} x_3$  in groups).

# Pointed cube terms

**Example:** Maltsev-term  $(x_1 x_2^{-1} x_3$  in groups).

$$m \begin{pmatrix} x & y & y \\ y & y & x \end{pmatrix} \approx \begin{pmatrix} x \\ x \end{pmatrix},$$

# Pointed cube terms

**Example:** Maltsev-term  $(x_1 x_2^{-1} x_3$  in groups).

$$m \begin{pmatrix} x & y & y \\ y & y & x \end{pmatrix} \approx \begin{pmatrix} x \\ x \end{pmatrix},$$

$$\Sigma \models F(M) \approx \begin{pmatrix} x \\ \vdots \\ x \end{pmatrix}.$$

# Pointed cube terms

**Example:** Maltsev-term  $(x_1 x_2^{-1} x_3$  in groups).

$$m \begin{pmatrix} x & y & y \\ y & y & x \end{pmatrix} \approx \begin{pmatrix} x \\ x \end{pmatrix},$$

there is a  $k \times m$  matrix  $M$  consisting of variables and  $p$  distinct constant symbols,

$$\Sigma \models F(M) \approx \begin{pmatrix} x \\ \vdots \\ x \end{pmatrix}.$$

# Pointed cube terms

**Example:** Maltsev-term  $(x_1 x_2^{-1} x_3$  in groups).

$$m \begin{pmatrix} x & y & y \\ y & y & x \end{pmatrix} \approx \begin{pmatrix} x \\ x \end{pmatrix},$$

there is a  $k \times m$  matrix  $M$  consisting of variables and  $p$  distinct constant symbols, with every column of  $M$  containing a symbol different from  $x$ , such that

$$\Sigma \models F(M) \approx \begin{pmatrix} x \\ \vdots \\ x \end{pmatrix}.$$

# Pointed cube terms

**Example:** Maltsev-term  $(x_1 x_2^{-1} x_3$  in groups).

$$m \begin{pmatrix} x & y & y \\ y & y & x \end{pmatrix} \approx \begin{pmatrix} x \\ x \end{pmatrix},$$

$p$ -pointed,

there is a  $k \times m$  matrix  $M$  consisting of variables and  $p$  distinct constant symbols, with every column of  $M$  containing a symbol different from  $x$ , such that

$$\Sigma \models F(M) \approx \begin{pmatrix} x \\ \vdots \\ x \end{pmatrix}.$$

# Pointed cube terms

**Example:** Maltsev-term  $(x_1 x_2^{-1} x_3$  in groups).

$$m \begin{pmatrix} x & y & y \\ y & y & x \end{pmatrix} \approx \begin{pmatrix} x \\ x \end{pmatrix},$$

**$p$ -pointed,  $k$ -cube term**

there is a  $k \times m$  matrix  $M$  consisting of variables and  $p$  distinct constant symbols, with every column of  $M$  containing a symbol different from  $x$ , such that

$$\Sigma \models F(M) \approx \begin{pmatrix} x \\ \vdots \\ x \end{pmatrix}.$$



# Pointed cube terms

**Example:** Maltsev-term ( $x_1 x_2^{-1} x_3$  in groups).

$$m \begin{pmatrix} x & y & y \\ y & y & x \end{pmatrix} \approx \begin{pmatrix} x \\ x \end{pmatrix},$$

If  $\Sigma$  is a set of identities in a language  $\mathcal{L}$ , then an  $\mathcal{L}$ -term  $F(x_1, \dots, x_m)$  is a  $p$ -pointed,  $k$ -cube term for the variety axiomatized by  $\Sigma$  if there is a  $k \times m$  matrix  $M$  consisting of variables and  $p$  distinct constant symbols, with every column of  $M$  containing a symbol different from  $x$ , such that

$$\Sigma \models F(M) \approx \begin{pmatrix} x \\ \vdots \\ x \end{pmatrix}.$$

# Pointed cube terms

**Example:** Maltsev-term  $(x_1 x_2^{-1} x_3$  in groups).

$$m \begin{pmatrix} x & y & y \\ y & y & x \end{pmatrix} \approx \begin{pmatrix} x \\ x \end{pmatrix},$$

witnesses that  $m(x_1, x_2, x_3)$  is a 3-ary,

If  $\Sigma$  is a set of identities in a language  $\mathcal{L}$ , then an  $\mathcal{L}$ -term  $F(x_1, \dots, x_m)$  is a  $p$ -pointed,  $k$ -cube term for the variety axiomatized by  $\Sigma$  if there is a  $k \times m$  matrix  $M$  consisting of variables and  $p$  distinct constant symbols, with every column of  $M$  containing a symbol different from  $x$ , such that

$$\Sigma \models F(M) \approx \begin{pmatrix} x \\ \vdots \\ x \end{pmatrix}.$$

# Pointed cube terms

**Example:** Maltsev-term  $(x_1 x_2^{-1} x_3$  in groups).

$$m \begin{pmatrix} x & y & y \\ y & y & x \end{pmatrix} \approx \begin{pmatrix} x \\ x \end{pmatrix},$$

witnesses that  $m(x_1, x_2, x_3)$  is a 3-ary, 0-pointed,

If  $\Sigma$  is a set of identities in a language  $\mathcal{L}$ , then an  $\mathcal{L}$ -term  $F(x_1, \dots, x_m)$  is a  $p$ -pointed,  $k$ -cube term for the variety axiomatized by  $\Sigma$  if there is a  $k \times m$  matrix  $M$  consisting of variables and  $p$  distinct constant symbols, with every column of  $M$  containing a symbol different from  $x$ , such that

$$\Sigma \models F(M) \approx \begin{pmatrix} x \\ \vdots \\ x \end{pmatrix}.$$

# Pointed cube terms

**Example:** Maltsev-term  $(x_1 x_2^{-1} x_3$  in groups).

$$m \begin{pmatrix} x & y & y \\ y & y & x \end{pmatrix} \approx \begin{pmatrix} x \\ x \end{pmatrix},$$

witnesses that  $m(x_1, x_2, x_3)$  is a 3-ary, 0-pointed, 2-cube term.

If  $\Sigma$  is a set of identities in a language  $\mathcal{L}$ , then an  $\mathcal{L}$ -term  $F(x_1, \dots, x_m)$  is a  $p$ -pointed,  $k$ -cube term for the variety axiomatized by  $\Sigma$  if there is a  $k \times m$  matrix  $M$  consisting of variables and  $p$  distinct constant symbols, with every column of  $M$  containing a symbol different from  $x$ , such that

$$\Sigma \models F(M) \approx \begin{pmatrix} x \\ \vdots \\ x \end{pmatrix}.$$

# Growth restrictions imposed by identities

## Theorem (KKSz)

Let  $\mathbf{A}$  be an algebra with an  $m$ -ary,  $p \geq 1$ -pointed,  $k$ -cube term, with at least one constant symbol appearing in the cube identities.

# Growth restrictions imposed by identities

## Theorem (KKSz)

Let  $\mathbf{A}$  be an algebra with an  $m$ -ary,  $p \geq 1$ -pointed,  $k$ -cube term, with at least one constant symbol appearing in the cube identities. If  $\mathbf{A}^{p+k-1}$  is finitely generated, then all finite powers of  $\mathbf{A}$  are finitely generated

# Growth restrictions imposed by identities

## Theorem (KKSz)

Let  $\mathbf{A}$  be an algebra with an  $m$ -ary,  $p \geq 1$ -pointed,  $k$ -cube term, with at least one constant symbol appearing in the cube identities. If  $\mathbf{A}^{p+k-1}$  is finitely generated, then all finite powers of  $\mathbf{A}$  are finitely generated and  $d_{\mathbf{A}}(n)$  is bounded above by a polynomial of degree at most  $\log_w(m)$ , where  $w = 2k/(2k-1)$ .

# Growth restrictions imposed by identities

## Theorem (KKSz)

Let  $\mathbf{A}$  be an algebra with an  $m$ -ary,  $p \geq 1$ -pointed,  $k$ -cube term, with at least one constant symbol appearing in the cube identities. If  $\mathbf{A}^{p+k-1}$  is finitely generated, then all finite powers of  $\mathbf{A}$  are finitely generated and  $d_{\mathbf{A}}(n)$  is bounded above by a polynomial of degree at most  $\log_w(m)$ , where  $w = 2k/(2k-1)$ .

- There exist finite algebras with pointed cube terms whose growth rate is  $\sim$  to a polynomial of any prescribed degree.



# Growth restrictions imposed by identities

## Theorem (KKSz)

Let  $\mathbf{A}$  be an algebra with an  $m$ -ary,  $p \geq 1$ -pointed,  $k$ -cube term, with at least one constant symbol appearing in the cube identities. If  $\mathbf{A}^{p+k-1}$  is finitely generated, then all finite powers of  $\mathbf{A}$  are finitely generated and  $d_{\mathbf{A}}(n)$  is bounded above by a polynomial of degree at most  $\log_w(m)$ , where  $w = 2k/(2k-1)$ .

- There exist finite algebras with pointed cube terms whose growth rate is  $\sim$  to a polynomial of any prescribed degree.
- The growth rate of any algebra with a pointed cube term arises as the growth rate of an algebra without a pointed cube term.

# Growth restrictions imposed by identities

## Theorem (KKSz)

Let  $\mathbf{A}$  be an algebra with an  $m$ -ary,  $p \geq 1$ -pointed,  $k$ -cube term, with at least one constant symbol appearing in the cube identities. If  $\mathbf{A}^{p+k-1}$  is finitely generated, then all finite powers of  $\mathbf{A}$  are finitely generated and  $d_{\mathbf{A}}(n)$  is bounded above by a polynomial of degree at most  $\log_w(m)$ , where  $w = 2k/(2k-1)$ .

- There exist finite algebras with pointed cube terms whose growth rate is  $\sim$  to a polynomial of any prescribed degree.
- The growth rate of any algebra with a pointed cube term arises as the growth rate of an algebra without a pointed cube term.
- If a **basic**  $\Sigma$  does not entail the existence of a pointed cube term, then  $\Sigma$  imposes no restriction on growth rates.  
“Basic” identity: at most one operation symbol on both sides.

# General Wiegold dichotomy

If **A** has a 0-pointed cube term, then it generates a congruence modular variety.

# General Wiegold dichotomy

If  $\mathbf{A}$  has a 0-pointed cube term, then it generates a congruence modular variety. We say that  $\mathbf{A}$  is **perfect**, if  $[1_{\mathbf{A}}, 1_{\mathbf{A}}] = 1_{\mathbf{A}}$  (in the sense of the modular commutator).

# General Wiegold dichotomy

If  $\mathbf{A}$  has a 0-pointed cube term, then it generates a congruence modular variety. We say that  $\mathbf{A}$  is **perfect**, if  $[1_{\mathbf{A}}, 1_{\mathbf{A}}] = 1_{\mathbf{A}}$  (in the sense of the modular commutator).

That is,  $\mathbf{A}$  is perfect iff it has no nontrivial abelian factor algebras.

# General Wiegold dichotomy

If  $\mathbf{A}$  has a 0-pointed cube term, then it generates a congruence modular variety. We say that  $\mathbf{A}$  is **perfect**, if  $[1_{\mathbf{A}}, 1_{\mathbf{A}}] = 1_{\mathbf{A}}$  (in the sense of the modular commutator). That is,  $\mathbf{A}$  is perfect iff it has no nontrivial abelian factor algebras.

## Theorem (KKSz)

Suppose that an algebra  $\mathbf{A}$  has a 0-pointed,  $k$ -cube term

# General Wiegold dichotomy

If  $\mathbf{A}$  has a 0-pointed cube term, then it generates a congruence modular variety. We say that  $\mathbf{A}$  is **perfect**, if  $[1_{\mathbf{A}}, 1_{\mathbf{A}}] = 1_{\mathbf{A}}$  (in the sense of the modular commutator).

That is,  $\mathbf{A}$  is perfect iff it has no nontrivial abelian factor algebras.

## Theorem (KKSz)

Suppose that an algebra  $\mathbf{A}$  has a 0-pointed,  $k$ -cube term and  $\mathbf{A}^k$  is finitely generated.

# General Wiegold dichotomy

If  $\mathbf{A}$  has a 0-pointed cube term, then it generates a congruence modular variety. We say that  $\mathbf{A}$  is **perfect**, if  $[1_{\mathbf{A}}, 1_{\mathbf{A}}] = 1_{\mathbf{A}}$  (in the sense of the modular commutator).

That is,  $\mathbf{A}$  is perfect iff it has no nontrivial abelian factor algebras.

## Theorem (KKSz)

Suppose that an algebra  $\mathbf{A}$  has a 0-pointed,  $k$ -cube term and  $\mathbf{A}^k$  is finitely generated.

- $\mathbf{A}$  perfect  $\implies d_{\mathbf{A}}(n) = O(\log(n))$ .



# General Wiegold dichotomy

If  $\mathbf{A}$  has a 0-pointed cube term, then it generates a congruence modular variety. We say that  $\mathbf{A}$  is **perfect**, if  $[1_{\mathbf{A}}, 1_{\mathbf{A}}] = 1_{\mathbf{A}}$  (in the sense of the modular commutator).

That is,  $\mathbf{A}$  is perfect iff it has no nontrivial abelian factor algebras.

## Theorem (KKSz)

Suppose that an algebra  $\mathbf{A}$  has a 0-pointed,  $k$ -cube term and  $\mathbf{A}^k$  is finitely generated.

- $\mathbf{A}$  perfect  $\implies d_{\mathbf{A}}(n) = O(\log(n))$ .
- $\mathbf{A}$  imperfect  $\implies d_{\mathbf{A}}(n) = O(n)$ .

# General Wiegold dichotomy

If  $\mathbf{A}$  has a 0-pointed cube term, then it generates a congruence modular variety. We say that  $\mathbf{A}$  is **perfect**, if  $[1_{\mathbf{A}}, 1_{\mathbf{A}}] = 1_{\mathbf{A}}$  (in the sense of the modular commutator).

That is,  $\mathbf{A}$  is perfect iff it has no nontrivial abelian factor algebras.

## Theorem (KKSz)

Suppose that an algebra  $\mathbf{A}$  has a 0-pointed,  $k$ -cube term and  $\mathbf{A}^k$  is finitely generated.

- $\mathbf{A}$  perfect  $\implies d_{\mathbf{A}}(n) = O(\log(n))$ .  
   $\mathbf{A}$  imperfect  $\implies d_{\mathbf{A}}(n) = O(n)$ .
- Suppose that  $\mathbf{A}$  is finite.

# General Wiegold dichotomy

If  $\mathbf{A}$  has a 0-pointed cube term, then it generates a congruence modular variety. We say that  $\mathbf{A}$  is **perfect**, if  $[1_{\mathbf{A}}, 1_{\mathbf{A}}] = 1_{\mathbf{A}}$  (in the sense of the modular commutator).

That is,  $\mathbf{A}$  is perfect iff it has no nontrivial abelian factor algebras.

## Theorem (KKSz)

Suppose that an algebra  $\mathbf{A}$  has a 0-pointed,  $k$ -cube term and  $\mathbf{A}^k$  is finitely generated.

- $\mathbf{A}$  perfect  $\implies d_{\mathbf{A}}(n) = O(\log(n))$ .  
   $\mathbf{A}$  imperfect  $\implies d_{\mathbf{A}}(n) = O(n)$ .
- Suppose that  $\mathbf{A}$  is finite.  $\mathbf{A}$  perfect  $\implies d_{\mathbf{A}}(n) = \Theta(\log(n))$ .

# General Wiegold dichotomy

If  $\mathbf{A}$  has a 0-pointed cube term, then it generates a congruence modular variety. We say that  $\mathbf{A}$  is **perfect**, if  $[1_{\mathbf{A}}, 1_{\mathbf{A}}] = 1_{\mathbf{A}}$  (in the sense of the modular commutator).

That is,  $\mathbf{A}$  is perfect iff it has no nontrivial abelian factor algebras.

## Theorem (KKSz)

Suppose that an algebra  $\mathbf{A}$  has a 0-pointed,  $k$ -cube term and  $\mathbf{A}^k$  is finitely generated.

- $\mathbf{A}$  perfect  $\implies d_{\mathbf{A}}(n) = O(\log(n))$ .  
   $\mathbf{A}$  imperfect  $\implies d_{\mathbf{A}}(n) = O(n)$ .
- Suppose that  $\mathbf{A}$  is finite.  $\mathbf{A}$  perfect  $\implies d_{\mathbf{A}}(n) = \Theta(\log(n))$ .  
   $\mathbf{A}$  imperfect  $\implies d_{\mathbf{A}}(n) = \Theta(n)$ .

# General Wiegold dichotomy

If  $\mathbf{A}$  has a 0-pointed cube term, then it generates a congruence modular variety. We say that  $\mathbf{A}$  is **perfect**, if  $[1_{\mathbf{A}}, 1_{\mathbf{A}}] = 1_{\mathbf{A}}$  (in the sense of the modular commutator).

That is,  $\mathbf{A}$  is perfect iff it has no nontrivial abelian factor algebras.

## Theorem (KKSz)

Suppose that an algebra  $\mathbf{A}$  has a 0-pointed,  $k$ -cube term and  $\mathbf{A}^k$  is finitely generated.

- $\mathbf{A}$  perfect  $\implies d_{\mathbf{A}}(n) = O(\log(n))$ .  
 $\mathbf{A}$  imperfect  $\implies d_{\mathbf{A}}(n) = O(n)$ .
- Suppose that  $\mathbf{A}$  is finite.  $\mathbf{A}$  perfect  $\implies d_{\mathbf{A}}(n) = \Theta(\log(n))$ .  
 $\mathbf{A}$  imperfect  $\implies d_{\mathbf{A}}(n) = \Theta(n)$ .

The proof uses a probabilistic argument of independent interest.

# Abelianness properties

By R. McKenzie and D. Hobby in tame congruence theory:

If  $\alpha, \beta, \delta \in \text{Con}(\mathbf{A})$ , then  $\alpha$  centralizes  $\beta$  modulo  $\delta$ , that is,  $C(\alpha, \beta; \delta)$  holds iff

# Abelianness properties

By R. McKenzie and D. Hobby in tame congruence theory:

If  $\alpha, \beta, \delta \in \text{Con}(\mathbf{A})$ , then  $\alpha$  centralizes  $\beta$  modulo  $\delta$ , that is,

$C(\alpha, \beta; \delta)$  holds iff for all polynomials  $t$  of  $\mathbf{A}$  we have

$$(\forall \mathbf{a} \equiv_{\alpha} \mathbf{b})(\forall \mathbf{c} \equiv_{\beta} \mathbf{d}) \quad t(\mathbf{a}, \mathbf{c}) \equiv_{\delta} t(\mathbf{a}, \mathbf{d}) \implies t(\mathbf{b}, \mathbf{c}) \equiv_{\delta} t(\mathbf{b}, \mathbf{d}).$$

# Abelianness properties

By R. McKenzie and D. Hobby in tame congruence theory:

If  $\alpha, \beta, \delta \in \text{Con}(\mathbf{A})$ , then  $\alpha$  centralizes  $\beta$  modulo  $\delta$ , that is,

$C(\alpha, \beta; \delta)$  holds iff for all polynomials  $t$  of  $\mathbf{A}$  we have

$$(\forall \mathbf{a} \equiv_{\alpha} \mathbf{b})(\forall \mathbf{c} \equiv_{\beta} \mathbf{d}) \quad t(\mathbf{a}, \mathbf{c}) \equiv_{\delta} t(\mathbf{a}, \mathbf{d}) \implies t(\mathbf{b}, \mathbf{c}) \equiv_{\delta} t(\mathbf{b}, \mathbf{d}).$$

The **commutator**:  $[\alpha, \beta] = \bigwedge \{ \delta \in \text{Con}(\mathbf{A}) : C(\alpha, \beta; \delta) \text{ holds} \}.$



# Abelianness properties

By R. McKenzie and D. Hobby in tame congruence theory:

If  $\alpha, \beta, \delta \in \text{Con}(\mathbf{A})$ , then  $\alpha$  centralizes  $\beta$  modulo  $\delta$ , that is,

$C(\alpha, \beta; \delta)$  holds iff for all polynomials  $t$  of  $\mathbf{A}$  we have

$$(\forall \mathbf{a} \equiv_{\alpha} \mathbf{b})(\forall \mathbf{c} \equiv_{\beta} \mathbf{d}) \quad t(\mathbf{a}, \mathbf{c}) \equiv_{\delta} t(\mathbf{a}, \mathbf{d}) \implies t(\mathbf{b}, \mathbf{c}) \equiv_{\delta} t(\mathbf{b}, \mathbf{d}).$$

The **commutator**:  $[\alpha, \beta] = \bigwedge \{ \delta \in \text{Con}(\mathbf{A}) : C(\alpha, \beta; \delta) \text{ holds} \}.$

$\mathbf{A}$  is **abelian** if  $[1_{\mathbf{A}}, 1_{\mathbf{A}}] = 0_{\mathbf{A}}$

# Abelianness properties

By R. McKenzie and D. Hobby in tame congruence theory:

If  $\alpha, \beta, \delta \in \text{Con}(\mathbf{A})$ , then  $\alpha$  centralizes  $\beta$  modulo  $\delta$ , that is,

$C(\alpha, \beta; \delta)$  holds iff for all polynomials  $t$  of  $\mathbf{A}$  we have

$$(\forall \mathbf{a} \equiv_{\alpha} \mathbf{b})(\forall \mathbf{c} \equiv_{\beta} \mathbf{d}) \quad t(\mathbf{a}, \mathbf{c}) \equiv_{\delta} t(\mathbf{a}, \mathbf{d}) \implies t(\mathbf{b}, \mathbf{c}) \equiv_{\delta} t(\mathbf{b}, \mathbf{d}).$$

The **commutator**:  $[\alpha, \beta] = \bigwedge \{ \delta \in \text{Con}(\mathbf{A}) : C(\alpha, \beta; \delta) \text{ holds} \}.$

$\mathbf{A}$  is **abelian** if  $[1_{\mathbf{A}}, 1_{\mathbf{A}}] = 0_{\mathbf{A}}$  (that is,  $C(1_{\mathbf{A}}, 1_{\mathbf{A}}; 0_{\mathbf{A}})$  holds).

# Abelianness properties

By R. McKenzie and D. Hobby in tame congruence theory:

If  $\alpha, \beta, \delta \in \text{Con}(\mathbf{A})$ , then  $\alpha$  centralizes  $\beta$  modulo  $\delta$ , that is,

$C(\alpha, \beta; \delta)$  holds iff for all polynomials  $t$  of  $\mathbf{A}$  we have

$$(\forall \mathbf{a} \equiv_{\alpha} \mathbf{b})(\forall \mathbf{c} \equiv_{\beta} \mathbf{d}) \quad t(\mathbf{a}, \mathbf{c}) \equiv_{\delta} t(\mathbf{a}, \mathbf{d}) \implies t(\mathbf{b}, \mathbf{c}) \equiv_{\delta} t(\mathbf{b}, \mathbf{d}).$$

The **commutator**:  $[\alpha, \beta] = \bigwedge \{ \delta \in \text{Con}(\mathbf{A}) : C(\alpha, \beta; \delta) \text{ holds} \}.$

$\mathbf{A}$  is **abelian** if  $[1_{\mathbf{A}}, 1_{\mathbf{A}}] = 0_{\mathbf{A}}$  (that is,  $C(1_{\mathbf{A}}, 1_{\mathbf{A}}; 0_{\mathbf{A}})$  holds).

Homomorphic images of abelian algebras are not always abelian.

# Abelianness properties

By R. McKenzie and D. Hobby in tame congruence theory:

If  $\alpha, \beta, \delta \in \text{Con}(\mathbf{A})$ , then  $\alpha$  centralizes  $\beta$  modulo  $\delta$ , that is,

$C(\alpha, \beta; \delta)$  holds iff for all polynomials  $t$  of  $\mathbf{A}$  we have

$$(\forall \mathbf{a} \equiv_{\alpha} \mathbf{b})(\forall \mathbf{c} \equiv_{\beta} \mathbf{d}) \quad t(\mathbf{a}, \mathbf{c}) \equiv_{\delta} t(\mathbf{a}, \mathbf{d}) \implies t(\mathbf{b}, \mathbf{c}) \equiv_{\delta} t(\mathbf{b}, \mathbf{d}).$$

The **commutator**:  $[\alpha, \beta] = \bigwedge \{ \delta \in \text{Con}(\mathbf{A}) : C(\alpha, \beta; \delta) \text{ holds} \}.$

$\mathbf{A}$  is **abelian** if  $[1_{\mathbf{A}}, 1_{\mathbf{A}}] = 0_{\mathbf{A}}$  (that is,  $C(1_{\mathbf{A}}, 1_{\mathbf{A}}; 0_{\mathbf{A}})$  holds).

Homomorphic images of abelian algebras are not always abelian.

$\mathbf{A}$  is **solvable**, if there is a chain of congruences

$$0_{\mathbf{A}} = \theta_0 < \theta_1 < \dots < \theta_n = 1_{\mathbf{A}}$$

# Abelianness properties

By R. McKenzie and D. Hobby in tame congruence theory:

If  $\alpha, \beta, \delta \in \text{Con}(\mathbf{A})$ , then  $\alpha$  centralizes  $\beta$  modulo  $\delta$ , that is,

$C(\alpha, \beta; \delta)$  holds iff for all polynomials  $t$  of  $\mathbf{A}$  we have

$$(\forall \mathbf{a} \equiv_{\alpha} \mathbf{b})(\forall \mathbf{c} \equiv_{\beta} \mathbf{d}) \quad t(\mathbf{a}, \mathbf{c}) \equiv_{\delta} t(\mathbf{a}, \mathbf{d}) \implies t(\mathbf{b}, \mathbf{c}) \equiv_{\delta} t(\mathbf{b}, \mathbf{d}).$$

The **commutator**:  $[\alpha, \beta] = \bigwedge \{ \delta \in \text{Con}(\mathbf{A}) : C(\alpha, \beta; \delta) \text{ holds} \}.$

$\mathbf{A}$  is **abelian** if  $[1_{\mathbf{A}}, 1_{\mathbf{A}}] = 0_{\mathbf{A}}$  (that is,  $C(1_{\mathbf{A}}, 1_{\mathbf{A}}; 0_{\mathbf{A}})$  holds).

Homomorphic images of abelian algebras are not always abelian.

$\mathbf{A}$  is **solvable**, if there is a chain of congruences

$0_{\mathbf{A}} = \theta_0 < \theta_1 < \dots < \theta_n = 1_{\mathbf{A}}$  such that each  $\theta_{i+1}/\theta_i$  is abelian.

# Abelianness properties

By R. McKenzie and D. Hobby in tame congruence theory:

If  $\alpha, \beta, \delta \in \text{Con}(\mathbf{A})$ , then  $\alpha$  centralizes  $\beta$  modulo  $\delta$ , that is,

$C(\alpha, \beta; \delta)$  holds iff for all polynomials  $t$  of  $\mathbf{A}$  we have

$$(\forall \mathbf{a} \equiv_{\alpha} \mathbf{b})(\forall \mathbf{c} \equiv_{\beta} \mathbf{d}) \quad t(\mathbf{a}, \mathbf{c}) \equiv_{\delta} t(\mathbf{a}, \mathbf{d}) \implies t(\mathbf{b}, \mathbf{c}) \equiv_{\delta} t(\mathbf{b}, \mathbf{d}).$$

The **commutator**:  $[\alpha, \beta] = \bigwedge \{ \delta \in \text{Con}(\mathbf{A}) : C(\alpha, \beta; \delta) \text{ holds} \}.$

$\mathbf{A}$  is **abelian** if  $[1_{\mathbf{A}}, 1_{\mathbf{A}}] = 0_{\mathbf{A}}$  (that is,  $C(1_{\mathbf{A}}, 1_{\mathbf{A}}; 0_{\mathbf{A}})$  holds).

Homomorphic images of abelian algebras are not always abelian.

$\mathbf{A}$  is **solvable**, if there is a chain of congruences

$0_{\mathbf{A}} = \theta_0 < \theta_1 < \dots < \theta_n = 1_{\mathbf{A}}$  such that each  $\theta_{i+1}/\theta_i$  is abelian.

Can be expressed with the commutator the same way as for groups.

# Abelianness properties

By R. McKenzie and D. Hobby in tame congruence theory:

If  $\alpha, \beta, \delta \in \text{Con}(\mathbf{A})$ , then  $\alpha$  centralizes  $\beta$  modulo  $\delta$ , that is,

$C(\alpha, \beta; \delta)$  holds iff for all polynomials  $t$  of  $\mathbf{A}$  we have

$$(\forall \mathbf{a} \equiv_{\alpha} \mathbf{b})(\forall \mathbf{c} \equiv_{\beta} \mathbf{d}) \quad t(\mathbf{a}, \mathbf{c}) \equiv_{\delta} t(\mathbf{a}, \mathbf{d}) \implies t(\mathbf{b}, \mathbf{c}) \equiv_{\delta} t(\mathbf{b}, \mathbf{d}).$$

The **commutator**:  $[\alpha, \beta] = \bigwedge \{ \delta \in \text{Con}(\mathbf{A}) : C(\alpha, \beta; \delta) \text{ holds} \}.$

$\mathbf{A}$  is **abelian** if  $[1_{\mathbf{A}}, 1_{\mathbf{A}}] = 0_{\mathbf{A}}$  (that is,  $C(1_{\mathbf{A}}, 1_{\mathbf{A}}; 0_{\mathbf{A}})$  holds).

Homomorphic images of abelian algebras are not always abelian.

$\mathbf{A}$  is **solvable**, if there is a chain of congruences

$0_{\mathbf{A}} = \theta_0 < \theta_1 < \dots < \theta_n = 1_{\mathbf{A}}$  such that each  $\theta_{i+1}/\theta_i$  is abelian.

Can be expressed with the commutator the same way as for groups.

Homomorphic images,

# Abelianness properties

By R. McKenzie and D. Hobby in tame congruence theory:

If  $\alpha, \beta, \delta \in \text{Con}(\mathbf{A})$ , then  $\alpha$  centralizes  $\beta$  modulo  $\delta$ , that is,

$C(\alpha, \beta; \delta)$  holds iff for all polynomials  $t$  of  $\mathbf{A}$  we have

$$(\forall \mathbf{a} \equiv_{\alpha} \mathbf{b})(\forall \mathbf{c} \equiv_{\beta} \mathbf{d}) \quad t(\mathbf{a}, \mathbf{c}) \equiv_{\delta} t(\mathbf{a}, \mathbf{d}) \implies t(\mathbf{b}, \mathbf{c}) \equiv_{\delta} t(\mathbf{b}, \mathbf{d}).$$

The **commutator**:  $[\alpha, \beta] = \bigwedge \{ \delta \in \text{Con}(\mathbf{A}) : C(\alpha, \beta; \delta) \text{ holds} \}.$

$\mathbf{A}$  is **abelian** if  $[1_{\mathbf{A}}, 1_{\mathbf{A}}] = 0_{\mathbf{A}}$  (that is,  $C(1_{\mathbf{A}}, 1_{\mathbf{A}}; 0_{\mathbf{A}})$  holds).

Homomorphic images of abelian algebras are not always abelian.

$\mathbf{A}$  is **solvable**, if there is a chain of congruences

$0_{\mathbf{A}} = \theta_0 < \theta_1 < \dots < \theta_n = 1_{\mathbf{A}}$  such that each  $\theta_{i+1}/\theta_i$  is abelian.

Can be expressed with the commutator the same way as for groups.

Homomorphic images, direct products



# Abelianness properties

By R. McKenzie and D. Hobby in tame congruence theory:

If  $\alpha, \beta, \delta \in \text{Con}(\mathbf{A})$ , then  $\alpha$  centralizes  $\beta$  modulo  $\delta$ , that is,

$C(\alpha, \beta; \delta)$  holds iff for all polynomials  $t$  of  $\mathbf{A}$  we have

$$(\forall \mathbf{a} \equiv_{\alpha} \mathbf{b})(\forall \mathbf{c} \equiv_{\beta} \mathbf{d}) \quad t(\mathbf{a}, \mathbf{c}) \equiv_{\delta} t(\mathbf{a}, \mathbf{d}) \implies t(\mathbf{b}, \mathbf{c}) \equiv_{\delta} t(\mathbf{b}, \mathbf{d}).$$

The **commutator**:  $[\alpha, \beta] = \bigwedge \{ \delta \in \text{Con}(\mathbf{A}) : C(\alpha, \beta; \delta) \text{ holds} \}.$

$\mathbf{A}$  is **abelian** if  $[1_{\mathbf{A}}, 1_{\mathbf{A}}] = 0_{\mathbf{A}}$  (that is,  $C(1_{\mathbf{A}}, 1_{\mathbf{A}}; 0_{\mathbf{A}})$  holds).

Homomorphic images of abelian algebras are not always abelian.

$\mathbf{A}$  is **solvable**, if there is a chain of congruences

$0_{\mathbf{A}} = \theta_0 < \theta_1 < \dots < \theta_n = 1_{\mathbf{A}}$  such that each  $\theta_{i+1}/\theta_i$  is abelian.

Can be expressed with the commutator the same way as for groups.

Homomorphic images, direct products and subalgebras of finite solvable algebras are solvable.

# Abelianness properties

By R. McKenzie and D. Hobby in tame congruence theory:

If  $\alpha, \beta, \delta \in \text{Con}(\mathbf{A})$ , then  $\alpha$  centralizes  $\beta$  modulo  $\delta$ , that is,

$C(\alpha, \beta; \delta)$  holds iff for all polynomials  $t$  of  $\mathbf{A}$  we have

$$(\forall \mathbf{a} \equiv_{\alpha} \mathbf{b})(\forall \mathbf{c} \equiv_{\beta} \mathbf{d}) \quad t(\mathbf{a}, \mathbf{c}) \equiv_{\delta} t(\mathbf{a}, \mathbf{d}) \implies t(\mathbf{b}, \mathbf{c}) \equiv_{\delta} t(\mathbf{b}, \mathbf{d}).$$

The **commutator**:  $[\alpha, \beta] = \bigwedge \{ \delta \in \text{Con}(\mathbf{A}) : C(\alpha, \beta; \delta) \text{ holds} \}.$

$\mathbf{A}$  is **abelian** if  $[1_{\mathbf{A}}, 1_{\mathbf{A}}] = 0_{\mathbf{A}}$  (that is,  $C(1_{\mathbf{A}}, 1_{\mathbf{A}}; 0_{\mathbf{A}})$  holds).

Homomorphic images of abelian algebras are not always abelian.

$\mathbf{A}$  is **solvable**, if there is a chain of congruences

$0_{\mathbf{A}} = \theta_0 < \theta_1 < \dots < \theta_n = 1_{\mathbf{A}}$  such that each  $\theta_{i+1}/\theta_i$  is abelian.

Can be expressed with the commutator the same way as for groups.

Homomorphic images, direct products and subalgebras of finite solvable algebras are solvable. A finite algebra is solvable iff only the types **1** and **2** of tame congruence theory occur in it.

# Nilpotence

Developed by K. Kearnes:

When  $\alpha \in \text{Con}(\mathbf{A})$  define  $(\alpha)^1 = [\alpha]^1 = \alpha$

# Nilpotence

Developed by K. Kearnes:

When  $\alpha \in \text{Con}(\mathbf{A})$  define  $(\alpha)^1 = [\alpha]^1 = \alpha$  and

$$(\alpha)^{k+1} = [\alpha, (\alpha)^k], \quad [\alpha]^{k+1} = [[\alpha]^k, \alpha].$$

# Nilpotence

Developed by K. Kearnes:

When  $\alpha \in \text{Con}(\mathbf{A})$  define  $(\alpha)^1 = [\alpha]^1 = \alpha$  and

$$(\alpha)^{k+1} = [\alpha, (\alpha)^k], \quad [\alpha]^{k+1} = [[\alpha]^k, \alpha].$$

If  $(\alpha)^{n+1} = 0$ , then  $\alpha$  is  $n$ -step left nilpotent,

# Nilpotence

Developed by K. Kearnes:

When  $\alpha \in \text{Con}(\mathbf{A})$  define  $(\alpha)^1 = [\alpha]^1 = \alpha$  and

$$(\alpha)^{k+1} = [\alpha, (\alpha)^k], \quad [\alpha]^{k+1} = [[\alpha]^k, \alpha].$$

If  $(\alpha)^{n+1} = 0$ , then  $\alpha$  is  $n$ -step left nilpotent,

if  $[\alpha]^{n+1} = 0$ , then  $\alpha$  is  $n$ -step right nilpotent.

# Nilpotence

Developed by K. Kearnes:

When  $\alpha \in \text{Con}(\mathbf{A})$  define  $(\alpha)^1 = [\alpha]^1 = \alpha$  and

$$(\alpha)^{k+1} = [\alpha, (\alpha)^k], \quad [\alpha]^{k+1} = [[\alpha]^k, \alpha].$$

If  $(\alpha)^{n+1} = 0$ , then  $\alpha$  is  $n$ -step left nilpotent,

if  $[\alpha]^{n+1} = 0$ , then  $\alpha$  is  $n$ -step right nilpotent.

Right nilpotent congruences are left nilpotent in finite algebras.

# Nilpotence

Developed by K. Kearnes:

When  $\alpha \in \text{Con}(\mathbf{A})$  define  $(\alpha)^1 = [\alpha]^1 = \alpha$  and

$$(\alpha)^{k+1} = [\alpha, (\alpha)^k], \quad [\alpha]^{k+1} = [[\alpha]^k, \alpha].$$

If  $(\alpha)^{n+1} = 0$ , then  $\alpha$  is  $n$ -step left nilpotent,

if  $[\alpha]^{n+1} = 0$ , then  $\alpha$  is  $n$ -step right nilpotent.

Right nilpotent congruences are left nilpotent in finite algebras.

Left nilpotence implies the following condition:

$C(1_{\mathbf{A}}, N^2; \delta)$  holds whenever  $\delta \prec \theta$  and  $N$  is a  $\langle \delta, \theta \rangle$ -trace.  $(\dagger)$



# Nilpotence

Developed by K. Kearnes:

When  $\alpha \in \text{Con}(\mathbf{A})$  define  $(\alpha)^1 = [\alpha]^1 = \alpha$  and

$$(\alpha)^{k+1} = [\alpha, (\alpha)^k], \quad [\alpha]^{k+1} = [[\alpha]^k, \alpha].$$

If  $(\alpha)^{n+1} = 0$ , then  $\alpha$  is  $n$ -step left nilpotent,

if  $[\alpha]^{n+1} = 0$ , then  $\alpha$  is  $n$ -step right nilpotent.

Right nilpotent congruences are left nilpotent in finite algebras.

Left nilpotence implies the following condition:

$C(1_{\mathbf{A}}, N^2; \delta)$  holds whenever  $\delta \prec \theta$  and  $N$  is a  $\langle \delta, \theta \rangle$ -trace.  $(\dagger)$

(Here  $N^2$  is considered as a binary relation, and centrality is defined naturally).

# Nilpotence

Developed by K. Kearnes:

When  $\alpha \in \text{Con}(\mathbf{A})$  define  $(\alpha)^1 = [\alpha]^1 = \alpha$  and

$$(\alpha)^{k+1} = [\alpha, (\alpha)^k], \quad [\alpha]^{k+1} = [[\alpha]^k, \alpha].$$

If  $(\alpha)^{n+1} = 0$ , then  $\alpha$  is  $n$ -step left nilpotent,

if  $[\alpha]^{n+1} = 0$ , then  $\alpha$  is  $n$ -step right nilpotent.

Right nilpotent congruences are left nilpotent in finite algebras.

Left nilpotence implies the following condition:

$C(1_{\mathbf{A}}, N^2; \delta)$  holds whenever  $\delta \prec \theta$  and  $N$  is a  $\langle \delta, \theta \rangle$ -trace.  $(\dagger)$

(Here  $N^2$  is considered as a binary relation, and centrality is defined naturally). This condition is still stronger than solvability.

# Nilpotence

Developed by K. Kearnes:

When  $\alpha \in \text{Con}(\mathbf{A})$  define  $(\alpha)^1 = [\alpha]^1 = \alpha$  and

$$(\alpha)^{k+1} = [\alpha, (\alpha)^k], \quad [\alpha]^{k+1} = [[\alpha]^k, \alpha].$$

If  $(\alpha)^{n+1} = 0$ , then  $\alpha$  is  $n$ -step left nilpotent,

if  $[\alpha]^{n+1} = 0$ , then  $\alpha$  is  $n$ -step right nilpotent.

Right nilpotent congruences are left nilpotent in finite algebras.

Left nilpotence implies the following condition:

$C(1_{\mathbf{A}}, N^2; \delta)$  holds whenever  $\delta \prec \theta$  and  $N$  is a  $\langle \delta, \theta \rangle$ -trace.  $(\dagger)$

(Here  $N^2$  is considered as a binary relation, and centrality is defined naturally). This condition is still stronger than solvability.

Theorem (K. Kearnes)

Homomorphic images of finite abelian algebras are right nilpotent.

# The Hamiltonian property

**A** is **Hamiltonian**: every subalgebra is a congruence block.

# The Hamiltonian property

**A** is **Hamiltonian**: every subalgebra is a congruence block.

**quasi-Hamiltonian**: every maximal subalgebra is a congruence-block.

# The Hamiltonian property

**A** is **Hamiltonian**: every subalgebra is a congruence block.

**quasi-Hamiltonian**: every maximal subalgebra is a congruence-block.

Theorem (E. W. Kiss, M. Valeriote)

A locally finite variety is abelian iff it is Hamiltonian.

# The Hamiltonian property

**A** is **Hamiltonian**: every subalgebra is a congruence block.

**quasi-Hamiltonian**: every maximal subalgebra is a congruence-block.

**Theorem (E. W. Kiss, M. Valeriote)**

A locally finite variety is abelian iff it is Hamiltonian.

**Wielandt**: A finite group is quasi-Hamiltonian (that is, every maximal subgroup is normal) iff it is nilpotent.

# The Hamiltonian property

**A** is **Hamiltonian**: every subalgebra is a congruence block.

**quasi-Hamiltonian**: every maximal subalgebra is a congruence-block.

Theorem (E. W. Kiss, M. Valeriote)

A locally finite variety is abelian iff it is Hamiltonian.

**Wielandt**: A finite group is quasi-Hamiltonian (that is, every maximal subgroup is normal) iff it is nilpotent.

Theorem (K. Kearnes)

If a finite algebra **A** satisfies  $(\dagger)$ , then it is quasi-Hamiltonian.



# The Hamiltonian property

**A** is **Hamiltonian**: every subalgebra is a congruence block.

**quasi-Hamiltonian**: every maximal subalgebra is a congruence-block.

Theorem (E. W. Kiss, M. Valeriote)

A locally finite variety is abelian iff it is Hamiltonian.

**Wielandt**: A finite group is quasi-Hamiltonian (that is, every maximal subgroup is normal) iff it is nilpotent.

Theorem (K. Kearnes)

If a finite algebra **A** satisfies  $(\dagger)$ , then it is quasi-Hamiltonian.

A variety generated by a finite left nilpotent algebra is quasi-Hamiltonian.

# The Hamiltonian property

**A** is **Hamiltonian**: every subalgebra is a congruence block.

**quasi-Hamiltonian**: every maximal subalgebra is a congruence-block.

Theorem (E. W. Kiss, M. Valerioté)

A locally finite variety is abelian iff it is Hamiltonian.

**Wielandt**: A finite group is quasi-Hamiltonian (that is, every maximal subgroup is normal) iff it is nilpotent.

Theorem (K. Kearnes)

If a finite algebra **A** satisfies  $(\dagger)$ , then it is quasi-Hamiltonian.

A variety generated by a finite left nilpotent algebra is quasi-Hamiltonian. Conversely, if  $\mathbf{A}^2$  is quasi-Hamiltonian, then  $V(\mathbf{A})$  is quasi-Hamiltonian,

# The Hamiltonian property

**A** is **Hamiltonian**: every subalgebra is a congruence block.

**quasi-Hamiltonian**: every maximal subalgebra is a congruence-block.

Theorem (E. W. Kiss, M. Valerioté)

A locally finite variety is abelian iff it is Hamiltonian.

**Wielandt**: A finite group is quasi-Hamiltonian (that is, every maximal subgroup is normal) iff it is nilpotent.

Theorem (K. Kearnes)

If a finite algebra **A** satisfies  $(\dagger)$ , then it is quasi-Hamiltonian.

A variety generated by a finite left nilpotent algebra is quasi-Hamiltonian. Conversely, if  $\mathbf{A}^2$  is quasi-Hamiltonian, then  $\mathbf{V}(\mathbf{A})$  is quasi-Hamiltonian, and its finite members satisfy  $(\dagger)$ .

# Strongly abelian algebras

An algebra **A** is strongly abelian, if for all polynomials  $t$  we have

$$(\forall \mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}, \mathbf{e}) \quad t(\mathbf{a}, \mathbf{c}) = t(\mathbf{b}, \mathbf{d}) \implies t(\mathbf{e}, \mathbf{c}) = t(\mathbf{e}, \mathbf{d}).$$

# Strongly abelian algebras

An algebra **A** is strongly abelian, if for all polynomials  $t$  we have  
 $(\forall \mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}, \mathbf{e}) \quad t(\mathbf{a}, \mathbf{c}) = t(\mathbf{b}, \mathbf{d}) \implies t(\mathbf{e}, \mathbf{c}) = t(\mathbf{e}, \mathbf{d}).$

Let **A** be a nontrivial finite algebra and let **B** be a nontrivial homomorphic image of  $\mathbf{A}^k$  for some  $k$ .

# Strongly abelian algebras

An algebra **A** is strongly abelian, if for all polynomials  $t$  we have  
 $(\forall \mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}, \mathbf{e}) \quad t(\mathbf{a}, \mathbf{c}) = t(\mathbf{b}, \mathbf{d}) \implies t(\mathbf{e}, \mathbf{c}) = t(\mathbf{e}, \mathbf{d}).$

Let **A** be a nontrivial finite algebra and let **B** be a nontrivial homomorphic image of  $\mathbf{A}^k$  for some  $k$ .

- If **B** is strongly abelian, then  $d_{\mathbf{A}}(n) = 2^{\Theta(n)}$  (exponential).

# Strongly abelian algebras

An algebra **A** is strongly abelian, if for all polynomials  $t$  we have  
 $(\forall \mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}, \mathbf{e}) \quad t(\mathbf{a}, \mathbf{c}) = t(\mathbf{b}, \mathbf{d}) \implies t(\mathbf{e}, \mathbf{c}) = t(\mathbf{e}, \mathbf{d}).$

Let **A** be a nontrivial finite algebra and let **B** be a nontrivial homomorphic image of  $\mathbf{A}^k$  for some  $k$ .

- If **B** is strongly abelian, then  $d_{\mathbf{A}}(n) = 2^{\Theta(n)}$  (exponential).
- If **B** is abelian, then  $d_{\mathbf{A}}(n) = \Omega(n)$  (at least linear).

# Strongly abelian algebras

An algebra **A** is strongly abelian, if for all polynomials  $t$  we have  
 $(\forall \mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}, \mathbf{e}) \quad t(\mathbf{a}, \mathbf{c}) = t(\mathbf{b}, \mathbf{d}) \implies t(\mathbf{e}, \mathbf{c}) = t(\mathbf{e}, \mathbf{d}).$

Let **A** be a nontrivial finite algebra and let **B** be a nontrivial homomorphic image of  $\mathbf{A}^k$  for some  $k$ .

- If **B** is strongly abelian, then  $d_{\mathbf{A}}(n) = 2^{\Theta(n)}$  (exponential).
- If **B** is abelian, then  $d_{\mathbf{A}}(n) = \Omega(n)$  (at least linear).

This holds, because the free algebras in the first case have polynomially bounded size,



# Strongly abelian algebras

An algebra **A** is strongly abelian, if for all polynomials  $t$  we have  
 $(\forall \mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}, \mathbf{e}) \quad t(\mathbf{a}, \mathbf{c}) = t(\mathbf{b}, \mathbf{d}) \implies t(\mathbf{e}, \mathbf{c}) = t(\mathbf{e}, \mathbf{d}).$

Let **A** be a nontrivial finite algebra and let **B** be a nontrivial homomorphic image of  $\mathbf{A}^k$  for some  $k$ .

- If **B** is strongly abelian, then  $d_{\mathbf{A}}(n) = 2^{\Theta(n)}$  (exponential).
- If **B** is abelian, then  $d_{\mathbf{A}}(n) = \Omega(n)$  (at least linear).

This holds, because the free algebras in the first case have polynomially bounded size, and in the second case their size is in  $2^{O(n)}$  by a result of J. Berman and R. McKenzie.

# Strongly abelian algebras

An algebra **A** is strongly abelian, if for all polynomials  $t$  we have  
 $(\forall \mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}, \mathbf{e}) \quad t(\mathbf{a}, \mathbf{c}) = t(\mathbf{b}, \mathbf{d}) \implies t(\mathbf{e}, \mathbf{c}) = t(\mathbf{e}, \mathbf{d}).$

Let **A** be a nontrivial finite algebra and let **B** be a nontrivial homomorphic image of  $\mathbf{A}^k$  for some  $k$ .

- If **B** is strongly abelian, then  $d_{\mathbf{A}}(n) = 2^{\Theta(n)}$  (exponential).
- If **B** is abelian, then  $d_{\mathbf{A}}(n) = \Omega(n)$  (at least linear).

This holds, because the free algebras in the first case have polynomially bounded size, and in the second case their size is in  $2^{O(n)}$  by a result of J. Berman and R. McKenzie.

Each simple factoralgebra of a finite solvable algebra **A** is either abelian or strongly abelian,

# Strongly abelian algebras

An algebra **A** is strongly abelian, if for all polynomials  $t$  we have

$$(\forall \mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}, \mathbf{e}) \quad t(\mathbf{a}, \mathbf{c}) = t(\mathbf{b}, \mathbf{d}) \implies t(\mathbf{e}, \mathbf{c}) = t(\mathbf{e}, \mathbf{d}).$$

Let **A** be a nontrivial finite algebra and let **B** be a nontrivial homomorphic image of  $\mathbf{A}^k$  for some  $k$ .

- If **B** is strongly abelian, then  $d_{\mathbf{A}}(n) = 2^{\Theta(n)}$  (exponential).
- If **B** is abelian, then  $d_{\mathbf{A}}(n) = \Omega(n)$  (at least linear).

This holds, because the free algebras in the first case have polynomially bounded size, and in the second case their size is in  $2^{O(n)}$  by a result of J. Berman and R. McKenzie.

Each simple factoralgebra of a finite solvable algebra **A** is either abelian or strongly abelian, so  $d_{\mathbf{A}}(n)$  is at least linear.

# The hierarchy of abelianness properties

(1) **A** is solvable.

# The hierarchy of abelianness properties

- (1)  $\mathbf{A}$  is solvable.
- (2)  $\mathbf{A}$  is (left) nilpotent.

# The hierarchy of abelianness properties

- (1)  $\mathbf{A}$  is solvable.
- (2)  $\mathbf{A}$  is (left) nilpotent.
- (3)  $\mathbf{A}$  is abelian.

# The hierarchy of abelianness properties

- (1)  $\mathbf{A}$  is solvable.
- (2)  $\mathbf{A}$  is (left) nilpotent.
- (3)  $\mathbf{A}$  is abelian.
- (4)  $\mathbf{A}$  is a subdirect product of simple abelian algebras.

# The hierarchy of abelianness properties

- (1)  $\mathbf{A}$  is solvable.
- (2)  $\mathbf{A}$  is (left) nilpotent.
- (3)  $\mathbf{A}$  is abelian.
- (4)  $\mathbf{A}$  is a subdirect product of simple abelian algebras.
- (5)  $\mathbf{A}$  generates an abelian variety.



# The hierarchy of abelianness properties

- (1) **A** is solvable.
- (2) **A** is (left) nilpotent.
- (3) **A** is abelian.
- (4) **A** is a subdirect product of simple abelian algebras.
- (5) **A** generates an abelian variety.

We have  $(4) \Rightarrow (3) \Rightarrow (2) \Rightarrow (1)$

# The hierarchy of abelianness properties

- (1) **A** is solvable.
- (2) **A** is (left) nilpotent.
- (3) **A** is abelian.
- (4) **A** is a subdirect product of simple abelian algebras.
- (5) **A** generates an abelian variety.

We have  $(4) \Rightarrow (3) \Rightarrow (2) \Rightarrow (1)$  and  $(5) \Rightarrow (3)$ .

# The hierarchy of abelianness properties

- (1) **A** is solvable.
- (2) **A** is (left) nilpotent.
- (3) **A** is abelian.
- (4) **A** is a subdirect product of simple abelian algebras.
- (5) **A** generates an abelian variety.

We have  $(4) \Rightarrow (3) \Rightarrow (2) \Rightarrow (1)$  and  $(5) \Rightarrow (3)$ .

No other implications hold (except the formal consequences).

# The hierarchy of abelianness properties

- (1) **A** is solvable.
- (2) **A** is (left) nilpotent.
- (3) **A** is abelian.
- (4) **A** is a subdirect product of simple abelian algebras.
- (5) **A** generates an abelian variety.

We have  $(4) \Rightarrow (3) \Rightarrow (2) \Rightarrow (1)$  and  $(5) \Rightarrow (3)$ .

No other implications hold (except the formal consequences).

We prove that **stronger abelianness properties** yield

# The hierarchy of abelianness properties

- (1) **A** is solvable.
- (2) **A** is (left) nilpotent.
- (3) **A** is abelian.
- (4) **A** is a subdirect product of simple abelian algebras.
- (5) **A** generates an abelian variety.

We have  $(4) \Rightarrow (3) \Rightarrow (2) \Rightarrow (1)$  and  $(5) \Rightarrow (3)$ .

No other implications hold (except the formal consequences).

We prove that **stronger abelianness properties** yield  
a **closer relationship between various growth-restricting conditions**.

# The hierarchy of abelianness properties

- (1) **A** is solvable.
- (2) **A** is (left) nilpotent.
- (3) **A** is abelian.
- (4) **A** is a subdirect product of simple abelian algebras.
- (5) **A** generates an abelian variety.

We have  $(4) \Rightarrow (3) \Rightarrow (2) \Rightarrow (1)$  and  $(5) \Rightarrow (3)$ .

No other implications hold (except the formal consequences).

We prove that **stronger abelianness properties** yield a **closer relationship between various growth-restricting conditions**.

**Example:** Both (5) and (4) imply that the growth rate is non-exponential iff

# The hierarchy of abelianness properties

- (1) **A** is solvable.
- (2) **A** is (left) nilpotent.
- (3) **A** is abelian.
- (4) **A** is a subdirect product of simple abelian algebras.
- (5) **A** generates an abelian variety.

We have  $(4) \Rightarrow (3) \Rightarrow (2) \Rightarrow (1)$  and  $(5) \Rightarrow (3)$ .

No other implications hold (except the formal consequences).

We prove that **stronger abelianness properties** yield a **closer relationship between various growth-restricting conditions**.

**Example:** Both (5) and (4) imply that the growth rate is non-exponential iff **A** has a Maltsev term

# The hierarchy of abelianness properties

- (1) **A** is solvable.
- (2) **A** is (left) nilpotent.
- (3) **A** is abelian.
- (4) **A** is a subdirect product of simple abelian algebras.
- (5) **A** generates an abelian variety.

We have  $(4) \Rightarrow (3) \Rightarrow (2) \Rightarrow (1)$  and  $(5) \Rightarrow (3)$ .

No other implications hold (except the formal consequences).

We prove that **stronger abelianness properties** yield a **closer relationship between various growth-restricting conditions**.

**Example:** Both (5) and (4) imply that the growth rate is non-exponential iff **A** has a Maltsev term (in which case the growth rate is linear),



# The hierarchy of abelianness properties

- (1) **A** is solvable.
- (2) **A** is (left) nilpotent.
- (3) **A** is abelian.
- (4) **A** is a subdirect product of simple abelian algebras.
- (5) **A** generates an abelian variety.

We have  $(4) \Rightarrow (3) \Rightarrow (2) \Rightarrow (1)$  and  $(5) \Rightarrow (3)$ .

No other implications hold (except the formal consequences).

We prove that **stronger abelianness properties** yield a **closer relationship between various growth-restricting conditions**.

**Example:** Both (5) and (4) imply that the growth rate is non-exponential iff **A** has a Maltsev term (in which case the growth rate is linear), but (2) does not.

# The hierarchy of growth-restricting conditions

(i) **A** has a Maltsev polynomial.

# The hierarchy of growth-restricting conditions

- (i) **A** has a Maltsev polynomial.
- (ii) **A** has a pointed cube polynomial.

# The hierarchy of growth-restricting conditions

- (i) **A** has a Maltsev polynomial.
- (ii) **A** has a pointed cube polynomial.
- (iii) **A** is a spread of its type **2** minimal sets (see later).

# The hierarchy of growth-restricting conditions

- (i) **A** has a Maltsev polynomial.
- (ii) **A** has a pointed cube polynomial.
- (iii) **A** is a spread of its type **2** minimal sets (see later).
- (iv)  $d_{\mathbf{A}}(n) \in O(n)$ .

# The hierarchy of growth-restricting conditions

- (i) **A** has a Maltsev polynomial.
- (ii) **A** has a pointed cube polynomial.
- (iii) **A** is a spread of its type **2** minimal sets (see later).
- (iv)  $d_{\mathbf{A}}(n) \in O(n)$ .
- (v)  $d_{\mathbf{A}}(n) \notin 2^{\Omega(n)}$ .

# The hierarchy of growth-restricting conditions

- (i)  $\mathbf{A}$  has a Maltsev polynomial.
- (ii)  $\mathbf{A}$  has a pointed cube polynomial.
- (iii)  $\mathbf{A}$  is a spread of its type **2** minimal sets (see later).
- (iv)  $d_{\mathbf{A}}(n) \in O(n)$ .
- (v)  $d_{\mathbf{A}}(n) \notin 2^{\Omega(n)}$ .
- (vi)  $\mathbf{A}^n$  has no nontrivial strongly abelian factor (for all  $n$ ).

# The hierarchy of growth-restricting conditions

- (i)  $\mathbf{A}$  has a Maltsev polynomial.
- (ii)  $\mathbf{A}$  has a pointed cube polynomial.
- (iii)  $\mathbf{A}$  is a spread of its type  $\mathbf{2}$  minimal sets (see later).
- (iv)  $d_{\mathbf{A}}(n) \in O(n)$ .
- (v)  $d_{\mathbf{A}}(n) \notin 2^{\Omega(n)}$ .
- (vi)  $\mathbf{A}^n$  has no nontrivial strongly abelian factor (for all  $n$ ).

We have  $(i) \Rightarrow (iv)$ ,



# The hierarchy of growth-restricting conditions

- (i)  $\mathbf{A}$  has a Maltsev polynomial.
- (ii)  $\mathbf{A}$  has a pointed cube polynomial.
- (iii)  $\mathbf{A}$  is a spread of its type  $\mathbf{2}$  minimal sets (see later).
- (iv)  $d_{\mathbf{A}}(n) \in O(n)$ .
- (v)  $d_{\mathbf{A}}(n) \notin 2^{\Omega(n)}$ .
- (vi)  $\mathbf{A}^n$  has no nontrivial strongly abelian factor (for all  $n$ ).

We have  $(i) \Rightarrow (iv)$ ,  $(i) \Rightarrow (ii) \Rightarrow (v)$ ,

# The hierarchy of growth-restricting conditions

- (i)  $\mathbf{A}$  has a Maltsev polynomial.
- (ii)  $\mathbf{A}$  has a pointed cube polynomial.
- (iii)  $\mathbf{A}$  is a spread of its type  $\mathbf{2}$  minimal sets (see later).
- (iv)  $d_{\mathbf{A}}(n) \in O(n)$ .
- (v)  $d_{\mathbf{A}}(n) \notin 2^{\Omega(n)}$ .
- (vi)  $\mathbf{A}^n$  has no nontrivial strongly abelian factor (for all  $n$ ).

We have  $(i) \Rightarrow (iv)$ ,  $(i) \Rightarrow (ii) \Rightarrow (v)$ , and  $(iii) \Rightarrow (iv) \Rightarrow (v) \Rightarrow (vi)$ .

# The hierarchy of growth-restricting conditions

- (i)  $\mathbf{A}$  has a Maltsev polynomial.
- (ii)  $\mathbf{A}$  has a pointed cube polynomial.
- (iii)  $\mathbf{A}$  is a spread of its type  $\mathbf{2}$  minimal sets (see later).
- (iv)  $d_{\mathbf{A}}(n) \in O(n)$ .
- (v)  $d_{\mathbf{A}}(n) \notin 2^{\Omega(n)}$ .
- (vi)  $\mathbf{A}^n$  has no nontrivial strongly abelian factor (for all  $n$ ).

We have  $(i) \Rightarrow (iv)$ ,  $(i) \Rightarrow (ii) \Rightarrow (v)$ , and  $(iii) \Rightarrow (iv) \Rightarrow (v) \Rightarrow (vi)$ .

No other implications hold for general finite algebras.

# The hierarchy of growth-restricting conditions

- (i)  $\mathbf{A}$  has a Maltsev polynomial.
- (ii)  $\mathbf{A}$  has a pointed cube polynomial.
- (iii)  $\mathbf{A}$  is a spread of its type  $\mathbf{2}$  minimal sets (see later).
- (iv)  $d_{\mathbf{A}}(n) \in O(n)$ .
- (v)  $d_{\mathbf{A}}(n) \notin 2^{\Omega(n)}$ .
- (vi)  $\mathbf{A}^n$  has no nontrivial strongly abelian factor (for all  $n$ ).

We have (i) $\Rightarrow$ (iv), (i) $\Rightarrow$ (ii) $\Rightarrow$ (v), and (iii) $\Rightarrow$ (iv) $\Rightarrow$ (v) $\Rightarrow$ (vi).

No other implications hold for general finite algebras.

## Open problem

Is the growth rate of each finite solvable  $\mathbf{A}$  linear or exponential?

# The hierarchy of growth-restricting conditions

- (i)  $\mathbf{A}$  has a Maltsev polynomial.
- (ii)  $\mathbf{A}$  has a pointed cube polynomial.
- (iii)  $\mathbf{A}$  is a spread of its type  $\mathbf{2}$  minimal sets (see later).
- (iv)  $d_{\mathbf{A}}(n) \in O(n)$ .
- (v)  $d_{\mathbf{A}}(n) \notin 2^{\Omega(n)}$ .
- (vi)  $\mathbf{A}^n$  has no nontrivial strongly abelian factor (for all  $n$ ).

We have (i) $\Rightarrow$ (iv), (i) $\Rightarrow$ (ii) $\Rightarrow$ (v), and (iii) $\Rightarrow$ (iv) $\Rightarrow$ (v) $\Rightarrow$ (vi).

No other implications hold for general finite algebras.

## Open problem

Is the growth rate of each finite solvable  $\mathbf{A}$  linear or exponential?

True if  $\mathbf{A}$  is nilpotent;

# The hierarchy of growth-restricting conditions

- (i)  $\mathbf{A}$  has a Maltsev polynomial.
- (ii)  $\mathbf{A}$  has a pointed cube polynomial.
- (iii)  $\mathbf{A}$  is a spread of its type **2** minimal sets (see later).
- (iv)  $d_{\mathbf{A}}(n) \in O(n)$ .
- (v)  $d_{\mathbf{A}}(n) \notin 2^{\Omega(n)}$ .
- (vi)  $\mathbf{A}^n$  has no nontrivial strongly abelian factor (for all  $n$ ).

We have (i) $\Rightarrow$ (iv), (i) $\Rightarrow$ (ii) $\Rightarrow$ (v), and (iii) $\Rightarrow$ (iv) $\Rightarrow$ (v) $\Rightarrow$ (vi).  
No other implications hold for general finite algebras.

## Open problem

Is the growth rate of each finite solvable  $\mathbf{A}$  linear or exponential?

True if  $\mathbf{A}$  is nilpotent; would follow from (vi) $\Rightarrow$ (iii) for solvable  $\mathbf{A}$ .

# Spreads

## Definition

Let  $\mathbf{A}$  be an algebra and  $\mathcal{U}$  a collection of subsets of  $A$ .

# Spreads

## Definition

Let  $\mathbf{A}$  be an algebra and  $\mathcal{U}$  a collection of subsets of  $A$ .

A subset  $S \subseteq A$  is a **spread** with respect to  $\mathcal{U}$  if there exists a polynomial  $p$  of  $\mathbf{A}$



# Spreads

## Definition

Let  $\mathbf{A}$  be an algebra and  $\mathcal{U}$  a collection of subsets of  $A$ .

A subset  $S \subseteq A$  is a **spread** with respect to  $\mathcal{U}$  if there exists a polynomial  $p$  of  $\mathbf{A}$  and (not necessarily distinct) elements  $U_1, \dots, U_k \in \mathcal{U}$

# Spreads

## Definition

Let  $\mathbf{A}$  be an algebra and  $\mathcal{U}$  a collection of subsets of  $A$ .

A subset  $S \subseteq A$  is a **spread** with respect to  $\mathcal{U}$  if there exists a polynomial  $p$  of  $\mathbf{A}$  and (not necessarily distinct) elements  $U_1, \dots, U_k \in \mathcal{U}$  such that  $p(U_1, \dots, U_k) = S$ .

# Spreads

## Definition

Let  $\mathbf{A}$  be an algebra and  $\mathcal{U}$  a collection of subsets of  $A$ .  
A subset  $S \subseteq A$  is a **spread** with respect to  $\mathcal{U}$  if there exists a polynomial  $p$  of  $\mathbf{A}$  and (not necessarily distinct) elements  $U_1, \dots, U_k \in \mathcal{U}$  such that  $p(U_1, \dots, U_k) = S$ .

## Claim

If a finite algebra  $\mathbf{A}$  is a spread of a family of subsets on which the induced algebras have Maltsev polynomials

# Spreads

## Definition

Let  $\mathbf{A}$  be an algebra and  $\mathcal{U}$  a collection of subsets of  $A$ .  
A subset  $S \subseteq A$  is a **spread** with respect to  $\mathcal{U}$  if there exists a polynomial  $p$  of  $\mathbf{A}$  and (not necessarily distinct) elements  $U_1, \dots, U_k \in \mathcal{U}$  such that  $p(U_1, \dots, U_k) = S$ .

## Claim

If a finite algebra  $\mathbf{A}$  is a spread of a family of subsets on which the induced algebras have Maltsev polynomials (like type 2 minimal sets),

# Spreads

## Definition

Let  $\mathbf{A}$  be an algebra and  $\mathcal{U}$  a collection of subsets of  $A$ .  
A subset  $S \subseteq A$  is a **spread** with respect to  $\mathcal{U}$  if there exists a polynomial  $p$  of  $\mathbf{A}$  and (not necessarily distinct) elements  $U_1, \dots, U_k \in \mathcal{U}$  such that  $p(U_1, \dots, U_k) = S$ .

## Claim

If a finite algebra  $\mathbf{A}$  is a spread of a family of subsets on which the induced algebras have Maltsev polynomials (like type 2 minimal sets), then the growth rate of  $\mathbf{A}$  is at most linear.

# Spreads

## Definition

Let  $\mathbf{A}$  be an algebra and  $\mathcal{U}$  a collection of subsets of  $A$ .  
A subset  $S \subseteq A$  is a **spread** with respect to  $\mathcal{U}$  if there exists a polynomial  $p$  of  $\mathbf{A}$  and (not necessarily distinct) elements  $U_1, \dots, U_k \in \mathcal{U}$  such that  $p(U_1, \dots, U_k) = S$ .

## Claim

If a finite algebra  $\mathbf{A}$  is a spread of a family of subsets on which the induced algebras have Maltsev polynomials (like type **2** minimal sets), then the growth rate of  $\mathbf{A}$  is at most linear.

## Theorem (KKSz)

If  $\mathbf{A}$  is a finite solvable algebra with a Maltsev polynomial, then  $\mathbf{A}$  is a spread of its type **2** minimal sets.

# Solvable algebras

## Theorem (KKSz)

Let  $\mathbf{A}$  be a finite solvable algebra that has a pointed cube term.

# Solvable algebras

## Theorem (KKSz)

Let  $\mathbf{A}$  be a finite solvable algebra that has a pointed cube term.  
Then  $d_{\mathbf{A}}(n) = \Theta(n)$ .



# Solvable algebras

## Theorem (KKSz)

Let  $\mathbf{A}$  be a finite solvable algebra that has a pointed cube term.  
Then  $d_{\mathbf{A}}(n) = \Theta(n)$ .

Tool used: a new characterization of solvability.

# Solvable algebras

## Theorem (KKSz)

Let  $\mathbf{A}$  be a finite solvable algebra that has a pointed cube term.  
Then  $d_{\mathbf{A}}(n) = \Theta(n)$ .

Tool used: a new characterization of solvability.

Let  $\mathbf{A}$  be an algebra and  $p$  an idempotent polynomial of  $\mathbf{A}$ .

# Solvable algebras

## Theorem (KKSz)

Let  $\mathbf{A}$  be a finite solvable algebra that has a pointed cube term.  
Then  $d_{\mathbf{A}}(n) = \Theta(n)$ .

Tool used: a new characterization of solvability.

Let  $\mathbf{A}$  be an algebra and  $p$  an idempotent polynomial of  $\mathbf{A}$ .  
The **translation-digraph**  $\text{Tr}(p)$  on  $\mathbf{A}$  has directed edges  
 $(c, c') = (p(c, c, \dots, c), p(c, \dots, c, d, c, \dots, c))$ ,

# Solvable algebras

## Theorem (KKSz)

Let  $\mathbf{A}$  be a finite solvable algebra that has a pointed cube term.  
Then  $d_{\mathbf{A}}(n) = \Theta(n)$ .

Tool used: a new characterization of solvability.

Let  $\mathbf{A}$  be an algebra and  $p$  an idempotent polynomial of  $\mathbf{A}$ .  
The **translation-digraph**  $\text{Tr}(p)$  on  $A$  has directed edges  
 $(c, c') = (p(c, c, \dots, c), p(c, \dots, c, d, c, \dots, c))$ , where  $c, d \in A$ .

# Solvable algebras

## Theorem (KKSz)

Let  $\mathbf{A}$  be a finite solvable algebra that has a pointed cube term.  
Then  $d_{\mathbf{A}}(n) = \Theta(n)$ .

Tool used: a new characterization of solvability.

Let  $\mathbf{A}$  be an algebra and  $p$  an idempotent polynomial of  $\mathbf{A}$ .  
The **translation-digraph**  $\text{Tr}(p)$  on  $\mathbf{A}$  has directed edges  
 $(c, c') = (p(c, c, \dots, c), p(c, \dots, c, d, c, \dots, c))$ , where  $c, d \in A$ .

## Theorem (KKSz)

A finite algebra  $\mathbf{A}$  is solvable if and only if for every neighborhood  $U$  of  $\mathbf{A}$ ,

# Solvable algebras

## Theorem (KKSz)

Let  $\mathbf{A}$  be a finite solvable algebra that has a pointed cube term.  
Then  $d_{\mathbf{A}}(n) = \Theta(n)$ .

Tool used: a new characterization of solvability.

Let  $\mathbf{A}$  be an algebra and  $p$  an idempotent polynomial of  $\mathbf{A}$ .  
The **translation-digraph**  $\text{Tr}(p)$  on  $\mathbf{A}$  has directed edges  
 $(c, c') = (p(c, c, \dots, c), p(c, \dots, c, d, c, \dots, c))$ , where  $c, d \in A$ .

## Theorem (KKSz)

A finite algebra  $\mathbf{A}$  is solvable if and only if for every neighborhood  $U$  of  $\mathbf{A}$ , and every idempotent polynomial  $p$  of the induced algebra  $\mathbf{A}|_U$ ,

# Solvable algebras

## Theorem (KKSz)

Let  $\mathbf{A}$  be a finite solvable algebra that has a pointed cube term.  
Then  $d_{\mathbf{A}}(n) = \Theta(n)$ .

Tool used: a new characterization of solvability.

Let  $\mathbf{A}$  be an algebra and  $p$  an idempotent polynomial of  $\mathbf{A}$ .  
The **translation-digraph**  $\mathbf{Tr}(p)$  on  $A$  has directed edges  
 $(c, c') = (p(c, c, \dots, c), p(c, \dots, c, d, c, \dots, c))$ , where  $c, d \in A$ .

## Theorem (KKSz)

A finite algebra  $\mathbf{A}$  is solvable if and only if for every neighborhood  $U$  of  $\mathbf{A}$ , and every idempotent polynomial  $p$  of the induced algebra  $\mathbf{A}|_U$ , the directed graph  $\mathbf{Tr}(p)$  is strongly connected.

# Nilpotent algebras

## Theorem (KKSz)

A finite left nilpotent algebra has a Maltsev polynomial iff it has a pointed cube polynomial.



# Nilpotent algebras

## Theorem (KKSz)

A finite left nilpotent algebra has a Maltsev polynomial iff it has a pointed cube polynomial. Hence a finite abelian algebra has a pointed cube polynomial iff it is affine

# Nilpotent algebras

## Theorem (KKSz)

A finite left nilpotent algebra has a Maltsev polynomial iff it has a pointed cube polynomial. Hence a finite abelian algebra has a pointed cube polynomial iff it is affine (so has a Maltsev-term).

# Nilpotent algebras

## Theorem (KKSz)

A finite left nilpotent algebra has a Maltsev polynomial iff it has a pointed cube polynomial. Hence a finite abelian algebra has a pointed cube polynomial iff it is affine (so has a Maltsev-term).

## Theorem (KKSz)

If  $\mathbf{A}$  is a finite, left nilpotent algebra, and  $\mathbf{A}^{|\mathbf{A}|}$  does not have a nontrivial strongly abelian quotient algebra,

# Nilpotent algebras

## Theorem (KKSz)

A finite left nilpotent algebra has a Maltsev polynomial iff it has a pointed cube polynomial. Hence a finite abelian algebra has a pointed cube polynomial iff it is affine (so has a Maltsev-term).

## Theorem (KKSz)

If  $\mathbf{A}$  is a finite, left nilpotent algebra, and  $\mathbf{A}^{|\mathbf{A}|}$  does not have a nontrivial strongly abelian quotient algebra, then  $\mathbf{A}$  is a spread of its type 2 minimal sets

# Nilpotent algebras

## Theorem (KKSz)

A finite left nilpotent algebra has a Maltsev polynomial iff it has a pointed cube polynomial. Hence a finite abelian algebra has a pointed cube polynomial iff it is affine (so has a Maltsev-term).

## Theorem (KKSz)

If  $\mathbf{A}$  is a finite, left nilpotent algebra, and  $\mathbf{A}^{|\mathbf{A}|}$  does not have a nontrivial strongly abelian quotient algebra, then  $\mathbf{A}$  is a spread of its type 2 minimal sets (hence linear).

# Nilpotent algebras

## Theorem (KKSz)

A finite left nilpotent algebra has a Maltsev polynomial iff it has a pointed cube polynomial. Hence a finite abelian algebra has a pointed cube polynomial iff it is affine (so has a Maltsev-term).

## Theorem (KKSz)

If  $\mathbf{A}$  is a finite, left nilpotent algebra, and  $\mathbf{A}^{|\mathbf{A}|}$  does not have a nontrivial strongly abelian quotient algebra, then  $\mathbf{A}$  is a spread of its type 2 minimal sets (hence linear).

The proof uses the quasi-Hamiltonian property for the subalgebras of  $\mathbf{A}^{|\mathbf{A}|}$ .

# Abelian varieties

## Theorem (KKSz)

Let  $\mathbf{A}$  be an algebra, which is a spread of subsets whose induced algebras are affine. Then the following hold.

# Abelian varieties

## Theorem (KKSz)

Let  $\mathbf{A}$  be an algebra, which is a spread of subsets whose induced algebras are affine. Then the following hold.

- If  $H(\mathbf{A}^2)$  is abelian, then there is an abelian group operation on  $\mathbf{A}$  that is compatible with all operations of  $\mathbf{A}$ ,



# Abelian varieties

## Theorem (KKSz)

Let  $\mathbf{A}$  be an algebra, which is a spread of subsets whose induced algebras are affine. Then the following hold.

- If  $H(\mathbf{A}^2)$  is abelian, then there is an abelian group operation on  $A$  that is compatible with all operations of  $\mathbf{A}$ , and preserves all congruences of  $\mathbf{A}$ .

# Abelian varieties

## Theorem (KKSz)

Let  $\mathbf{A}$  be an algebra, which is a spread of subsets whose induced algebras are affine. Then the following hold.

- If  $H(\mathbf{A}^2)$  is abelian, then there is an abelian group operation on  $\mathbf{A}$  that is compatible with all operations of  $\mathbf{A}$ , and preserves all congruences of  $\mathbf{A}$ .
- If the variety  $V(\mathbf{A})$  generated by  $\mathbf{A}$  is abelian, then  $\mathbf{A}$  is affine.

# Abelian varieties

## Theorem (KKSz)

Let  $\mathbf{A}$  be an algebra, which is a spread of subsets whose induced algebras are affine. Then the following hold.

- If  $H(\mathbf{A}^2)$  is abelian, then there is an abelian group operation on  $\mathbf{A}$  that is compatible with all operations of  $\mathbf{A}$ , and preserves all congruences of  $\mathbf{A}$ .
- If the variety  $V(\mathbf{A})$  generated by  $\mathbf{A}$  is abelian, then  $\mathbf{A}$  is affine.

## Examples

An 8-element quasi-affine algebra shows that in the second statement the assumption that  $V(\mathbf{A})$  is abelian cannot be dropped.

# Abelian varieties

## Theorem (KKSz)

Let  $\mathbf{A}$  be an algebra, which is a spread of subsets whose induced algebras are affine. Then the following hold.

- If  $H(\mathbf{A}^2)$  is abelian, then there is an abelian group operation on  $\mathbf{A}$  that is compatible with all operations of  $\mathbf{A}$ , and preserves all congruences of  $\mathbf{A}$ .
- If the variety  $V(\mathbf{A})$  generated by  $\mathbf{A}$  is abelian, then  $\mathbf{A}$  is affine.

## Examples

An 8-element quasi-affine algebra shows that in the second statement the assumption that  $V(\mathbf{A})$  is abelian cannot be dropped. Another 8-element abelian algebra shows that in the first statement it is not sufficient to assume only that  $H(\mathbf{A})$  is abelian.

# Semisimple algebras

## Theorem (KKSz)

Let  $\mathbf{A}$  be a finite solvable algebra and  $\beta$  the intersection of all maximal congruences of  $\mathbf{A}$ .

# Semisimple algebras

## Theorem (KKSz)

Let  $\mathbf{A}$  be a finite solvable algebra and  $\beta$  the intersection of all maximal congruences of  $\mathbf{A}$ . If the growth rate of  $\mathbf{A}/\beta$  is linear, then  $\mathbf{A}/\beta$  has a Maltsev polynomial.

# Semisimple algebras

## Theorem (KKSz)

Let  $\mathbf{A}$  be a finite solvable algebra and  $\beta$  the intersection of all maximal congruences of  $\mathbf{A}$ . If the growth rate of  $\mathbf{A}/\beta$  is linear, then  $\mathbf{A}/\beta$  has a Maltsev polynomial. In particular, if  $\mathbf{A}$  is (linear, and) a direct product of simple abelian algebras, then  $\mathbf{A}$  is Maltsev.

# Semisimple algebras

## Theorem (KKSz)

Let  $\mathbf{A}$  be a finite solvable algebra and  $\beta$  the intersection of all maximal congruences of  $\mathbf{A}$ . If the growth rate of  $\mathbf{A}/\beta$  is linear, then  $\mathbf{A}/\beta$  has a Maltsev polynomial. In particular, if  $\mathbf{A}$  is (linear, and) a direct product of simple abelian algebras, then  $\mathbf{A}$  is Maltsev.

The proof shows that  $\mathbf{A}/\beta$  is a direct product, and not just a subdirect product of simple abelian algebras.



# Semisimple algebras

## Theorem (KKSz)

Let  $\mathbf{A}$  be a finite solvable algebra and  $\beta$  the intersection of all maximal congruences of  $\mathbf{A}$ . If the growth rate of  $\mathbf{A}/\beta$  is linear, then  $\mathbf{A}/\beta$  has a Maltsev polynomial. In particular, if  $\mathbf{A}$  is (linear, and) a direct product of simple abelian algebras, then  $\mathbf{A}$  is Maltsev.

The proof shows that  $\mathbf{A}/\beta$  is a direct product, and not just a subdirect product of simple abelian algebras.

## Example

There exist a 16-element algebra that is a direct product of two, 4-element affine (hence abelian, Maltsev) algebras,

# Semisimple algebras

## Theorem (KKSz)

Let  $\mathbf{A}$  be a finite solvable algebra and  $\beta$  the intersection of all maximal congruences of  $\mathbf{A}$ . If the growth rate of  $\mathbf{A}/\beta$  is linear, then  $\mathbf{A}/\beta$  has a Maltsev polynomial. In particular, if  $\mathbf{A}$  is (linear, and) a direct product of simple abelian algebras, then  $\mathbf{A}$  is Maltsev.

The proof shows that  $\mathbf{A}/\beta$  is a direct product, and not just a subdirect product of simple abelian algebras.

## Example

There exist a 16-element algebra that is a direct product of two, 4-element affine (hence abelian, Maltsev) algebras, has a linear growth rate,

# Semisimple algebras

## Theorem (KKSz)

Let  $\mathbf{A}$  be a finite solvable algebra and  $\beta$  the intersection of all maximal congruences of  $\mathbf{A}$ . If the growth rate of  $\mathbf{A}/\beta$  is linear, then  $\mathbf{A}/\beta$  has a Maltsev polynomial. In particular, if  $\mathbf{A}$  is (linear, and) a direct product of simple abelian algebras, then  $\mathbf{A}$  is Maltsev.

The proof shows that  $\mathbf{A}/\beta$  is a direct product, and not just a subdirect product of simple abelian algebras.

## Example

There exist a 16-element algebra that is a direct product of two, 4-element affine (hence abelian, Maltsev) algebras, has a linear growth rate, but does not have a Maltsev polynomial.

# Summary: arbitrary

- (i)  $\mathbf{A}$  has a Maltsev polynomial.
- (ii)  $\mathbf{A}$  has a pointed cube polynomial.
- (iii)  $\mathbf{A}$  is a spread of its type 2 minimal sets.
- (iv)  $d_{\mathbf{A}}(n) \in O(n)$ .
- (v)  $d_{\mathbf{A}}(n) \notin 2^{\Omega(n)}$ .
- (vi)  $\mathbf{A}^n$  has no nontrivial strongly abelian factor (for all  $n$ ).

All are equivalent if  $\mathbf{A}$  is semisimple or if  $\mathbf{V}(\mathbf{A})$  is abelian.

$$(i) \implies (ii)$$



$$(iii) \implies (iv) \implies (v) \implies (vi).$$

For arbitrary finite algebras

# Summary: solvable

- (i)  $\mathbf{A}$  has a Maltsev polynomial.
- (ii)  $\mathbf{A}$  has a pointed cube polynomial.
- (iii)  $\mathbf{A}$  is a spread of its type 2 minimal sets.
- (iv)  $d_{\mathbf{A}}(n) \in O(n)$ .
- (v)  $d_{\mathbf{A}}(n) \notin 2^{\Omega(n)}$ .
- (vi)  $\mathbf{A}^n$  has no nontrivial strongly abelian factor (for all  $n$ ).

All are equivalent if  $\mathbf{A}$  is semisimple or if  $\mathbf{V}(\mathbf{A})$  is abelian.

$$(i) \implies (ii)$$



$$(iii) \implies (iv) \implies (v) \implies (vi).$$

For finite, solvable algebras

# Summary: nilpotent

- (i)  $\mathbf{A}$  has a Maltsev polynomial.
- (ii)  $\mathbf{A}$  has a pointed cube polynomial.
- (iii)  $\mathbf{A}$  is a spread of its type 2 minimal sets.
- (iv)  $d_{\mathbf{A}}(n) \in O(n)$ .
- (v)  $d_{\mathbf{A}}(n) \notin 2^{\Omega(n)}$ .
- (vi)  $\mathbf{A}^n$  has no nontrivial strongly abelian factor (for all  $n$ ).

All are equivalent if  $\mathbf{A}$  is semisimple or if  $\mathbf{V}(\mathbf{A})$  is abelian.

$$(i) \iff (ii)$$



$$(iii) \iff (iv) \iff (v) \iff (vi).$$

For finite, left nilpotent algebras

# Open problems

Is there a finite algebra  $\mathbf{A}$  such that  $d_{\mathbf{A}}(n) \notin \Omega(n)$  and  $d_{\mathbf{A}}(n) \notin O(\log(n))$ ? That is, whose growth rate is between logarithmic and linear?

# Open problems

Is there a finite algebra  $\mathbf{A}$  such that  $d_{\mathbf{A}}(n) \notin \Omega(n)$  and  $d_{\mathbf{A}}(n) \notin O(\log(n))$ ? That is, whose growth rate is between logarithmic and linear? Open for 2-element partial algebras, too.



# Open problems

Is there a finite algebra  $\mathbf{A}$  such that  $d_{\mathbf{A}}(n) \notin \Omega(n)$  and  $d_{\mathbf{A}}(n) \notin O(\log(n))$ ? That is, whose growth rate is between logarithmic and linear? Open for 2-element partial algebras, too.

Is it true that a finite algebra with a 2-sided unit for some binary term has logarithmic or linear growth?

# Open problems

Is there a finite algebra  $\mathbf{A}$  such that  $d_{\mathbf{A}}(n) \notin \Omega(n)$  and  $d_{\mathbf{A}}(n) \notin O(\log(n))$ ? That is, whose growth rate is between logarithmic and linear? Open for 2-element partial algebras, too.

Is it true that a finite algebra with a 2-sided unit for some binary term has logarithmic or linear growth? (Note that the identities  $x * 1 = 1 * x = x$  show that  $*$  is a 1-pointed 2-cube term.)

# Open problems

Is there a finite algebra  $\mathbf{A}$  such that  $d_{\mathbf{A}}(n) \notin \Omega(n)$  and  $d_{\mathbf{A}}(n) \notin O(\log(n))$ ? That is, whose growth rate is between logarithmic and linear? Open for 2-element partial algebras, too.

Is it true that a finite algebra with a 2-sided unit for some binary term has logarithmic or linear growth? (Note that the identities  $x * 1 = 1 * x = x$  show that  $*$  is a 1-pointed 2-cube term.)

Does (ii) $\Rightarrow$ (iii) hold for finite solvable algebras?

# Open problems

Is there a finite algebra  $\mathbf{A}$  such that  $d_{\mathbf{A}}(n) \notin \Omega(n)$  and  $d_{\mathbf{A}}(n) \notin O(\log(n))$ ? That is, whose growth rate is between logarithmic and linear? Open for 2-element partial algebras, too.

Is it true that a finite algebra with a 2-sided unit for some binary term has logarithmic or linear growth? (Note that the identities  $x * 1 = 1 * x = x$  show that  $*$  is a 1-pointed 2-cube term.)

Does  $(ii) \Rightarrow (iii)$  hold for finite solvable algebras?

Which of the true implications  $(iii) \Rightarrow (iv) \Rightarrow (v) \Rightarrow (vi)$  can be reversed for finite solvable algebras?

# Open problems

Is there a finite algebra  $\mathbf{A}$  such that  $d_{\mathbf{A}}(n) \notin \Omega(n)$  and  $d_{\mathbf{A}}(n) \notin O(\log(n))$ ? That is, whose growth rate is between logarithmic and linear? Open for 2-element partial algebras, too.

Is it true that a finite algebra with a 2-sided unit for some binary term has logarithmic or linear growth? (Note that the identities  $x * 1 = 1 * x = x$  show that  $*$  is a 1-pointed 2-cube term.)

Does  $(ii) \Rightarrow (iii)$  hold for finite solvable algebras?

Which of the true implications  $(iii) \Rightarrow (iv) \Rightarrow (v) \Rightarrow (vi)$  can be reversed for finite solvable algebras? In particular, is the growth rate of a finite solvable algebra always linear or exponential?

# Literature

- Hobby, David, McKenzie, Ralph, *The structure of finite algebras*. Contemporary Mathematics, **76**. American Mathematical Society, Providence, RI, 1988.

# Literature

- Hobby, David, McKenzie, Ralph, *The structure of finite algebras*. Contemporary Mathematics, **76**. American Mathematical Society, Providence, RI, 1988.
- Kearnes, Keith, *An order-theoretic property of the commutator*. Internat. J. Algebra Comput. **3** (1993), 491–533.

# Literature

- Hobby, David, McKenzie, Ralph, *The structure of finite algebras*. Contemporary Mathematics, **76**. American Mathematical Society, Providence, RI, 1988.
- Kearnes, Keith, *An order-theoretic property of the commutator*. Internat. J. Algebra Comput. **3** (1993), 491–533.
- Kearnes, Keith, *A Hamiltonian property for nilpotent algebras*. Algebra Universalis **37** (1997), 403–421.



# Literature

- Hobby, David, McKenzie, Ralph, *The structure of finite algebras*. Contemporary Mathematics, **76**. American Mathematical Society, Providence, RI, 1988.
- Kearnes, Keith, *An order-theoretic property of the commutator*. Internat. J. Algebra Comput. **3** (1993), 491–533.
- Kearnes, Keith, *A Hamiltonian property for nilpotent algebras*. Algebra Universalis **37** (1997), 403–421.
- KKSz: *Growth rates of algebras I: pointed cube terms*.  
[Arxiv](#): 1311.2352

# Literature

- Hobby, David, McKenzie, Ralph, *The structure of finite algebras*. Contemporary Mathematics, **76**. American Mathematical Society, Providence, RI, 1988.
- Kearnes, Keith, *An order-theoretic property of the commutator*. Internat. J. Algebra Comput. **3** (1993), 491–533.
- Kearnes, Keith, *A Hamiltonian property for nilpotent algebras*. Algebra Universalis **37** (1997), 403–421.
- KKSz: *Growth rates of algebras I: pointed cube terms*. [Arxiv](#): 1311.2352
- KKSz: *Growth rates of algebras II: Wiegold dichotomy*. [Arxiv](#): 1311.6189

# Literature

- Hobby, David, McKenzie, Ralph, *The structure of finite algebras*. Contemporary Mathematics, **76**. American Mathematical Society, Providence, RI, 1988.
- Kearnes, Keith, *An order-theoretic property of the commutator*. Internat. J. Algebra Comput. **3** (1993), 491–533.
- Kearnes, Keith, *A Hamiltonian property for nilpotent algebras*. Algebra Universalis **37** (1997), 403–421.
- KKSz: *Growth rates of algebras I: pointed cube terms*. [Arxiv: 1311.2352](#)
- KKSz: *Growth rates of algebras II: Wiegold dichotomy*. [Arxiv: 1311.6189](#)
- KKSz: *Growth rates of algebras III: Solvable algebras*. [Arxiv: 1311.2359](#)

# Eötvös University, Faculty of Natural Sciences



Thank you for your attention.