

Reconstructing the topology of the polymorphism clone of the random graph

András Pongrácz

Laboratoire d'Informatique, École Polytechnique

joint work with Manuel Bodirsky and Michael Pinsker

Prague, 2014

The general problem

The general problem

Δ : a countable relational structure (typically ω -categorical)

The general problem

Δ : a countable relational structure (typically ω -categorical)

$\mathcal{C} = \text{Pol}(\Delta)$, closed clone

The general problem

Δ : a countable relational structure (typically ω -categorical)

$\mathcal{C} = \text{Pol}(\Delta)$, closed clone

Question 1 (Up to ...) can we reconstruct Δ from $\text{Pol}(\Delta)$ viewed as an abstract clone?

The general problem

Δ : a countable relational structure (typically ω -categorical)

$\mathcal{C} = \text{Pol}(\Delta)$, closed clone

Question 1 (Up to ...) can we reconstruct Δ from $\text{Pol}(\Delta)$ viewed as an **abstract clone**? (same for groups and monoids)

The general problem

Δ : a countable relational structure (typically ω -categorical)

$\mathcal{C} = \text{Pol}(\Delta)$, closed clone

Question 1 (Up to ...) can we reconstruct Δ from $\text{Pol}(\Delta)$ viewed as an **abstract clone**? (same for groups and monoids)

Question 2 Can we reconstruct the topology on $\text{Pol}(\Delta)$ from the algebraic structure of $\text{Pol}(\Delta)$?

The general problem

Δ : a countable relational structure (typically ω -categorical)

$\mathcal{C} = \text{Pol}(\Delta)$, closed clone

Question 1 (Up to ...) can we reconstruct Δ from $\text{Pol}(\Delta)$ viewed as an **abstract clone**? (same for groups and monoids)

Question 2 Can we reconstruct the topology on $\text{Pol}(\Delta)$ from the algebraic structure of $\text{Pol}(\Delta)$? (same for groups and monoids)

Theorem [C. Ryll-Nardzewski]

Δ is ω -categorical iff $\text{Aut}(\Delta)$ is oligomorphic.

Theorem [C. Ryll-Nardzewski]

Δ is ω -categorical iff $\text{Aut}(\Delta)$ is oligomorphic.

Moreover, Δ and Γ are first-order interdefinable iff $\text{Aut}(\Delta) = \text{Aut}(\Gamma)$.

Theorem [C. Ryll-Nardzewski]

Δ is ω -categorical iff $\text{Aut}(\Delta)$ is oligomorphic.

Moreover, Δ and Γ are first-order interdefinable iff $\text{Aut}(\Delta) = \text{Aut}(\Gamma)$.

Theorem [G. Ahlbrandt, M. Ziegler]

If Δ is ω -categorical, then Δ and Γ are first-order bi-interpretable iff $\text{Aut}(\Delta) \cong \text{Aut}(\Gamma)$ as topological groups.

$\text{Aut}(\Delta)$

Permutation group \leftrightarrow first-order interdefinability

Topological group \leftrightarrow first-order bi-interpretability

Abstract group \leftrightarrow ?

$\text{Aut}(\Delta)$

Permutation group \leftrightarrow first-order interdefinability

Topological group \leftrightarrow first-order bi-interpretability

Abstract group \leftrightarrow The same?

Groups

Δ : a countable structure (typically ω -categorical).

Groups

Δ : a countable structure (typically ω -categorical).

Automatic homeomorphicity

For any closed group G and any isomorphism $\xi : \text{Aut}(\Delta) \rightarrow G$, ξ is a homeomorphism.

Groups

Δ : a countable structure (typically ω -categorical).

Automatic homeomorphicity

For any closed group G and any isomorphism $\xi : \text{Aut}(\Delta) \rightarrow G$, ξ is a homeomorphism.

Reconstruction

For any closed group G , if there exists an isomorphism $\xi : \text{Aut}(\Delta) \rightarrow G$, then there exists (possibly another) isomorphism $\xi' : \text{Aut}(\Delta) \rightarrow G$ which is a homeomorphism.

Groups

The following groups have automatic homeomorphicity.

The following groups have automatic homeomorphicity.

[E. B. Rabinovich (1977)] S_∞

Groups

The following groups have automatic homeomorphicity.

[E. B. Rabinovich (1977)] S_∞

[D. M. Evans (1986)] $\text{Aut}(V_\infty)$, where $V_\infty = GF(q)^\omega$ for some finite q .

Groups

The following groups have automatic homeomorphicity.

[E. B. Rabinovich (1977)] S_∞

[D. M. Evans (1986)] $\text{Aut}(V_\infty)$, where $V_\infty = GF(q)^\omega$ for some finite q .

[J. Truss (1989)] $\text{Aut}(\mathbb{Q}, <)$

Groups

The following groups have automatic homeomorphicity.

[E. B. Rabinovich (1977)] S_∞

[D. M. Evans (1986)] $\text{Aut}(V_\infty)$, where $V_\infty = GF(q)^\omega$ for some finite q .

[J. Truss (1989)] $\text{Aut}(\mathbb{Q}, <)$

[W. Hodges, I. Hodkinson, D. Lascar, S. Shelah (1993)] $\text{Aut}(V, E)$

Groups

The following groups have automatic homeomorphicity.

[E. B. Rabinovich (1977)] S_∞

[D. M. Evans (1986)] $\text{Aut}(V_\infty)$, where $V_\infty = GF(q)^\omega$ for some finite q .

[J. Truss (1989)] $\text{Aut}(\mathbb{Q}, <)$

[W. Hodges, I. Hodkinson, D. Lascar, S. Shelah (1993)] $\text{Aut}(V, E)$

[M. Rubin (1994)] Generic poset, universal tournament, etc.

Groups

The following groups have automatic homeomorphicity.

[E. B. Rabinovich (1977)] S_∞

[D. M. Evans (1986)] $\text{Aut}(V_\infty)$, where $V_\infty = GF(q)^\omega$ for some finite q .

[J. Truss (1989)] $\text{Aut}(\mathbb{Q}, <)$

[W. Hodges, I. Hodkinson, D. Lascar, S. Shelah (1993)] $\text{Aut}(V, E)$

[M. Rubin (1994)] Generic poset, universal tournament, etc.

[S. Barbina, D. Macpherson (2004)] Random k -uniform hypergraphs, Henson digraphs, etc.

Groups

The following groups have automatic homeomorphicity.

[E. B. Rabinovich (1977)] S_∞

[D. M. Evans (1986)] $\text{Aut}(V_\infty)$, where $V_\infty = GF(q)^\omega$ for some finite q .

[J. Truss (1989)] $\text{Aut}(\mathbb{Q}, <)$

[W. Hodges, I. Hodkinson, D. Lascar, S. Shelah (1993)] $\text{Aut}(V, E)$

[M. Rubin (1994)] Generic poset, universal tournament, etc.

[S. Barbina, D. Macpherson (2004)] Random k -uniform hypergraphs, Henson digraphs, etc.

Fact: It is consistent with ZF that for every ω -categorical structure Δ the topological group $\text{Aut}(\Delta)$ has automatic homeomorphicity.

$\text{Aut}(\Delta)$

Permutation group \leftrightarrow first-order interdefinability

Topological group \leftrightarrow first-order bi-interpretability

Abstract group \leftrightarrow The same?

$\text{Pol}(\Delta)$

Function clone \leftrightarrow primitive positive interdefinability

Topological clone \leftrightarrow primitive positive bi-interpretability

Abstract clone \leftrightarrow The same?

$\text{End}(\Delta)$

Transf. monoid \leftrightarrow existential positive interdefinability

Topo. monoid \leftrightarrow existential positive bi-interpretability

Abstract monoid \leftrightarrow The same?

$\text{Aut}(\Delta)$

Permutation group \leftrightarrow first-order interdefinability

Topological group \leftrightarrow first-order bi-interpretability

Abstract group \leftrightarrow The same?

Pol(Δ)

M. Bodirsky, J. Nesetril

Function clone \leftrightarrow primitive positive interdefinability

M. Bodirsky, M. Pinsker

Topological clone \leftrightarrow primitive positive bi-interpretability

Abstract clone \leftrightarrow The same?

End(Δ)

M. Bodirsky, M. Pinsker

Transf. monoid \leftrightarrow existential positive interdefinability

M. Bodirsky, M. Junker

Topo. monoid \leftrightarrow existential positive bi-interpretability

Abstract monoid \leftrightarrow The same?

Aut(Δ)

C. Ryll-Nardzewski

Permutation group \leftrightarrow first-order interdefinability

G. Ahlbrandt, M. Ziegler

Topological group \leftrightarrow first-order bi-interpretability

Abstract group \leftrightarrow The same?

Pol(Δ)

M. Bodirsky, J. Nesetril

Function clone \leftrightarrow primitive positive interdefinability

M. Bodirsky, M. Pinsker

Topological clone \leftrightarrow primitive positive bi-interpretability

Abstract clone \leftrightarrow The same?

End(Δ)

M. Bodirsky, M. Pinsker

Transf. monoid \leftrightarrow existential positive interdefinability

M. Bodirsky, M. Junker

Topo. monoid \leftrightarrow existential positive bi-interpretability

Abstract monoid \leftrightarrow The same?

Aut(Δ)

C. Ryll-Nardzewski

Permutation group \leftrightarrow first-order interdefinability

G. Ahlbrandt, M. Ziegler

Topological group \leftrightarrow first-order bi-interpretability

Abstract group \leftrightarrow The same?

Clones

Δ : a countable structure (typically ω -categorical).

Automatic homeomorphicity

For any closed clone \mathcal{C} and any isomorphism $\xi : \text{Pol}(\Delta) \rightarrow \mathcal{C}$, ξ is a homeomorphism.

Reconstruction

For any closed clone \mathcal{C} , if there exists an isomorphism $\xi : \text{Pol}(\Delta) \rightarrow \mathcal{C}$, then there exists (possibly another) isomorphism $\xi' : \text{Pol}(\Delta) \rightarrow \mathcal{C}$ which is a homeomorphism.

Pol(Δ)

M. Bodirsky, J. Nešetřil

Function clone \leftrightarrow primitive positive interdefinability

M. Bodirsky, M. Pinsker

Topological clone \leftrightarrow primitive positive bi-interpretability

Abstract clone \leftrightarrow The same?

End(Δ)

M. Bodirsky, M. Pinsker

Transf. monoid \leftrightarrow existential positive interdefinability

M. Bodirsky, M. Junker

Topo. monoid \leftrightarrow existential positive bi-interpretability

Abstract monoid \leftrightarrow The same?

Aut(Δ)

C. Ryll-Nardzewski

Permutation group \leftrightarrow first-order interdefinability

G. Ahlbrandt, M. Ziegler

Topological group \leftrightarrow first-order bi-interpretability

Abstract group \leftrightarrow The same?

From groups to monoids

\mathcal{O} : the clone of all operations

$\mathcal{O}^{(1)}$: its unary part

From groups to monoids

\mathcal{O} : the clone of all operations

$\mathcal{O}^{(1)}$: its unary part

Proposition

Let \mathcal{M} and \mathcal{M}' be closed submonoids of $\mathcal{O}^{(1)}$ with dense subsets of invertibles \mathcal{G} and \mathcal{G}' . Let $\xi : \mathcal{G} \rightarrow \mathcal{G}'$ be a continuous isomorphism. Then ξ extends to an isomorphism $\bar{\xi} : \mathcal{M} \rightarrow \mathcal{M}'$ which is a homeomorphism.

From groups to monoids

Proposition (M. Bodirsky, M. Pinsker, AP '13)

Let \mathcal{M} be a closed submonoid of $\mathcal{O}^{(1)}$ whose group of invertible elements \mathcal{G} is dense in \mathcal{M} and has automatic homeomorphicity. Assume that the only injective endomorphism of \mathcal{M} that fixes every element of \mathcal{G} is the identity function $\text{id}_{\mathcal{M}}$ on \mathcal{M} . Then \mathcal{M} has automatic homeomorphicity.

Automatic homeomorphicity of monoids

Theorem (M. Bodirsky, M. Pinsker, AP '13)

Let Δ be a countable homogeneous relational structure such that $\text{Aut}(\Delta)$ has no algebraicity and with the joint extension property such that $\text{Aut}(\Delta)$ has automatic homeomorphicity. Then the monoid $\overline{\text{Aut}(\Delta)}$ of self-embeddings of Δ has automatic homeomorphicity.

Automatic homeomorphicity of monoids

Theorem (M. Bodirsky, M. Pinsker, AP '13)

Let Δ be a countable homogeneous relational structure such that $\text{Aut}(\Delta)$ has no algebraicity and with the joint extension property such that $\text{Aut}(\Delta)$ has automatic homeomorphicity. Then the monoid $\overline{\text{Aut}(\Delta)}$ of self-embeddings of Δ has automatic homeomorphicity.

Examples: (V, E) , random tournament, random hypergraphs, $(\omega, =)$

Automatic homeomorphicity of monoids

Theorem (M. Bodirsky, M. Pinsker, AP '13)

Let Δ be a countable homogeneous relational structure such that $\text{Aut}(\Delta)$ has no algebraicity and with the joint extension property such that $\text{Aut}(\Delta)$ has automatic homeomorphicity. Then the monoid $\overline{\text{Aut}(\Delta)}$ of self-embeddings of Δ has automatic homeomorphicity.

Examples: (V, E) , random tournament, random hypergraphs, $(\omega, =)$

Non-examples: $(\mathbb{Q}, <)$, Henson graphs

Horn clone

Horn clone

Let \mathcal{H} be the closed clone generated by

Horn clone

Let \mathcal{H} be the closed clone generated by

- all unary injections, and

Horn clone

Let \mathcal{H} be the closed clone generated by

- all unary injections, and
- a binary injection $f: \omega^2 \rightarrow \omega$.

Horn clone

Let \mathcal{H} be the closed clone generated by

- all unary injections, and
- a binary injection $f: \omega^2 \rightarrow \omega$.

\mathcal{H} consists of all injections $\omega^n \rightarrow \omega$

Horn clone

Let \mathcal{H} be the closed clone generated by

- all unary injections, and
- a binary injection $f: \omega^2 \rightarrow \omega$.

\mathcal{H} consists of all injections $\omega^n \rightarrow \omega$ (almost. . .)

Horn clone

Let \mathcal{H} be the closed clone generated by

- all unary injections, and
- a binary injection $f: \omega^2 \rightarrow \omega$.

\mathcal{H} consists of all injections $\omega^n \rightarrow \omega$ (almost. . .)

Given an isomorphism $\xi: \mathcal{H} \rightarrow \mathcal{C}$. If $g: \omega^k \rightarrow \omega$ is bijective, then any k -ary $h \in \mathcal{H}$ is of the form $\alpha \circ g$.

Horn clone

Let \mathcal{H} be the closed clone generated by

- all unary injections, and
- a binary injection $f: \omega^2 \rightarrow \omega$.

\mathcal{H} consists of all injections $\omega^n \rightarrow \omega$ (almost...)

Given an isomorphism $\xi: \mathcal{H} \rightarrow \mathcal{C}$. If $g: \omega^k \rightarrow \omega$ is bijective, then any k -ary $h \in \mathcal{H}$ is of the form $\alpha \circ g$.

Sequences $(h_n)_{n \in \mathbb{N}} \subseteq \mathcal{H}^{(k)}$ are sequences $(\alpha_n \circ g)_{n \in \mathbb{N}}$.

Horn clone

Let \mathcal{H} be the closed clone generated by

- all unary injections, and
- a binary injection $f: \omega^2 \rightarrow \omega$.

\mathcal{H} consists of all injections $\omega^n \rightarrow \omega$ (almost...)

Given an isomorphism $\xi: \mathcal{H} \rightarrow \mathcal{C}$. If $g: \omega^k \rightarrow \omega$ is bijective, then any k -ary $h \in \mathcal{H}$ is of the form $\alpha \circ g$.

Sequences $(h_n)_{n \in \mathbb{N}} \subseteq \mathcal{H}^{(k)}$ are sequences $(\alpha_n \circ g)_{n \in \mathbb{N}}$.

$\lim \xi(h_n) = \lim(\xi(\alpha_n)) \circ \xi(g) = \xi(\lim(\alpha_n) \circ g) = \xi(\lim h_n)$

Horn clone

Let \mathcal{H} be the closed clone generated by

- all unary injections, and
- a binary injection $f: \omega^2 \rightarrow \omega$.

\mathcal{H} consists of all injections $\omega^n \rightarrow \omega$ (almost...)

Given an isomorphism $\xi: \mathcal{H} \rightarrow \mathcal{C}$. If $g: \omega^k \rightarrow \omega$ is bijective, then any k -ary $h \in \mathcal{H}$ is of the form $\alpha \circ g$.

Sequences $(h_n)_{n \in \mathbb{N}} \subseteq \mathcal{H}^{(k)}$ are sequences $(\alpha_n \circ g)_{n \in \mathbb{N}}$.

$\lim \xi(h_n) = \lim(\xi(\alpha_n)) \circ \xi(g) = \xi(\lim(\alpha_n) \circ g) = \xi(\lim h_n)$

Proposition

Every isomorphism $\xi: \mathcal{H} \rightarrow \mathcal{C}$ is continuous.

Gate coverings

Gate coverings

Definition

A *gate covering* of a topological clone \mathcal{C} consists of

Gate coverings

Definition

A *gate covering* of a topological clone \mathcal{C} consists of

- an open covering \mathcal{U} of \mathcal{C} ;
- for every $U \in \mathcal{U}$ a function $f_U \in U$;

Gate coverings

Definition

A *gate covering* of a topological clone \mathcal{C} consists of

- an open covering \mathcal{U} of \mathcal{C} ;
- for every $U \in \mathcal{U}$ a function $f_U \in U$;

such that for all converging sequences $(g^i)_{i \in \omega}$ in \mathcal{C} (say in some $U \in \mathcal{U}$, and say of arity n)

Gate coverings

Definition

A *gate covering* of a topological clone \mathcal{C} consists of

- an open covering \mathcal{U} of \mathcal{C} ;
- for every $U \in \mathcal{U}$ a function $f_U \in U$;

such that for all converging sequences $(g^i)_{i \in \omega}$ in \mathcal{C} (say in some $U \in \mathcal{U}$, and say of arity n)

there exist unary $(\alpha^i)_{i \in \mathbb{N}}$ and $(\beta_1^i)_{i \in \mathbb{N}}, \dots, (\beta_n^i)_{i \in \mathbb{N}}$ in \mathcal{C} with

- $g^i(x_1, \dots, x_n) = \alpha^i(f_U(\beta_1^i(x_1), \dots, \beta_n^i(x_n)))$ and
- $(\alpha^i)_{i \in \mathbb{N}}$ and $(\beta_1^i)_{i \in \mathbb{N}}, \dots, (\beta_n^i)_{i \in \mathbb{N}}$ converge.

Gate coverings II

Gate coverings II

Theorem (M. Bodirsky, M. Pinsker, AP '13)

If \mathcal{C} is a closed subclone of \mathcal{O} such that

Gate coverings II

Theorem (M. Bodirsky, M. Pinsker, AP '13)

If \mathcal{C} is a closed subclone of \mathcal{O} such that

- \mathcal{C} acts transitively;

Gate coverings II

Theorem (M. Bodirsky, M. Pinsker, AP '13)

If \mathcal{C} is a closed subclone of \mathcal{O} such that

- \mathcal{C} acts transitively;
- $\mathcal{C}^{(1)}$ has automatic homeomorphicity;

Gate coverings II

Theorem (M. Bodirsky, M. Pinsker, AP '13)

If \mathcal{C} is a closed subclone of \mathcal{O} such that

- \mathcal{C} acts transitively;
- $\mathcal{C}^{(1)}$ has automatic homeomorphicity;
- \mathcal{C} has a gate covering;

Theorem (M. Bodirsky, M. Pinsker, AP '13)

If \mathcal{C} is a closed subclone of \mathcal{O} such that

- \mathcal{C} acts transitively;
- $\mathcal{C}^{(1)}$ has automatic homeomorphicity;
- \mathcal{C} has a gate covering;

Then \mathcal{C} has automatic homeomorphicity.

Gate coverings II

Theorem (M. Bodirsky, M. Pinsker, AP '13)

If \mathcal{C} is a closed subclone of \mathcal{O} such that

- \mathcal{C} acts transitively;
- $\mathcal{C}^{(1)}$ has automatic homeomorphicity;
- \mathcal{C} has a gate covering;

Then \mathcal{C} has automatic homeomorphicity.

Corollary

\mathcal{H} has automatic homeomorphicity.

Automatic homeomorphicity of $\text{Pol}(V, E)$

Techniques

Automatic homeomorphicity of $\text{Pol}(V, E)$

Techniques

Gate coverings

Automatic homeomorphicity of $\text{Pol}(V, E)$

Techniques

Gate coverings

Topological Birkhoff

Automatic homeomorphicity of $\text{Pol}(V, E)$

Techniques

Gate coverings

Topological Birkhoff

Cheating!

$\text{Pol}(V, E)$ has automatic homeomorphicity

Every isomorphism $\xi : \text{End}(V, E) \rightarrow \mathcal{M}$ is continuous.

$\text{Pol}(V, E)$ has automatic homeomorphicity

Every isomorphism $\xi : \text{End}(V, E) \rightarrow \mathcal{M}$ is continuous.
 $\text{Pol}(V, E) \cap \mathcal{H}$ has automatic homeomorphicity.

$\text{Pol}(V, E)$ has automatic homeomorphicity

Every isomorphism $\xi : \text{End}(V, E) \rightarrow \mathcal{M}$ is continuous.

$\text{Pol}(V, E) \cap \mathcal{H}$ has automatic homeomorphicity.

Every $f \in \text{Pol}(V, E)$ decomposes as $f = h \circ g$ with $h \in \text{End}(V, E)$, $g \in \text{Pol}(V, E) \cap \mathcal{H}$.

$\text{Pol}(V, E)$ has automatic homeomorphicity

Every isomorphism $\xi : \text{End}(V, E) \rightarrow \mathcal{M}$ is continuous.

$\text{Pol}(V, E) \cap \mathcal{H}$ has automatic homeomorphicity.

Every $f \in \text{Pol}(V, E)$ decomposes as $f = h \circ g$ with $h \in \text{End}(V, E)$, $g \in \text{Pol}(V, E) \cap \mathcal{H}$.

Moreover, if $(f_i)_{i \in \mathbb{N}}$ is Cauchy, then \exists decompositions $f_i = h_i \circ g_i$ with $h_i \in \text{End}(V, E)$, $g_i \in \text{Pol}(V, E) \cap \mathcal{H}$ Cauchy sequences.

$\text{Pol}(V, E)$ has automatic homeomorphicity

Every isomorphism $\xi : \text{End}(V, E) \rightarrow \mathcal{M}$ is continuous.

$\text{Pol}(V, E) \cap \mathcal{H}$ has automatic homeomorphicity.

Every $f \in \text{Pol}(V, E)$ decomposes as $f = h \circ g$ with $h \in \text{End}(V, E)$, $g \in \text{Pol}(V, E) \cap \mathcal{H}$.

Moreover, if $(f_i)_{i \in \mathbb{N}}$ is Cauchy, then \exists decompositions $f_i = h_i \circ g_i$ with $h_i \in \text{End}(V, E)$, $g_i \in \text{Pol}(V, E) \cap \mathcal{H}$ Cauchy sequences.

All isomorphisms $\xi : \text{Pol}(V, E) \rightarrow \mathcal{C}$ are continuous.

$\text{Pol}(V, E)$ has automatic homeomorphicity

Every isomorphism $\xi : \text{End}(V, E) \rightarrow \mathcal{M}$ is continuous.

$\text{Pol}(V, E) \cap \mathcal{H}$ has automatic homeomorphicity.

Every $f \in \text{Pol}(V, E)$ decomposes as $f = h \circ g$ with $h \in \text{End}(V, E)$, $g \in \text{Pol}(V, E) \cap \mathcal{H}$.

Moreover, if $(f_i)_{i \in \mathbb{N}}$ is Cauchy, then \exists decompositions $f_i = h_i \circ g_i$ with $h_i \in \text{End}(V, E)$, $g_i \in \text{Pol}(V, E) \cap \mathcal{H}$ Cauchy sequences.

All isomorphisms $\xi : \text{Pol}(V, E) \rightarrow \mathcal{C}$ are continuous.

Openness follows from **TB** + **CH**.

Open problems

\exists ω -categorical Δ with a discontinuous automorphism of $\text{End}(\Delta)$.

Open problems

\exists ω -categorical Δ with a discontinuous automorphism of $\text{End}(\Delta)$.

Problem 1 Is there an ω -categorical Δ such that $\text{End}(\Delta)$ (or $\text{Pol}(\Delta)$) does not have reconstruction?

Open problems

\exists ω -categorical Δ with a discontinuous automorphism of $\text{End}(\Delta)$.

Problem 1 Is there an ω -categorical Δ such that $\text{End}(\Delta)$ (or $\text{Pol}(\Delta)$) does not have reconstruction?

Problem 2 Does $\text{End}(H_n, E)$ (or $\text{Pol}(H_n, E)$) have automatic homeomorphicity for (any) $n \geq 3$?