

Complexity of variety membership

Marcel Jackson
La Trobe University, Batmania, Australia



Prague ALCfest, June 2014

Var-mem

Var-Mem(**A**)

Instance: a finite algebra **B** of the same signature as **A**

Question: is $\mathbf{B} \in \mathbf{HSP}(\mathbf{A})$?

Var-mem

Var-Mem(**A**)

Instance: a finite algebra **B** of the same signature as **A**

Question: is $\mathbf{B} \in \mathbf{HSP}(\mathbf{A})$?

Var_{si}-Mem(**A**)

Instance: a finite algebra **B** of the same signature as **A**

Question: is $\mathbf{B} \in [\mathbf{HSP}(\mathbf{A})]_{\text{si}}$?

Var-mem

Var-Mem(**A**)

Instance: a finite algebra **B** of the same signature as **A**

Question: is $\mathbf{B} \in \mathbf{HSP}(\mathbf{A})$?

Var_{si}-Mem(**A**)

Instance: a finite algebra **B** of the same signature as **A**

Question: is $\mathbf{B} \in [\mathbf{HSP}(\mathbf{A})]_{\text{si}}$?

▷ *decomposition into subdirectly irreducibles can be performed in polynomial time (Demlová, 1982)*

Var-mem

Var-Mem(**A**)

Instance: a finite algebra **B** of the same signature as **A**

Question: is $\mathbf{B} \in \mathbf{HSP}(\mathbf{A})$?

Var_{si}-Mem(**A**)

Instance: a finite algebra **B** of the same signature as **A**

Question: is $\mathbf{B} \in [\mathbf{HSP}(\mathbf{A})]_{\text{si}}$?

- ▷ *decomposition into subdirectly irreducibles can be performed in polynomial time (Demlová, 1982)*
- ▷ *Var-Mem is solvable in 2-EXPTIME (Bergman and Slutzki, 2000).*

Var-mem(**A**) can be:

- NP-complete [Szekely, 7 (or 6) elements; 2002]

Var-mem(**A**) can be:

- NP-complete [Szekely, 7 (or 6) elements; 2002]
- NP-hard for a semigroup [J and McKenzie, 55 elements; 2006]
- NP-hard for an involuted semigroup [J and Volkov, 26 elements; 2010]

Var-mem(**A**) can be:

- NP-complete [Szekely, 7 (or 6) elements; 2002]
- NP-hard for a semigroup [J and McKenzie, 55 elements; 2006]
- NP-hard for an involuted semigroup [J and Volkov, 26 elements; 2010]
- PSPACE-complete [Kozik; 2007]

Var-mem(**A**) can be:

- NP-complete [Szekely, 7 (or 6) elements; 2002]
- NP-hard for a semigroup [J and McKenzie, 55 elements; 2006]
- NP-hard for an involuted semigroup [J and Volkov, 26 elements; 2010]
- PSPACE-complete [Kozik; 2007]
- EXPSPACE-complete [Kozik; 2007]

Var-mem(**A**) can be:

- NP-complete [Szekely, 7 (or 6) elements; 2002]
- NP-hard for a semigroup [J and McKenzie, 55 elements; 2006]
- NP-hard for an involuted semigroup [J and Volkov, 26 elements; 2010]
- PSPACE-complete [Kozik; 2007]
- EXSPACE-complete [Kozik; 2007]
- 2-EXPTIME-complete [Kozik; 2009]

Connections with finite basis problem

If \mathbf{A} has a finite basis for its equations

$\text{Var-Mem}(\mathbf{A})$ is solvable in polynomial time

Connections with finite basis problem

If \mathbf{A} has a finite basis for its equations

$\text{Var-Mem}(\mathbf{A})$ is solvable in polynomial time

The converse not true.

One example: Lyndon's algebra \mathbf{L}

Connections with finite basis problem

If \mathbf{A} has a finite basis for its equations

$\text{Var-Mem}(\mathbf{A})$ is solvable in polynomial time

The converse not true.

One example: Lyndon's algebra \mathbf{L}

$\text{Var-Mem}(\mathbf{L})$ is in NL

- ▶ (J and McNulty, 2011)

Eilenberg-Schützenberger

A is a finite algebra of finite signature

If $\text{HSP}_{\text{fin}}(\mathbf{A})$ is definable by a finite system of equations, is $\text{HSP}(\mathbf{A})$ finitely axiomatisable?

Eilenberg-Schützenberger

A is a finite algebra of finite signature

If $\text{HSP}_{\text{fin}}(\mathbf{A})$ is definable by a finite system of equations, is $\text{HSP}(\mathbf{A})$ finitely axiomatisable?

If **C** a counterexample

$\text{Var-Mem}(\mathbf{C}) \in \text{F0}$, even though $\text{HSP}(\mathbf{C})$ is not finitely axiomatisable?!?!?

Eilenberg-Schützenberger

A is a finite algebra of finite signature

If $\text{HSP}_{\text{fin}}(\mathbf{A})$ is definable by a finite system of equations, is $\text{HSP}(\mathbf{A})$ finitely axiomatisable?

First order Eilenberg-Schützenberger

Does $\text{HSP}_{\text{fin}}(\mathbf{A})$ first order definable imply $\text{HSP}(\mathbf{A})$ is first order definable?

Eilenberg-Schützenberger

A is a finite algebra of finite signature

If $\text{HSP}_{\text{fin}}(\mathbf{A})$ is definable by a finite system of equations, is $\text{HSP}(\mathbf{A})$ finitely axiomatisable?

First order Eilenberg-Schützenberger

Does $\text{HSP}_{\text{fin}}(\mathbf{A})$ first order definable imply $\text{HSP}(\mathbf{A})$ is first order definable?

A result: “no”

Eilenberg-Schützenberger

A is a finite algebra of finite signature

If $\text{HSP}_{\text{fin}}(\mathbf{A})$ is definable by a finite system of equations, is $\text{HSP}(\mathbf{A})$ finitely axiomatisable?

First order Eilenberg-Schützenberger

Does $\text{HSP}_{\text{fin}}(\mathbf{A})$ first order definable imply $\text{HSP}(\mathbf{A})$ is first order definable?

A result: “no”

There exists a finite algebra \mathbf{A} for which $\text{HSP}_{\text{fin}}(\mathbf{A})$ is definable by a $\forall^*\exists^*\forall^*$ sentence, but not by a $\forall^*\exists^*$ -sentence, nor by any finite system of pseudo-equations. In particular, $\text{HSP}(\mathbf{A})$ is not first order definable.

Preservation Theorems

Łos-Tarski Theorem

If $K = \text{Mod}(\phi)$ and is closed under taking substructures, then ϕ is equivalent to a \forall^* -sentence

Preservation Theorems

Łos-Tarski Theorem

If $K = \text{Mod}(\phi)$ and is closed under taking substructures, then ϕ is equivalent to a \forall^* -sentence

Tait (1959), Gurevich-Shelah (1980s)

Łos-Tarski Theorem is not true when restricted to finite structures

Preservation Theorems

Łos-Tarski Theorem

If $K = \text{Mod}(\phi)$ and is closed under taking substructures, then ϕ is equivalent to a \forall^* -sentence

Tait (1959), Gurevich-Shelah (1980s)

Łos-Tarski Theorem is not true when restricted to finite structures

Homomorphism Preservation

Rossman (2005/8) showed that the homomorphism preservation theorem *does* hold at the finite level

Preservation Theorems

Łos-Tarski Theorem

If $K = \text{Mod}(\phi)$ and is closed under taking substructures, then ϕ is equivalent to a \forall^* -sentence

Tait (1959), Gurevich-Shelah (1980s)

Łos-Tarski Theorem is not true when restricted to finite structures

Homomorphism Preservation

Rossman (2005/8) showed that the homomorphism preservation theorem *does* hold at the finite level

This fact is crucial in the classification of first order definable CSPs

ISP preservation?

Problem 1, Alechina and Gurevich 1997

If $K = \text{Mod}_{\text{fin}}(\phi)$ is closed under SP_{fin} , then is it true that ϕ is equivalent to a universal Horn sentence on finite structures?

ISP preservation?

Problem 1, Alechina and Gurevich 1997

If $K = \text{Mod}_{\text{fin}}(\phi)$ is closed under SP_{fin} , then is it true that ϕ is equivalent to a universal Horn sentence on finite structures?

No

- $\text{SP}(M_{3-3})$ is not finitely based (Belkin, 1978)

ISP preservation?

Problem 1, Alechina and Gurevich 1997

If $K = \text{Mod}_{\text{fin}}(\phi)$ is closed under SP_{fin} , then is it true that ϕ is equivalent to a universal Horn sentence on finite structures?

No

- $\text{SP}(M_{3-3})$ is not finitely based (Belkin, 1978)
- $\text{S}_c\text{P}(M_{3-3})$ (where M_{3-3} has discrete topology) is first order definable amongst Boolean topological algebras (Clark, Davey, J, Pitkethly, 2008)

ISP preservation?

Problem 1, Alechina and Gurevich 1997

If $K = \text{Mod}_{\text{fin}}(\phi)$ is closed under SP_{fin} , then is it true that ϕ is equivalent to a universal Horn sentence on finite structures?

No

- $\text{SP}(M_{3-3})$ is not finitely based (Belkin, 1978)
- $\text{S}_c\text{P}(M_{3-3})$ (where M_{3-3} has discrete topology) is first order definable amongst Boolean topological algebras (Clark, Davey, J, Pitkethly, 2008)
- $[\text{S}_c\text{P}(M_{3-3})]_{\text{fin}}$ and $[\text{SP}(M_{3-3})]_{\text{fin}}$ coincide

ISP preservation?

Problem 1, Alechina and Gurevich 1997

If $K = \text{Mod}_{\text{fin}}(\phi)$ is closed under SP_{fin} , then is it true that ϕ is equivalent to a universal Horn sentence on finite structures?

No

- $\text{SP}(M_{3-3})$ is not finitely based (Belkin, 1978)
- $\text{S}_c\text{P}(M_{3-3})$ (where M_{3-3} has discrete topology) is first order definable amongst Boolean topological algebras (Clark, Davey, J, Pitkethly, 2008)
- $[\text{S}_c\text{P}(M_{3-3})]_{\text{fin}}$ and $[\text{SP}(M_{3-3})]_{\text{fin}}$ coincide
- $[\text{SP}(M_{3-3})]_{\text{fin}}$ is not definable by a universal Horn sentence

$\forall^* \exists^* \forall^*$ for finite semilattice-based algebras

S finite, with a semilattice term \wedge

T $\in \mathbb{SP}_{\text{fin}}(\mathbf{S})$ if and only if

$\forall^* \exists^* \forall^*$ for finite semilattice-based algebras

S finite, with a semilattice term \wedge

T $\in \mathbb{SP}_{\text{fin}}(\mathbf{S})$ if and only if \wedge is a semilattice operation on **T** and

$\forall^* \exists^* \forall^*$ for finite semilattice-based algebras

S finite, with a semilattice term \wedge

T $\in \mathbb{SP}_{\text{fin}}(\mathbf{S})$ if and only if \wedge is a semilattice operation on **T** and for each $a \neq b$ in T there is a subalgebra **U** of **S** and (distinct) $t_u \in T$ for each $u \in U$

$\forall^* \exists^* \forall^*$ for finite semilattice-based algebras

S finite, with a semilattice term \wedge

T $\in \mathsf{SP}_{\text{fin}}(\mathbf{S})$ if and only if \wedge is a semilattice operation on **T** and for each $a \neq b$ in T there is a subalgebra **U** of **S** and (distinct) $t_u \in T$ for each $u \in U$ such that the equivalence relation $x \equiv y$ defined by $t_u \leq x \Leftrightarrow t_u \leq y$ is a congruence relation of **T**


$\forall^* \exists^* \forall^*$ for finite semilattice-based algebras

S finite, with a semilattice term \wedge

T $\in \mathsf{SP}_{\text{fin}}(\mathbf{S})$ if and only if \wedge is a semilattice operation on **T** and for each $a \neq b$ in T there is a subalgebra **U** of **S** and (distinct) $t_u \in T$ for each $u \in U$ such that the equivalence relation $x \equiv y$ defined by $t_u \leq x \Leftrightarrow t_u \leq y$ is a congruence relation of **T** with $a \not\equiv b$ and with **T**/ \equiv isomorphic to **U**.


$\text{HSP}_{\text{fin}}(\mathbf{C})$ finitely axiomatisable but
 $\text{HSP}(\mathbf{C})$ not finitely axiomatisable

Example: $\mathbf{C} := M_{3-3}^b$



$\text{HSP}_{\text{fin}}(\mathbf{C})$ finitely axiomatisable but
 $\text{HSP}(\mathbf{C})$ not finitely axiomatisable

Example: $\mathbf{C} := M_{3-3}^b$



$\text{HSP}_{\text{fin}}(\mathbf{C})$ finitely axiomatisable but
 $\text{HSP}(\mathbf{C})$ not finitely axiomatisable

Example: $\mathbf{C} := M_{3-3}^b$

$\text{HSP}_{\text{fin}}(\mathbf{C})$ definable by a $\forall^*\exists^*\forall^*$ -sentence, but not by any $\forall^*\exists^*$ -sentence

$\text{HSP}_{\text{fin}}(\mathbf{C})$ finitely axiomatisable but
 $\text{HSP}(\mathbf{C})$ not finitely axiomatisable

Example: $\mathbf{C} := M_{3-3}^b$

$\text{HSP}_{\text{fin}}(\mathbf{C})$ definable by a $\forall^*\exists^*\forall^*$ -sentence, but not by any $\forall^*\exists^*$ -sentence

▷ *Simultaneous failure of both the ISP-Preservation Theorem, and the Łos-Tarski Theorem at the finite level!* (And the HSP -preservation theorem at the finite level)

$\text{HSP}_{\text{fin}}(\mathbf{C})$ finitely axiomatisable but
 $\text{HSP}(\mathbf{C})$ not finitely axiomatisable

Example: $\mathbf{C} := M_{3-3}^b$

$\text{HSP}_{\text{fin}}(\mathbf{C})$ definable by a $\forall^*\exists^*\forall^*$ -sentence, but not by any $\forall^*\exists^*$ -sentence

▷ *Simultaneous failure of both the ISP-Preservation Theorem, and the Łos-Tarski Theorem at the finite level!* (And the HSP -preservation theorem at the finite level)

▷ Shows $\text{Var-Mem}(\mathbf{C}) \in \text{FO}$ not equivalent to \mathbf{C} being finitely based.

Related examples

Theorem

For every finite relational structure \mathbf{R} of finite signature, there is a finite algebra $\mathbf{A}_{\mathbf{R}}$ such that $\text{CSP}(\mathbf{R})$ is first order equivalent to both $\text{Var-Mem}(\mathbf{A}_{\mathbf{R}})$ and $\text{Var}_{\text{si-Mem}}(\mathbf{A}_{\mathbf{R}})$

Related examples

Theorem

For every finite relational structure \mathbf{R} of finite signature, there is a finite algebra $\mathbf{A}_{\mathbf{R}}$ such that $\text{CSP}(\mathbf{R})$ is first order equivalent to both $\text{Var-Mem}(\mathbf{A}_{\mathbf{R}})$ and $\text{Var}_{\text{si-Mem}}(\mathbf{A}_{\mathbf{R}})$

Corollary

$\text{Var-Mem}(\mathbf{A}_{\mathbf{R}})$ can be L-complete, NL-complete, P-complete, Mod_pL -complete, etc (infinitely many tractable complexities)

Related examples

Theorem

For every finite relational structure \mathbf{R} of finite signature, there is a finite algebra $\mathbf{A}_{\mathbf{R}}$ such that $\text{CSP}(\mathbf{R})$ is first order equivalent to both $\text{Var-Mem}(\mathbf{A}_{\mathbf{R}})$ and $\text{Var}_{\text{si-Mem}}(\mathbf{A}_{\mathbf{R}})$

Corollary

$\text{Var-Mem}(\mathbf{A}_{\mathbf{R}})$ can be L-complete, NL-complete, P-complete, Mod_pL -complete, etc (infinitely many tractable complexities)

Interesting problem for future

Is it possible that if $\text{Var-Mem}(A)$ not in P then it is NP-hard with respect to many-one reductions?

For any complexity class $K \subsetneq \text{PSPACE}$

Given finite algebra \mathbf{A} it is undecidable whether or not $\text{Var-Mem}(\mathbf{A})$ is in K .

For any complexity class $K \subsetneq PSPACE$

Given finite algebra \mathbf{A} it is undecidable whether or not $\text{Var-Mem}(\mathbf{A})$ is in K .

(*In contrast, it is NP-complete to decide if $\text{CSP}(\mathbf{R})$ is in FO* ; Larose, Loten, Tardif 2006)

Central technique

First order definable subdirect decomposition

There is a formula $\pi(x, y, u, v)$ such that $\{(x, y) \mid \mathbf{A} \models \pi(x, y, a, b)\}$ is a congruence $\theta_{a,b}$ maximal with respect to $(a, b) \notin \theta_{a,b}$

Consequence

$\mathbf{A}/\theta_{a,b}$ is first order definable from \mathbf{A} for each $a \neq b$

Pointed semi-discriminator varieties

1. If \mathbf{P} has a projection term $t(x, y) \approx x$, then

$$[\text{HSP}(\mathbf{P}^b)]_{\text{si}} = [\text{SPP}_{\text{u}}(\mathbf{P})]^b$$

Pointed semi-discriminator varieties

1. If \mathbf{P} has a projection term $t(x, y) \approx x$, then

$$[\text{HSP}(\mathbf{P}^b)]_{\text{si}} = [\text{SP}(\mathbf{P})]^b$$

Pointed semi-discriminator varieties

1. If \mathbf{P} has a projection term $t(x, y) \approx x$, then

$$[\text{HSP}(\mathbf{P}^b)]_{\text{si}} = [\text{SP}(\mathbf{P})]^b$$

$\Rightarrow \text{Var}_{\text{si-Mem}}(\mathbf{P}^b)$ is FO-equivalent to membership problem for $\text{SP}_{\text{fin}}(\mathbf{P})$

Pointed semi-discriminator varieties

1. If \mathbf{P} has a projection term $t(x, y) \approx x$, then

$$[\mathbf{HSP}(\mathbf{P}^b)]_{\text{si}} = [\mathbf{SP}(\mathbf{P})]^b$$

$\Rightarrow \text{Var}_{\text{si-Mem}}(\mathbf{P}^b)$ is FO-equivalent to membership problem for $\mathbf{SP}_{\text{fin}}(\mathbf{P})$

2. $\mathbb{V}(\mathbf{P}^b)$ has first order definable subdirect decomposition: for $a \not\leq b$

$$\pi(x, y, a, b) = (\neg t(x, a) \approx a \ \& \ \neg t(y, a) \approx a) \ \mathbb{V}(t(x \wedge y, a) \approx a)$$

Pointed semi-discriminator varieties

1. If \mathbf{P} has a projection term $t(x, y) \approx x$, then

$$[\text{HSP}(\mathbf{P}^b)]_{\text{si}} = [\text{SP}(\mathbf{P})]^b$$

$\Rightarrow \text{Var}_{\text{si-Mem}}(\mathbf{P}^b)$ is FO-equivalent to membership problem for $\text{SP}_{\text{fin}}(\mathbf{P})$

2. $\mathbb{V}(\mathbf{P}^b)$ has first order definable subdirect decomposition: for $a \not\leq b$

$$\pi(x, y, a, b) = (\neg t(x, a) \approx a \wedge \neg t(y, a) \approx a) \vee (t(x \wedge y, a) \approx a)$$

$\Rightarrow \text{Var-Mem}(\mathbf{P}^b)$ reduces to $\text{Var}_{\text{si-Mem}}(\mathbf{P}^b)$ by a sort of “first order Turing reduction”

Pointed semi-discriminator varieties

1. If \mathbf{P} has a projection term $t(x, y) \approx x$, then

$$[\text{HSP}(\mathbf{P}^b)]_{\text{si}} = [\text{SP}(\mathbf{P})]^b$$

$\Rightarrow \text{Var}_{\text{si-Mem}}(\mathbf{P}^b)$ is FO-equivalent to membership problem for $\text{SP}_{\text{fin}}(\mathbf{P})$

2. $\mathbb{V}(\mathbf{P}^b)$ has first order definable subdirect decomposition: for $a \not\leq b$

$$\pi(x, y, a, b) = (\neg t(x, a) \approx a \wedge \neg t(y, a) \approx a) \vee (t(x \wedge y, a) \approx a)$$

$\Rightarrow \text{Var-Mem}(\mathbf{P}^b)$ reduces to $\text{Var}_{\text{si-Mem}}(\mathbf{P}^b)$ by a sort of “first order Turing reduction”

3. If \mathbf{P} comes from a relational structure, then a further trick enables a genuine first order reduction from $\text{Var-Mem}(\mathbf{P}^b)$ to $\text{Var}_{\text{si-Mem}}(\mathbf{P}^b)$

Pointed semi-discriminator varieties

1. If \mathbf{P} has a projection term $t(x, y) \approx x$, then

$$[\text{HSP}(\mathbf{P}^b)]_{\text{si}} = [\text{SP}(\mathbf{P})]^b$$

$\Rightarrow \text{Var}_{\text{si-Mem}}(\mathbf{P}^b)$ is FO-equivalent to membership problem for $\text{SP}_{\text{fin}}(\mathbf{P})$

2. $\mathbb{V}(\mathbf{P}^b)$ has first order definable subdirect decomposition: for $a \not\leq b$

$$\pi(x, y, a, b) = (\neg t(x, a) \approx a \wedge \neg t(y, a) \approx a) \vee (t(x \wedge y, a) \approx a)$$

$\Rightarrow \text{Var-Mem}(\mathbf{P}^b)$ reduces to $\text{Var}_{\text{si-Mem}}(\mathbf{P}^b)$ by a sort of “first order Turing reduction”

3. If \mathbf{P} comes from a relational structure, then a further trick enables a genuine first order reduction from $\text{Var-Mem}(\mathbf{P}^b)$ to $\text{Var}_{\text{si-Mem}}(\mathbf{P}^b)$
4. But first replace \mathbf{P} by \mathbf{P}^\sharp

For M_{3-3}

If ϕ defines $[\text{SP}_{\text{fin}}(\mathbf{P})]^b$, then a finite system of equations plus

$$\forall u \forall v \quad u \not\approx v \rightarrow \phi_{u,v}$$

defines $\text{HSP}_{\text{fin}}(\mathbf{P}^b)$, where $\phi_{u,v}$ has all instances of $\dots \approx \dots$ replaced by $\pi(\dots, \dots, u, v)$

For M_{3-3}

If ϕ defines $[\text{SP}_{\text{fin}}(\mathbf{P})]^b$, then a finite system of equations plus

$$\forall u \forall v \quad u \not\approx v \rightarrow \phi_{u,v}$$

defines $\text{HSP}_{\text{fin}}(\mathbf{P}^b)$, where $\phi_{u,v}$ has all instances of $\dots \approx \dots$ replaced by $\pi(\dots, \dots, u, v)$

ϕ is a $\forall^* \exists^* \forall^*$ -sentence for $\mathbf{P} = M_{3-3}$

For M_{3-3}

If ϕ defines $[\mathbb{SP}_{\text{fin}}(\mathbf{P})]^b$, then a finite system of equations plus

$$\forall u \forall v \quad u \not\approx v \rightarrow \phi_{u,v}$$

defines $\mathbb{HSP}_{\text{fin}}(\mathbf{P}^b)$, where $\phi_{u,v}$ has all instances of $\dots \approx \dots$ replaced by $\pi(\dots, \dots, u, v)$

ϕ is a $\forall^* \exists^* \forall^*$ -sentence for $\mathbf{P} = M_{3-3}$

A newish preservation theorem

If K is a uniformly locally finite \mathbb{SP}_{fin} -class of structures contained in some locally finite and finitely axiomatisable \mathbb{SP} -class, then K is definable by a $\forall^* \exists^*$ -sentence if and only if it is definable by a universal Horn sentence