

The structure of medial quandles

Přemysl Jedlička¹, Agata Pilitowska², David Stanovský³, Anna Zamojska-Dzienio²

¹Department of Mathematics, Faculty of Engineering, Czech University of Life Sciences,

²Faculty of Mathematics and Information Science, Warsaw University of Technology,

³Department of Algebra, Faculty of Mathematics and Physics, Charles University

Prague, July 2014

Quandles

A binary algebra (Q, \cdot) is called a **quandle** if it is:

- **idempotent**: $xx = x$ for each $x \in Q$,
- **left distributive**: $x(yz) = (xy)(xz)$ for every $x, y, z \in Q$,
- a **left quasigroup**: the equation $xu = y$ has a unique solution $u \in Q$ for every $x, y \in Q$.

Quandles

A binary algebra (Q, \cdot) is called a **quandle** if it is:

- **idempotent**: $xx = x$ for each $x \in Q$,
- **left distributive**: $x(yz) = (xy)(xz)$ for every $x, y, z \in Q$,
- a **left quasigroup**: the equation $xu = y$ has a unique solution $u \in Q$ for every $x, y \in Q$.

A quandle Q is called **medial** if

$$(xy)(uv) = (xu)(yv)$$

for every $x, y, u, v \in Q$.

Affine quandles

Affine quandles

Example

Let A be an abelian group, k its endomorphism, and define an operation on the set A by

$$a * b = (1 - k)(a) + k(b).$$

The resulting algebra $\text{Aff}(A, k) = (A, *)$ is called **affine** over the group A . It is idempotent and medial. If k is an automorphism then it is a medial quandle, called **affine quandle** over A .

Affine quandles

Example

Let A be an abelian group, k its endomorphism, and define an operation on the set A by

$$a * b = (1 - k)(a) + k(b).$$

The resulting algebra $\text{Aff}(A, k) = (A, *)$ is called **affine** over the group A . It is idempotent and medial. If k is an automorphism then it is a medial quandle, called **affine quandle** over A .

What is the role of affine quandles in the class of medial quandles?

Two important groups

Two important groups

The (left) **multiplication group** of a quandle Q is the permutation group generated by left translations, i.e.,

$$\text{LMlt}(Q) = \langle L_a \mid a \in Q \rangle \leq S_Q.$$

$$L_a: Q \rightarrow Q; \quad L_a(x) = ax$$

Two important groups

The (left) **multiplication group** of a quandle Q is the permutation group generated by left translations, i.e.,

$$\text{LMlt}(Q) = \langle L_a \mid a \in Q \rangle \leq S_Q.$$

$$L_a: Q \rightarrow Q; \quad L_a(x) = ax$$

The **group of displacements** is the subgroup

$$\text{Dis}(Q) = \langle L_a L_b^{-1} \mid a, b \in Q \rangle.$$

Two important groups

The (left) **multiplication group** of a quandle Q is the permutation group generated by left translations, i.e.,

$$\text{LMlt}(Q) = \langle L_a \mid a \in Q \rangle \leq S_Q.$$

$$L_a: Q \rightarrow Q; \quad L_a(x) = ax$$

The **group of displacements** is the subgroup

$$\text{Dis}(Q) = \langle L_a L_b^{-1} \mid a, b \in Q \rangle.$$

Both groups are normal subgroups of $\text{Aut}(Q)$.

Two important groups

Proposition

Let Q be a quandle. Then

- ❶ $\text{Dis}(Q) = \{L_{a_1}^{k_1} \dots L_{a_n}^{k_n} : a_1, \dots, a_n \in Q \text{ and } \sum_{i=1}^n k_i = 0\};$
- ❷ *the natural actions of $\text{LMlt}(Q)$ and $\text{Dis}(Q)$ on Q have the same orbits;*
- ❸ *Q is medial if and only if $\text{Dis}(Q)$ is abelian.*

Orbit decomposition

Orbits of Q are the orbits of transitivity of the groups $\text{LMlt}(Q)$ and $\text{Dis}(Q)$ denoted

$$Qe = \{\alpha(e) \mid \alpha \in \text{LMlt}(Q)\} = \{\alpha(e) \mid \alpha \in \text{Dis}(Q)\},$$

i.e. Qe is the orbit containing an element $e \in Q$.

Orbit decomposition

Orbits of Q are the orbits of transitivity of the groups $\text{LMlt}(Q)$ and $\text{Dis}(Q)$ denoted

$$Qe = \{\alpha(e) \mid \alpha \in \text{LMlt}(Q)\} = \{\alpha(e) \mid \alpha \in \text{Dis}(Q)\},$$

i.e. Qe is the orbit containing an element $e \in Q$.

Orbits are subquandles of Q .

Orbit decomposition

Orbits of Q are the orbits of transitivity of the groups $\text{LMlt}(Q)$ and $\text{Dis}(Q)$ denoted

$$Qe = \{\alpha(e) \mid \alpha \in \text{LMlt}(Q)\} = \{\alpha(e) \mid \alpha \in \text{Dis}(Q)\},$$

i.e. Qe is the orbit containing an element $e \in Q$.

Orbits are subquandles of Q .

Let $\alpha(e), \beta(e) \in Qe$ with $\alpha, \beta \in \text{Dis}(Q)$ and put

$$\alpha(e) + \beta(e) = \alpha\beta(e) \quad \text{and} \quad -\alpha(e) = \alpha^{-1}(e).$$

Then $\text{Orb}_Q(e) = (Qe, +, -, e)$ is an abelian group, called the **orbit group** for Qe .

Orbit decomposition

Orbits of Q are the orbits of transitivity of the groups $\text{LMlt}(Q)$ and $\text{Dis}(Q)$ denoted

$$Qe = \{\alpha(e) \mid \alpha \in \text{LMlt}(Q)\} = \{\alpha(e) \mid \alpha \in \text{Dis}(Q)\},$$

i.e. Qe is the orbit containing an element $e \in Q$.

Orbits are subquandles of Q .

Let $\alpha(e), \beta(e) \in Qe$ with $\alpha, \beta \in \text{Dis}(Q)$ and put

$$\alpha(e) + \beta(e) = \alpha\beta(e) \quad \text{and} \quad -\alpha(e) = \alpha^{-1}(e).$$

Then $\text{Orb}_Q(e) = (Qe, +, -, e)$ is an abelian group, called the **orbit group** for Qe .

Every orbit of a medial quandle is an affine quandle:

$$Qe = \text{Aff}(\text{Orb}_Q(e), L_e).$$

Construction

An **affine mesh** over a non-empty set I is a triple

$$\mathcal{A} = ((A_i)_{i \in I}, (\varphi_{i,j})_{i,j \in I}, (c_{i,j})_{i,j \in I})$$

where A_i are abelian groups, $\varphi_{i,j} : A_i \rightarrow A_j$ homomorphisms, and $c_{i,j} \in A_j$ constants, satisfying the following conditions for every $i, j, j', k \in I$:

(M1) $1 - \varphi_{i,i}$ is an automorphism of A_i ;

(M2) $c_{i,i} = 0$;

(M3) $\varphi_{j,k} \varphi_{i,j} = \varphi_{j',k} \varphi_{i,j'}$, i.e., the following diagram commutes:

$$\begin{array}{ccc} A_i & \xrightarrow{\varphi_{i,j}} & A_j \\ \downarrow \varphi_{i,j'} & & \downarrow \varphi_{j,k} \\ A_{j'} & \xrightarrow{\varphi_{j',k}} & A_k \end{array}$$

(M4) $\varphi_{j,k}(c_{i,j}) = \varphi_{k,k}(c_{i,k} - c_{j,k})$.

Construction

The **sum of an affine mesh** $((A_i)_{i \in I}, (\varphi_{i,j})_{i,j \in I}, (c_{i,j})_{i,j \in I})$ over a set I is a binary algebra defined on the disjoint union of the sets A_i , with operation

$$a * b = c_{i,j} + \varphi_{i,j}(a) + (1 - \varphi_{j,j})(b)$$

for every $a \in A_i$ and $b \in A_j$.

Construction

The **sum of an affine mesh** $((A_i)_{i \in I}, (\varphi_{i,j})_{i,j \in I}, (c_{i,j})_{i,j \in I})$ over a set I is a binary algebra defined on the disjoint union of the sets A_i , with operation

$$a * b = c_{i,j} + \varphi_{i,j}(a) + (1 - \varphi_{j,j})(b)$$

for every $a \in A_i$ and $b \in A_j$.

Every fibre $(A_i, *)$ is a subquandle of the sum. Moreover it is affine and equal to $\text{Aff}(A_i, 1 - \varphi_{i,i})$.

Construction

The **sum of an affine mesh** $((A_i)_{i \in I}, (\varphi_{i,j})_{i,j \in I}, (c_{i,j})_{i,j \in I})$ over a set I is a binary algebra defined on the disjoint union of the sets A_i , with operation

$$a * b = c_{i,j} + \varphi_{i,j}(a) + (1 - \varphi_{j,j})(b)$$

for every $a \in A_i$ and $b \in A_j$.

Every fibre $(A_i, *)$ is a subquandle of the sum. Moreover it is affine and equal to $\text{Aff}(A_i, 1 - \varphi_{i,i})$.

Lemma

The sum of an affine mesh is a medial quandle.

Representation theorem

Representation theorem

An affine mesh $((A_i)_{i \in I}, (\varphi_{i,j})_{i,j \in I}, (c_{i,j})_{i,j \in I})$ over a set I is called **indecomposable** if

$$A_j = \left\langle \bigcup_{i \in I} (c_{i,j} + \text{Im}(\varphi_{i,j})) \right\rangle,$$

for every $j \in I$.

Representation theorem

An affine mesh $((A_i)_{i \in I}, (\varphi_{i,j})_{i,j \in I}, (c_{i,j})_{i,j \in I})$ over a set I is called **indecomposable** if

$$A_j = \left\langle \bigcup_{i \in I} (c_{i,j} + \text{Im}(\varphi_{i,j})) \right\rangle,$$

for every $j \in I$.

Theorem

A binary algebra is a medial quandle if and only if it is the sum of an indecomposable affine mesh. The orbits of the quandle coincide with the groups of the mesh.

Homologous meshes

Homologous meshes

Two affine meshes $\mathcal{A} = ((A_i)_{i \in I}, (\varphi_{i,j})_{i,j \in I}, (c_{i,j})_{i,j \in I})$ and $\mathcal{A}' = ((A'_i)_{i \in I}, (\varphi'_{i,j})_{i,j \in I}, (c'_{i,j})_{i,j \in I})$, over the same index set I , are **homologous**, if there is a permutation π of the set I , group isomorphisms $\psi_i : A_i \mapsto A'_{\pi i}$, and constants $d_i \in A'_{\pi i}$, such that, for every $i, j \in I$,

(H1) $\psi_j \varphi_{i,j} = \varphi'_{\pi i, \pi j} \psi_i$, i.e., the following diagram commutes:

$$\begin{array}{ccc} A_i & \xrightarrow{\varphi_{i,j}} & A_j \\ \downarrow \psi_i & & \downarrow \psi_j \\ A'_{\pi i} & \xrightarrow{\varphi'_{\pi i, \pi j}} & A'_{\pi j} \end{array}$$

(H2) $\psi_j(c_{i,j}) = c'_{\pi i, \pi j} + \varphi'_{\pi i, \pi j}(d_i) - \varphi'_{\pi j, \pi j}(d_j)$.

Isomorphism theorem

Theorem

Let $\mathcal{A} = ((A_i)_{i \in I}, (\varphi_{i,j})_{i,j \in I}, (c_{i,j})_{i,j \in I})$ and $\mathcal{A}' = ((A'_i)_{i \in I}, (\varphi'_{i,j})_{i,j \in I}, (c'_{i,j})_{i,j \in I})$ be two indecomposable affine meshes, over the same index set I . Then the sums of \mathcal{A} and \mathcal{A}' are isomorphic quandles if and only if the meshes \mathcal{A} , \mathcal{A}' are homologous.