

On the complexity of testing for Jónsson terms

Alexandr Kazda Matt Valeriote

Vanderbilt University

McMaster University

30 June 2014

Congruence Distributivity

Theorem (Jónsson)

\mathcal{V} is *congruence distributive* if and only if for some $k > 1$ it has terms $p_i(x, y, z)$ for $0 \leq i \leq k$ which satisfy the identities:

$$p_0(x, y, z) \approx x$$

$$p_i(x, y, x) \approx x \text{ for all } i$$

$$p_i(x, x, y) \approx p_{i+1}(x, x, y) \text{ for all } i \text{ even}$$

$$p_i(x, y, y) \approx p_{i+1}(x, y, y) \text{ for all } i \text{ odd}$$

$$p_k(x, y, z) \approx z$$

Congruence Distributivity

Theorem (Jónsson)

\mathcal{V} is *congruence distributive* if and only if for some $k > 1$ it has terms $p_i(x, y, z)$ for $0 \leq i \leq k$ which satisfy the identities:

$$p_0(x, y, z) \approx x$$

$$p_i(x, y, x) \approx x \text{ for all } i$$

$$p_i(x, x, y) \approx p_{i+1}(x, x, y) \text{ for all } i \text{ even}$$

$$p_i(x, y, y) \approx p_{i+1}(x, y, y) \text{ for all } i \text{ odd}$$

$$p_k(x, y, z) \approx z$$

Definition

For $k > 1$, an algebra \mathbf{A} is said to be a *CD(k) algebra* if it has a sequence of term operations $p_i(x, y, z)$, $0 \leq i \leq k$, that satisfy the equations in the above theorem.

Testing for Jónsson terms

Question

Given a finite algebra \mathbf{A} and $k > 1$, how hard is it to determine if \mathbf{A} is a $\text{CD}(k)$ algebra?

Testing for Jónsson terms

Question

Given a finite algebra \mathbf{A} and $k > 1$, how hard is it to determine if \mathbf{A} is a $\text{CD}(k)$ algebra?

Remark

There is a straightforward, but inefficient algorithm to settle this question: Compute the free algebra in $\mathbf{V}(\mathbf{A})$ generated by $\{x, y, z\}$ and look for a sequence of Jónsson terms of length $k + 1$.

Testing for Jónsson terms

Question

Given a finite algebra \mathbf{A} and $k > 1$, how hard is it to determine if \mathbf{A} is a $\text{CD}(k)$ algebra?

Remark

There is a straightforward, but inefficient algorithm to settle this question: Compute the free algebra in $\mathbf{V}(\mathbf{A})$ generated by $\{x, y, z\}$ and look for a sequence of Jónsson terms of length $k + 1$.

Testing for Jónsson terms

Question

Given a finite algebra \mathbf{A} and $k > 1$, how hard is it to determine if \mathbf{A} is a $\text{CD}(k)$ algebra?

Remark

There is a straightforward, but inefficient algorithm to settle this question: Compute the free algebra in $\mathbf{V}(\mathbf{A})$ generated by $\{x, y, z\}$ and look for a sequence of Jónsson terms of length $k + 1$. As a function of $|A|$, the run time of this algorithm grows exponentially.

Testing for Jónsson terms

Question

Given a finite algebra \mathbf{A} and $k > 1$, how hard is it to determine if \mathbf{A} is a $\text{CD}(k)$ algebra?

Remark

There is a straightforward, but inefficient algorithm to settle this question: Compute the free algebra in $\mathbf{V}(\mathbf{A})$ generated by $\{x, y, z\}$ and look for a sequence of Jónsson terms of length $k + 1$.

As a function of $|\mathbf{A}|$, the run time of this algorithm grows exponentially.

Theorem (Freese-Val., Horowitz)

Let \mathbf{A} be a finite algebra. The following decision problems are EXP-TIME complete:

Testing for Jónsson terms

Question

Given a finite algebra \mathbf{A} and $k > 1$, how hard is it to determine if \mathbf{A} is a $\text{CD}(k)$ algebra?

Remark

There is a straightforward, but inefficient algorithm to settle this question: Compute the free algebra in $\mathbf{V}(\mathbf{A})$ generated by $\{x, y, z\}$ and look for a sequence of Jónsson terms of length $k + 1$.

As a function of $|A|$, the run time of this algorithm grows exponentially.

Theorem (Freese-Val., Horowitz)

Let \mathbf{A} be a finite algebra. The following decision problems are EXP-TIME complete:

- *Is \mathbf{A} is $\text{CD}(k)$ for some $k > 1$?*

Testing for Jónsson terms

Question

Given a finite algebra \mathbf{A} and $k > 1$, how hard is it to determine if \mathbf{A} is a $\text{CD}(k)$ algebra?

Remark

There is a straightforward, but inefficient algorithm to settle this question: Compute the free algebra in $\mathbf{V}(\mathbf{A})$ generated by $\{x, y, z\}$ and look for a sequence of Jónsson terms of length $k + 1$.

As a function of $|A|$, the run time of this algorithm grows exponentially.

Theorem (Freese-Val., Horowitz)

Let \mathbf{A} be a finite algebra. The following decision problems are EXP-TIME complete:

- Is \mathbf{A} is $\text{CD}(k)$ for some $k > 1$?*
- For a fixed $k > 2$, is \mathbf{A} $\text{CD}(k)$?*

Idempotent Algebras

Definition

Idempotent Algebras

Definition

- An operation $f(x_1, x_2, \dots, x_n)$ on a set A is **idempotent** if for all $a \in A$, $f(a, a, \dots, a) = a$.

Idempotent Algebras

Definition

- An operation $f(x_1, x_2, \dots, x_n)$ on a set A is **idempotent** if for all $a \in A$, $f(a, a, \dots, a) = a$.
- An algebra is **idempotent** if all of its basic operations are idempotent.

Idempotent Algebras

Definition

- An operation $f(x_1, x_2, \dots, x_n)$ on a set A is **idempotent** if for all $a \in A$, $f(a, a, \dots, a) = a$.
- An algebra is **idempotent** if all of its basic operations are idempotent.

Remarks

Idempotent Algebras

Definition

- An operation $f(x_1, x_2, \dots, x_n)$ on a set A is **idempotent** if for all $a \in A$, $f(a, a, \dots, a) = a$.
- An algebra is **idempotent** if all of its basic operations are idempotent.

Remarks

- *Many of the properties of the variety generated by an algebra are determined by the clone of idempotent term operations of the algebra.*

Idempotent Algebras

Definition

- An operation $f(x_1, x_2, \dots, x_n)$ on a set A is **idempotent** if for all $a \in A$, $f(a, a, \dots, a) = a$.
- An algebra is **idempotent** if all of its basic operations are idempotent.

Remarks

- *Many of the properties of the variety generated by an algebra are determined by the clone of idempotent term operations of the algebra.*
- *In general, it is hard to determine if a given finite algebra has any non-trivial idempotent term operations.*

Idempotent Algebras

Definition

- An operation $f(x_1, x_2, \dots, x_n)$ on a set A is **idempotent** if for all $a \in A$, $f(a, a, \dots, a) = a$.
- An algebra is **idempotent** if all of its basic operations are idempotent.

Remarks

- *Many of the properties of the variety generated by an algebra are determined by the clone of idempotent term operations of the algebra.*
- *In general, it is hard to determine if a given finite algebra has any non-trivial idempotent term operations.*

Fact

Płonka showed that the clone of idempotent operations of a group $(G, \cdot, {}^{-1}, e)$ is generated by the two idempotent ternary operations $x^{-1}yz$ and xyz^{-1} .

The Idempotent case

Theorem (Freese-Val.)

Let \mathbf{A} be a finite idempotent algebra. The following problems can be solved by algorithms whose run time can be bounded by a polynomial in the size of \mathbf{A} .

The Idempotent case

Theorem (Freese-Val.)

Let \mathbf{A} be a finite idempotent algebra. The following problems can be solved by algorithms whose run time can be bounded by a polynomial in the size of \mathbf{A} .

- *Is \mathbf{A} a $CD(k)$ algebra for some $k > 1$?*

The Idempotent case

Theorem (Freese-Val.)

Let \mathbf{A} be a finite idempotent algebra. The following problems can be solved by algorithms whose run time can be bounded by a polynomial in the size of \mathbf{A} .

- *Is \mathbf{A} a $CD(k)$ algebra for some $k > 1$?*
- *Is \mathbf{A} a $CD(2)$ algebra?*

The Idempotent case

Theorem (Freese-Val.)

Let \mathbf{A} be a finite idempotent algebra. The following problems can be solved by algorithms whose run time can be bounded by a polynomial in the size of \mathbf{A} .

- Is \mathbf{A} a $CD(k)$ algebra for some $k > 1$?
- Is \mathbf{A} a $CD(2)$ algebra?

Idea of proof

Take an algebra of minimal size in $\mathbf{V}(\mathbf{A})$ whose congruence lattice is not distributive and hammer away at it until it can be shown to lie in $\mathbf{HS}(\mathbf{A}^3)$.

The Idempotent case

Theorem (Freese-Val.)

Let \mathbf{A} be a finite idempotent algebra. The following problems can be solved by algorithms whose run time can be bounded by a polynomial in the size of \mathbf{A} .

- Is \mathbf{A} a $CD(k)$ algebra for some $k > 1$?
- Is \mathbf{A} a $CD(2)$ algebra?

Idea of proof

Take an algebra of minimal size in $\mathbf{V}(\mathbf{A})$ whose congruence lattice is not distributive and hammer away at it until it can be shown to lie in $\mathbf{HS}(\mathbf{A}^3)$.

The Idempotent case

Theorem (Freese-Val.)

Let \mathbf{A} be a finite idempotent algebra. The following problems can be solved by algorithms whose run time can be bounded by a polynomial in the size of \mathbf{A} .

- *Is \mathbf{A} a $CD(k)$ algebra for some $k > 1$?*
- *Is \mathbf{A} a $CD(2)$ algebra?*

Idea of proof

Take an algebra of minimal size in $\mathbf{V}(\mathbf{A})$ whose congruence lattice is not distributive and hammer away at it until it can be shown to lie in $\mathbf{HS}(\mathbf{A}^3)$. Our algorithm amounts to searching through the 3-generated subalgebras of \mathbf{A}^3 , looking for a failure of congruence distributivity.

A different (better?) approach

The CD(2) case

Let \mathbf{A} be a finite idempotent algebra.

A different (better?) approach

The CD(2) case

Let \mathbf{A} be a finite idempotent algebra.

- To determine if \mathbf{A} has a majority term one needs to find a term operation $m(x, y, z)$ of \mathbf{A} such that for all 6-tuples $\sigma = (a_1, a_2, a_3, b_1, b_2, b_3) \in A^6$,

$$m(a_1, a_1, b_1) = a_1, \quad m(a_2, b_2, a_2) = a_2, \quad \text{and} \quad m(b_3, a_3, a_3) = a_3.$$

A different (better?) approach

The CD(2) case

Let \mathbf{A} be a finite idempotent algebra.

- To determine if \mathbf{A} has a majority term one needs to find a term operation $m(x, y, z)$ of \mathbf{A} such that for all 6-tuples $\sigma = (a_1, a_2, a_3, b_1, b_2, b_3) \in A^6$,

$$m(a_1, a_1, b_1) = a_1, \quad m(a_2, b_2, a_2) = a_2, \quad \text{and} \quad m(b_3, a_3, a_3) = a_3.$$

- Independently, Horowitz and McKenzie observed that \mathbf{A} will have a majority term if and only if for every 6-tuple σ as above, \mathbf{A} has a term operation $m^\sigma(x, y, z)$ that satisfies the displayed equalities.

A different (better?) approach

The CD(2) case

Let \mathbf{A} be a finite idempotent algebra.

- To determine if \mathbf{A} has a majority term one needs to find a term operation $m(x, y, z)$ of \mathbf{A} such that for all 6-tuples $\sigma = (a_1, a_2, a_3, b_1, b_2, b_3) \in A^6$,

$$m(a_1, a_1, b_1) = a_1, \quad m(a_2, b_2, a_2) = a_2, \quad \text{and} \quad m(b_3, a_3, a_3) = a_3.$$

- Independently, Horowitz and McKenzie observed that \mathbf{A} will have a majority term if and only if for every 6-tuple σ as above, \mathbf{A} has a term operation $m^\sigma(x, y, z)$ that satisfies the displayed equalities.
- Thus, if \mathbf{A} has enough “local” majority term operations, then it will have a majority term operation.

Testing for local term operations

Proposition

For \mathbf{A} an algebra and $\sigma = (a_1, a_2, a_3, b_1, b_2, b_3) \in A^6$, \mathbf{A} will have a term operation m^σ satisfying

$$m^\sigma(a_1, a_1, b_1) = a_1, \quad m^\sigma(a_2, b_2, a_2) = a_2, \quad \text{and} \quad m^\sigma(b_3, a_3, a_3) = a_3$$

if and only if the tuple (a_1, a_2, a_3) is in the subuniverse of \mathbf{A}^3 generated by

$$\{(a_1, a_2, b_3), (a_1, b_2, a_3), (b_1, a_2, a_3)\}.$$

Testing for local term operations

Proposition

For \mathbf{A} an algebra and $\sigma = (a_1, a_2, a_3, b_1, b_2, b_3) \in A^6$, \mathbf{A} will have a term operation m^σ satisfying

$$m^\sigma(a_1, a_1, b_1) = a_1, \quad m^\sigma(a_2, b_2, a_2) = a_2, \quad \text{and} \quad m^\sigma(b_3, a_3, a_3) = a_3$$

if and only if the tuple (a_1, a_2, a_3) is in the subuniverse of \mathbf{A}^3 generated by

$$\{(a_1, a_2, b_3), (a_1, b_2, a_3), (b_1, a_2, a_3)\}.$$

Remark

It follows that to test that a finite idempotent algebra \mathbf{A} has a majority term, one need only search through A^6 , checking that the above subuniverse condition holds.

Testing for local term operations

Proposition

For \mathbf{A} an algebra and $\sigma = (a_1, a_2, a_3, b_1, b_2, b_3) \in A^6$, \mathbf{A} will have a term operation m^σ satisfying

$$m^\sigma(a_1, a_1, b_1) = a_1, \quad m^\sigma(a_2, b_2, a_2) = a_2, \quad \text{and} \quad m^\sigma(b_3, a_3, a_3) = a_3$$

if and only if the tuple (a_1, a_2, a_3) is in the subuniverse of \mathbf{A}^3 generated by

$$\{(a_1, a_2, b_3), (a_1, b_2, a_3), (b_1, a_2, a_3)\}.$$

Remark

It follows that to test that a finite idempotent algebra \mathbf{A} has a majority term, one need only search through A^6 , checking that the above subuniverse condition holds.

Testing for local term operations

Proposition

For \mathbf{A} an algebra and $\sigma = (a_1, a_2, a_3, b_1, b_2, b_3) \in A^6$, \mathbf{A} will have a term operation m^σ satisfying

$$m^\sigma(a_1, a_1, b_1) = a_1, \quad m^\sigma(a_2, b_2, a_2) = a_2, \quad \text{and} \quad m^\sigma(b_3, a_3, a_3) = a_3$$

if and only if the tuple (a_1, a_2, a_3) is in the subuniverse of \mathbf{A}^3 generated by

$$\{(a_1, a_2, b_3), (a_1, b_2, a_3), (b_1, a_2, a_3)\}.$$

Remark

It follows that to test that a finite idempotent algebra \mathbf{A} has a majority term, one need only search through A^6 , checking that the above subuniverse condition holds.

This can be done using a poly-time algorithm, since this sort of subuniverse generation can be performed quickly.

Definition

Let $k > 1$ and \mathbf{A} be a finite idempotent algebra and let

$$\sigma = (a_1, a_2, \dots, a_k, b_1, b_2, \dots, b_k, c_1, c_2, \dots, c_{k-1}, d_1, \dots, d_{k-1})$$

be a tuple from A^{4k-2} . The sequence of term operations $p_i^\sigma(x, y, z)$, $1 \leq i < k$ is a local Jónsson sequence for σ if

$$p_i^\sigma(c_i, d_i, c_i) = c_i \text{ for all } i, \text{ and}$$

$$a_1 = p_1^\sigma(a_1, a_1, b_1),$$

$$p_1^\sigma(a_2, b_2, b_2) = p_2^\sigma(a_2, b_2, b_2),$$

$$p_2^\sigma(a_3, a_3, b_3) = p_3^\sigma(a_3, a_3, b_3),$$

$$\vdots$$

CD(k) for $k > 2$

Theorem (Kazda-Val.)

Let \mathbf{A} be a finite idempotent algebra and $k > 1$. \mathbf{A} is a CD(k) algebra if and only if for every $4k - 2$ tuple σ from A , \mathbf{A} has a local Jónsson sequence for σ .

CD(k) for $k > 2$

Theorem (Kazda-Val.)

Let \mathbf{A} be a finite idempotent algebra and $k > 1$. \mathbf{A} is a CD(k) algebra if and only if for every $4k - 2$ tuple σ from A , \mathbf{A} has a local Jónsson sequence for σ .

Corollary

For $k > 1$, there is a poly-time algorithm to determine if a given finite idempotent algebra is CD(k).

CD(k) for $k > 2$

Theorem (Kazda-Val.)

Let \mathbf{A} be a finite idempotent algebra and $k > 1$. \mathbf{A} is a CD(k) algebra if and only if for every $4k - 2$ tuple σ from A , \mathbf{A} has a local Jónsson sequence for σ .

Corollary

For $k > 1$, there is a poly-time algorithm to determine if a given finite idempotent algebra is CD(k).

Remarks

CD(k) for $k > 2$

Theorem (Kazda-Val.)

Let \mathbf{A} be a finite idempotent algebra and $k > 1$. \mathbf{A} is a CD(k) algebra if and only if for every $4k - 2$ tuple σ from A , \mathbf{A} has a local Jónsson sequence for σ .

Corollary

For $k > 1$, there is a poly-time algorithm to determine if a given finite idempotent algebra is CD(k).

Remarks

- *The proof of the theorem is in the Krakow-Prague style and uses potatoes and coloured graphs.*

CD(k) for $k > 2$

Theorem (Kazda-Val.)

Let \mathbf{A} be a finite idempotent algebra and $k > 1$. \mathbf{A} is a CD(k) algebra if and only if for every $4k - 2$ tuple σ from A , \mathbf{A} has a local Jónsson sequence for σ .

Corollary

For $k > 1$, there is a poly-time algorithm to determine if a given finite idempotent algebra is CD(k).

Remarks

- The proof of the theorem is in the Krakow-Prague style and uses potatoes and coloured graphs.
- The corollary follows, since checking for local Jónsson sequences amounts to computing certain 3-generated subuniverses of \mathbf{A}^3 and checking for the existence of special elements.



Definition

An operation $t(x_1, x_2, \dots, x_n)$ is a **cyclic operation** on the set A if it is idempotent and for all $(a_1, a_2, \dots, a_n) \in A^n$,

$$t(a_1, a_2, \dots, a_n) = t(a_2, a_3, \dots, a_n, a_1) = \dots = t(a_n, a_1, \dots, a_{n-1}).$$

Cyclic terms

Definition

An operation $t(x_1, x_2, \dots, x_n)$ is a **cyclic operation** on the set A if it is idempotent and for all $(a_1, a_2, \dots, a_n) \in A^n$,

$$t(a_1, a_2, \dots, a_n) = t(a_2, a_3, \dots, a_n, a_1) = \dots = t(a_n, a_1, \dots, a_{n-1}).$$

Theorem (BKMMN)

Let $n > 1$. A finite idempotent algebra has a cyclic term operation of arity n if and only if it has enough local cyclic term operations.

Cyclic terms

Definition

An operation $t(x_1, x_2, \dots, x_n)$ is a **cyclic operation** on the set A if it is idempotent and for all $(a_1, a_2, \dots, a_n) \in A^n$,

$$t(a_1, a_2, \dots, a_n) = t(a_2, a_3, \dots, a_n, a_1) = \dots = t(a_n, a_1, \dots, a_{n-1}).$$

Theorem (BKMMN)

Let $n > 1$. A finite idempotent algebra has a cyclic term operation of arity n if and only if it has enough local cyclic term operations.

Corollary

For $n > 1$, there is a poly-time algorithm to determine if a given finite idempotent algebra has a cyclic term operation of arity n .

Cyclic terms for groups

Remarks

Cyclic terms for groups

Remarks

- *BKMMN showed that if \mathbf{A} is any finite algebra that generates a congruence modular variety, then for all prime numbers $p > |A|$, \mathbf{A} will have a p -ary cyclic term operation.*

Cyclic terms for groups

Remarks

- *BKMMN showed that if \mathbf{A} is any finite algebra that generates a congruence modular variety, then for all prime numbers $p > |A|$, \mathbf{A} will have a p -ary cyclic term operation.*
- *Barto and Kozik extended this result to any finite algebra that generates a variety that satisfies some non-trivial idempotent Maltsev condition.*

Cyclic terms for groups

Remarks

- *BKMMN showed that if \mathbf{A} is any finite algebra that generates a congruence modular variety, then for all prime numbers $p > |A|$, \mathbf{A} will have a p -ary cyclic term operation.*
- *Barto and Kozik extended this result to any finite algebra that generates a variety that satisfies some non-trivial idempotent Maltsev condition.*

Barto's 4 beers & 1 shot problem

Does the group S_5 have a cyclic term operation of arity less than 61?

Cyclic terms for groups

Remarks

- *BKMMN showed that if \mathbf{A} is any finite algebra that generates a congruence modular variety, then for all prime numbers $p > |A|$, \mathbf{A} will have a p -ary cyclic term operation.*
- *Barto and Kozik extended this result to any finite algebra that generates a variety that satisfies some non-trivial idempotent Maltsev condition.*

Barto's 4 beers & 1 shot problem

Does the group S_5 have a cyclic term operation of arity less than 61?

Theorem (Lai-Val.)

If \mathbf{G} is a finite group and $n > 1$, then \mathbf{G} will have a cyclic term operation of arity n if n and $|G|$ are relatively prime.

A conjecture

Conjecture

Let Σ be a *special Maltsev condition* (i.e., it is strong, idempotent and does not involve compositions of terms). There is a poly-time algorithm to determine, given a finite idempotent algebra \mathbf{A} , if $\mathbf{V}(\mathbf{A})$ satisfies Σ .

A conjecture

Conjecture

Let Σ be a *special Maltsev condition* (i.e., it is strong, idempotent and does not involve compositions of terms). There is a poly-time algorithm to determine, given a finite idempotent algebra \mathbf{A} , if $\mathbf{V}(\mathbf{A})$ satisfies Σ .

Remarks

A conjecture

Conjecture

Let Σ be a *special Maltsev condition* (i.e., it is strong, idempotent and does not involve compositions of terms). There is a poly-time algorithm to determine, given a finite idempotent algebra \mathbf{A} , if $\mathbf{V}(\mathbf{A})$ satisfies Σ .

Remarks

- Beside verifying the conjecture for the $CD(k)$ condition, it has also been verified for variants of it.

A conjecture

Conjecture

Let Σ be a *special Maltsev condition* (i.e., it is strong, idempotent and does not involve compositions of terms). There is a poly-time algorithm to determine, given a finite idempotent algebra \mathbf{A} , if $\mathbf{V}(\mathbf{A})$ satisfies Σ .

Remarks

- Beside verifying the conjecture for the $CD(k)$ condition, it has also been verified for variants of it.
- Valeriote and Willard have verified it for congruence n -permutability for any given $n > 1$.

A conjecture

Conjecture

Let Σ be a *special Maltsev condition* (i.e., it is strong, idempotent and does not involve compositions of terms). There is a poly-time algorithm to determine, given a finite idempotent algebra \mathbf{A} , if $\mathbf{V}(\mathbf{A})$ satisfies Σ .

Remarks

- Beside verifying the conjecture for the $CD(k)$ condition, it has also been verified for variants of it.
- Valeriote and Willard have verified it for congruence n -permutability for any given $n > 1$.
- Horowitz has verified it for k -edge terms and other types of cube terms.

A conjecture

Conjecture

Let Σ be a *special Maltsev condition* (i.e., it is strong, idempotent and does not involve compositions of terms). There is a poly-time algorithm to determine, given a finite idempotent algebra \mathbf{A} , if $\mathbf{V}(\mathbf{A})$ satisfies Σ .

Remarks

- Beside verifying the conjecture for the $CD(k)$ condition, it has also been verified for variants of it.
- Valeriote and Willard have verified it for congruence n -permutability for any given $n > 1$.
- Horowitz has verified it for k -edge terms and other types of cube terms.
- It also holds for Gumm terms of a fixed length.

Related conjectures & questions

Remark

A stronger version of the conjecture is that not only can special Maltsev conditions be tested in poly-time for finite idempotent algebras, but that this can be accomplished via checking for the existence of associated local term operations.

Related conjectures & questions

Remark

A stronger version of the conjecture is that not only can special Maltsev conditions be tested in poly-time for finite idempotent algebras, but that this can be accomplished via checking for the existence of associated local term operations.

Questions/Problems

Related conjectures & questions

Remark

A stronger version of the conjecture is that not only can special Maltsev conditions be tested in poly-time for finite idempotent algebras, but that this can be accomplished via checking for the existence of associated local term operations.

Questions/Problems

- Is it the case that if Σ is a strong idempotent Maltsev condition that is not equivalent to a special Maltsev condition, then testing for Σ is an EXP-TIME complete problem, even for idempotent algebras?

Related conjectures & questions

Remark

A stronger version of the conjecture is that not only can special Maltsev conditions be tested in poly-time for finite idempotent algebras, but that this can be accomplished via checking for the existence of associated local term operations.

Questions/Problems

- Is it the case that if Σ is a strong idempotent Maltsev condition that is not equivalent to a special Maltsev condition, then testing for Σ is an EXP-TIME complete problem, even for idempotent algebras?
- How hard is it to test if a finite idempotent algebra has a semilattice term operation?

Related conjectures & questions

Remark

A stronger version of the conjecture is that not only can special Maltsev conditions be tested in poly-time for finite idempotent algebras, but that this can be accomplished via checking for the existence of associated local term operations.

Questions/Problems

- Is it the case that if Σ is a strong idempotent Maltsev condition that is not equivalent to a special Maltsev condition, then testing for Σ is an EXP-TIME complete problem, even for idempotent algebras?
- How hard is it to test if a finite idempotent algebra has a semilattice term operation?
- Show that if $n > 1$, then there is a poly-time algorithm to test whether a given finite idempotent algebra has an n -ary weak near unanimity term operation.