

Term-equivalent semigroups

Oleg Košík¹ Peter Mayr²

¹University of Tartu

²JKU Linz

ALC fest 2014, Prague

Term-equivalence

A *term operation* of an algebra \mathbf{A} is any operation on A that can be constructed from basic operations and projection maps.

Term-equivalence

A *term operation* of an algebra \mathbf{A} is any operation on A that can be constructed from basic operations and projection maps.

Two algebras with the same universe are called *term-equivalent* if their clones of term operations are equal.

Term-equivalence

A *term operation* of an algebra **A** is any operation on A that can be constructed from basic operations and projection maps.

Two algebras with the same universe are called *term-equivalent* if their clones of term operations are equal.

Example

$(\mathbb{Z}_n, +)$ and $(\mathbb{Z}_n, -)$ are term-equivalent.

Term-equivalence

A *term operation* of an algebra **A** is any operation on A that can be constructed from basic operations and projection maps.

Two algebras with the same universe are called *term-equivalent* if their clones of term operations are equal.

Example

$(\mathbb{Z}_n, +)$ and $(\mathbb{Z}_n, -)$ are term-equivalent.

Indeed, $x - y = x + (n - 1)y$ and $x + y = x - ((y - y) - y)$.

Motivation

Motivation

Theorem (L. Zádori, 1997)

Finite groups are categorically equivalent iff they are weakly isomorphic.

Two algebras **A** and **B** are called weakly isomorphic, if there exists an isomorphism that maps **A** to an algebra term-equivalent to **B**.

Motivation

Theorem (L. Zádori, 1997)

Finite groups are categorically equivalent iff they are weakly isomorphic.

Two algebras **A** and **B** are called weakly isomorphic, if there exists an isomorphism that maps **A** to an algebra term-equivalent to **B**.

Problem (L. Zádori, 1997)

Are term-equivalent finite groups isomorphic?

Motivation

Theorem (L. Zádori, 1997)

Finite groups are categorically equivalent iff they are weakly isomorphic.

Two algebras **A** and **B** are called weakly isomorphic, if there exists an isomorphism that maps **A** to an algebra term-equivalent to **B**.

Problem (L. Zádori, 1997)

Are term-equivalent finite groups isomorphic?

However, already in 1967, A. P. Street constructed two non-isomorphic term-equivalent groups of the smallest order $2 \cdot 7 \cdot 43$.

Motivation

Theorem (L. Zádori, 1997)

Finite groups are categorically equivalent iff they are weakly isomorphic.

Two algebras **A** and **B** are called weakly isomorphic, if there exists an isomorphism that maps **A** to an algebra term-equivalent to **B**.

Problem (L. Zádori, 1997)

Are term-equivalent finite groups isomorphic?

However, already in 1967, A. P. Street constructed two non-isomorphic term-equivalent groups of the smallest order $2 \cdot 7 \cdot 43$.

In 2004, K. A. Kearnes and Á. Szendrei constructed two groups of the form $(\mathbb{Z}_7 \times \mathbb{Z}_{13}) \rtimes \mathbb{Z}_3$ that are term equivalent but not isomorphic.

Motivation

Theorem (L. Zádori, 1997)

Finite groups are categorically equivalent iff they are weakly isomorphic.

Two algebras **A** and **B** are called weakly isomorphic, if there exists an isomorphism that maps **A** to an algebra term-equivalent to **B**.

Problem (L. Zádori, 1997)

Are term-equivalent finite groups isomorphic?

However, already in 1967, A. P. Street constructed two non-isomorphic term-equivalent groups of the smallest order $2 \cdot 7 \cdot 43$.

In 2004, K. A. Kearnes and Á. Szendrei constructed two groups of the form $(\mathbb{Z}_7 \times \mathbb{Z}_{13}) \rtimes \mathbb{Z}_3$ that are term equivalent but not isomorphic.

Theorem (M. Behrisch, T. Waldhauser)

Finite semigroups are categorically equivalent iff they are weakly isomorphic.

First observations

Let $\langle S, \cdot \rangle$ be a semigroup and $a \in S$. By $\langle a \rangle$ we denote the subgroup generated by a .

First observations

Let $\langle S, \cdot \rangle$ be a semigroup and $a \in S$. By $\langle a \rangle$ we denote the subgroup generated by a . The *index* of $\langle a \rangle$ is the smallest m for which $a^m = a^r$ for some positive integer r .

First observations

Let $\langle S, \cdot \rangle$ be a semigroup and $a \in S$. By $\langle a \rangle$ we denote the subgroup generated by a . The *index* of $\langle a \rangle$ is the smallest m for which $a^m = a^r$ for some positive integer r . The *period* of $\langle a \rangle$ is the smallest n for which $a^{m+n} = a^m$ for some positive integer m .

First observations

Let $\langle S, \cdot \rangle$ be a semigroup and $a \in S$. By $\langle a \rangle$ we denote the subgroup generated by a . The *index* of $\langle a \rangle$ is the smallest m for which $a^m = a^r$ for some positive integer r . The *period* of $\langle a \rangle$ is the smallest n for which $a^{m+n} = a^m$ for some positive integer m . If $\langle a \rangle$ is infinite, then both index and period are infinite.

First observations

Let $\langle S, \cdot \rangle$ be a semigroup and $a \in S$. By $\langle a \rangle$ we denote the subgroup generated by a . The *index* of $\langle a \rangle$ is the smallest m for which $a^m = a^r$ for some positive integer r . The *period* of $\langle a \rangle$ is the smallest n for which $a^{m+n} = a^m$ for some positive integer m . If $\langle a \rangle$ is infinite, then both index and period are infinite.

Lemma

Let (A, \cdot) and $(A, +)$ be term equivalent semigroups. Then

First observations

Let $\langle S, \cdot \rangle$ be a semigroup and $a \in S$. By $\langle a \rangle$ we denote the subgroup generated by a . The *index* of $\langle a \rangle$ is the smallest m for which $a^m = a^r$ for some positive integer r . The *period* of $\langle a \rangle$ is the smallest n for which $a^{m+n} = a^m$ for some positive integer m . If $\langle a \rangle$ is infinite, then both index and period are infinite.

Lemma

Let (A, \cdot) and $(A, +)$ be term equivalent semigroups. Then

- The subsemigroups of (A, \cdot) and $(A, +)$ are the same and term equivalent.*

First observations

Let $\langle S, \cdot \rangle$ be a semigroup and $a \in S$. By $\langle a \rangle$ we denote the subgroup generated by a . The *index* of $\langle a \rangle$ is the smallest m for which $a^m = a^r$ for some positive integer r . The *period* of $\langle a \rangle$ is the smallest n for which $a^{m+n} = a^m$ for some positive integer m . If $\langle a \rangle$ is infinite, then both index and period are infinite.

Lemma

Let (A, \cdot) and $(A, +)$ be term equivalent semigroups. Then

- The subsemigroups of (A, \cdot) and $(A, +)$ are the same and term equivalent.*
- The ideals of (A, \cdot) and $(A, +)$ are the same.*

First observations

Let $\langle S, \cdot \rangle$ be a semigroup and $a \in S$. By $\langle a \rangle$ we denote the subgroup generated by a . The *index* of $\langle a \rangle$ is the smallest m for which $a^m = a^r$ for some positive integer r . The *period* of $\langle a \rangle$ is the smallest n for which $a^{m+n} = a^m$ for some positive integer m . If $\langle a \rangle$ is infinite, then both index and period are infinite.

Lemma

Let (A, \cdot) and $(A, +)$ be term equivalent semigroups. Then

- The subsemigroups of (A, \cdot) and $(A, +)$ are the same and term equivalent.*
- The ideals of (A, \cdot) and $(A, +)$ are the same.*
- The idempotents of (A, \cdot) and $(A, +)$ are the same.*

First observations

Let $\langle S, \cdot \rangle$ be a semigroup and $a \in S$. By $\langle a \rangle$ we denote the subgroup generated by a . The *index* of $\langle a \rangle$ is the smallest m for which $a^m = a^r$ for some positive integer r . The *period* of $\langle a \rangle$ is the smallest n for which $a^{m+n} = a^m$ for some positive integer m . If $\langle a \rangle$ is infinite, then both index and period are infinite.

Lemma

Let (A, \cdot) and $(A, +)$ be term equivalent semigroups. Then

- The subsemigroups of (A, \cdot) and $(A, +)$ are the same and term equivalent.*
- The ideals of (A, \cdot) and $(A, +)$ are the same.*
- The idempotents of (A, \cdot) and $(A, +)$ are the same.*
- For every $a \in A$ that generates a finite semigroup $\langle a \rangle$, the index $m(a)$ and the period $n(a)$ of $\langle a \rangle$ are the same with respect to \cdot and $+$.*

More observations

Lemma

Let (S, \cdot) and $(S, +)$ be term equivalent semigroups. Assume that $+$ does not depend on both arguments. Then $xy = x + y$ for all $x, y \in S$ or $xy = y + x$ for all $x, y \in S$.

More observations

Lemma

Let (S, \cdot) and $(S, +)$ be term equivalent semigroups. Assume that $+$ does not depend on both arguments. Then $xy = x + y$ for all $x, y \in S$ or $xy = y + x$ for all $x, y \in S$.

Lemma

Let (S, \cdot) and $(S, +)$ be term equivalent semigroups. Assume that for all $n \in \mathbb{N}$ there exists $a \in S$ such that $|\langle a \rangle| > n$ or a has index at least 3. Then $xy = x + y$ for all $x, y \in S$ or $xy = y + x$ for all $x, y \in S$.

By this lemma, two semigroups may be term equivalent without being identical or dual only if every element is contained in a group or has index 2. The order of every element is bounded from above by some fixed positive integer.

Nilpotent semigroups

Corollary

If a semigroup $(S, +)$ is term equivalent to a nilpotent semigroup (S, \cdot) , then both are identical or dual to each other.

Nilpotent semigroups

Corollary

If a semigroup $(S, +)$ is term equivalent to a nilpotent semigroup (S, \cdot) , then both are identical or dual to each other.

Proof. Let $s(x, y)$ be a term over $(S, +)$, let $t(x, y)$ be a term over (S, \cdot) such that

$$xy = s(x, y) \text{ and } x + y = t(x, y) \text{ for all } x, y \in S.$$

Assume that lengths of s and t are at least 3. Then we obtain

$$S^2 \subseteq S + S \subseteq S^3 \subseteq S^2$$

Thus $S^2 = S^3$ and consequently $S^2 = S^n$ for every $n \geq 3$. Since (S, \cdot) is nilpotent, it follows that $S^2 = 0$. Thus (S, \cdot) and $(S, +)$ are identical 0-semigroups.

Commutative semigroups

Theorem

If a semigroup (S, \cdot) is term equivalent to a commutative semigroup $(S, +)$, then both are identical.

Commutative semigroups

Theorem

If a semigroup (S, \cdot) is term equivalent to a commutative semigroup $(S, +)$, then both are identical.

Corollary

Let $(S, \cdot), (S, +)$ be term equivalent semigroups. Then for every $k \in \mathbb{N}$ and for every $x \in S$ we have $x^k = kx$.

Commutative semigroups

Theorem

If a semigroup (S, \cdot) is term equivalent to a commutative semigroup $(S, +)$, then both are identical.

Corollary

Let $(S, \cdot), (S, +)$ be term equivalent semigroups. Then for every $k \in \mathbb{N}$ and for every $x \in S$ we have $x^k = kx$.

Proof. Monogenic subsemigroups are commutative.

Some invariant properties

Theorem

*Let $(S, \cdot), (S, *)$ be term equivalent semigroups. Then we have:*

Some invariant properties

Theorem

*Let $(S, \cdot), (S, *)$ be term equivalent semigroups. Then we have:*

- 1 *Assume (S, \cdot) has a zero 0 . Then 0 is also a zero with respect to $*$.*

Some invariant properties

Theorem

*Let $(S, \cdot), (S, *)$ be term equivalent semigroups. Then we have:*

- ① *Assume (S, \cdot) has a zero 0 . Then 0 is also a zero with respect to $*$.*
- ② *Assume (S, \cdot) has an identity 1 . Then 1 is also an identity with respect to $*$, and (S, \cdot) and $(S, *)$ are identical, dual, or both completely regular (i.e. are union of groups).*

Some invariant properties

Theorem

*Let $(S, \cdot), (S, *)$ be term equivalent semigroups. Then we have:*

- ① *Assume (S, \cdot) has a zero 0 . Then 0 is also a zero with respect to $*$.*
- ② *Assume (S, \cdot) has an identity 1 . Then 1 is also an identity with respect to $*$, and (S, \cdot) and $(S, *)$ are identical, dual, or both completely regular (i.e. are union of groups).*
- ③ *Assume (S, \cdot) is finitely generated. Then $(S, *)$ is finitely generated.*

Some invariant properties

Theorem

*Let $(S, \cdot), (S, *)$ be term equivalent semigroups. Then we have:*

- ① *Assume (S, \cdot) has a zero 0 . Then 0 is also a zero with respect to $*$.*
- ② *Assume (S, \cdot) has an identity 1 . Then 1 is also an identity with respect to $*$, and (S, \cdot) and $(S, *)$ are identical, dual, or both completely regular (i.e. are union of groups).*
- ③ *Assume (S, \cdot) is finitely generated. Then $(S, *)$ is finitely generated.*
- ④ *Assume (S, \cdot) is regular. Then $(S, *)$ is regular.*

Some invariant properties

Theorem

*Let $(S, \cdot), (S, *)$ be term equivalent semigroups. Then we have:*

- ① *Assume (S, \cdot) has a zero 0 . Then 0 is also a zero with respect to $*$.*
- ② *Assume (S, \cdot) has an identity 1 . Then 1 is also an identity with respect to $*$, and (S, \cdot) and $(S, *)$ are identical, dual, or both completely regular (i.e. are union of groups).*
- ③ *Assume (S, \cdot) is finitely generated. Then $(S, *)$ is finitely generated.*
- ④ *Assume (S, \cdot) is regular. Then $(S, *)$ is regular.*
- ⑤ *Assume (S, \cdot) is inverse. Then $(S, *)$ is inverse.*

Some invariant properties

Theorem

*Let $(S, \cdot), (S, *)$ be term equivalent semigroups. Then we have:*

- ① *Assume (S, \cdot) has a zero 0 . Then 0 is also a zero with respect to $*$.*
- ② *Assume (S, \cdot) has an identity 1 . Then 1 is also an identity with respect to $*$, and (S, \cdot) and $(S, *)$ are identical, dual, or both completely regular (i.e. are union of groups).*
- ③ *Assume (S, \cdot) is finitely generated. Then $(S, *)$ is finitely generated.*
- ④ *Assume (S, \cdot) is regular. Then $(S, *)$ is regular.*
- ⑤ *Assume (S, \cdot) is inverse. Then $(S, *)$ is inverse.*
- ⑥ *Assume (S, \cdot) is a group. Then $(S, *)$ is a group.*

Term-equivalent semigroups that are not completely regular.

From the examples of term-equivalent groups we can obtain semigroups that are not regular and still term equivalent in a nontrivial way.

Term-equivalent semigroups that are not completely regular.

From the examples of term-equivalent groups we can obtain semigroups that are not regular and still term equivalent in a nontrivial way.

Example

Let $(G, \cdot), (G, *)$ be finite term equivalent, non-isomorphic groups with identity 1. Let a be distinct from the elements of G , and extend the operation $\circ \in \{\cdot, *\}$ to $S := G \cup \{a\}$ as follows:

$$\begin{aligned}x \circ a &= a \circ x = x \quad \forall x \in G, \\a \circ a &= 1.\end{aligned}$$

Then $(S, \cdot), (S, *)$ are term equivalent semigroups that are not regular, not isomorphic, and not dual to each other.

Completely regular semigroups

Problem

*Let $(S, *)$, (S, \cdot) be term equivalent completely regular semigroups such that $*$ and \cdot are identical on all subgroups. Are $(S, *)$ and (S, \cdot) identical or dual to each other?*

Completely regular semigroups

Problem

*Let $(S, *)$, (S, \cdot) be term equivalent completely regular semigroups such that $*$ and \cdot are identical on all subgroups. Are $(S, *)$ and (S, \cdot) identical or dual to each other?*

Generally, the answer to this question is negative.

Completely regular semigroups

Problem

*Let $(S, *)$, (S, \cdot) be term equivalent completely regular semigroups such that $*$ and \cdot are identical on all subgroups. Are $(S, *)$ and (S, \cdot) identical or dual to each other?*

Generally, the answer to this question is negative.

Example

Let G be a finite non-commutative group. Let $S = (G \times \mathbb{Z}_2)$ be an universe of semigroups (S, \cdot) and $(S, *)$, where

$$(g, a) \cdot (h, b) := (gh, b), \quad (g, a) * (h, b) := (gh, a).$$

Semigroups (S, \cdot) and $(S, *)$ are term-equivalent, since

$$x \cdot y = y^{|G|} * x * y, \quad x * y = x * y * x^{|G|}.$$

Completely regular semigroups

However we can give an affirmative answer for some special cases.

Completely regular semigroups

However we can give an affirmative answer for some special cases.

Theorem

If a semigroup $(S, +)$ is term equivalent to a band (S, \cdot) , then both are identical or dual to each other.

Completely regular semigroups

However we can give an affirmative answer for some special cases.

Theorem

If a semigroup $(S, +)$ is term equivalent to a band (S, \cdot) , then both are identical or dual to each other.

Proposition

*Let $(S, *)$, (S, \cdot) be completely regular and term equivalent inverse semigroups. Assume the operations are identical on all subgroups. Then $(S, *)$ and (S, \cdot) are identical.*

Thank you for your attention!



European Union
European Social Fund



Investing in your future