

Presentation for semigroups $x \approx x^k, k > 2$

Somnuek Worawiset (with Dr. Jörz Koppitz)

Department of Mathematics, Khon Kaen University, Thailand

July 3, 2014

Outline

The aim of this talk is to look at a construction of some semigroups satisfying the identity $x \approx x^3$.

Theorem

Let Y be a semilattice. For each $\alpha \in Y$ let S_α be a semigroup such that $S_\alpha \cap S_\beta = \emptyset$ if $\alpha \neq \beta$. Let $\alpha, \beta \in Y$ with $\beta \leq \alpha$, let $\{\phi_{\alpha,\beta} : S_\alpha \rightarrow S_\beta\}$ be a collection of homomorphisms such that

Theorem

Let Y be a semilattice. For each $\alpha \in Y$ let S_α be a semigroup such that $S_\alpha \cap S_\beta = \emptyset$ if $\alpha \neq \beta$. Let $\alpha, \beta \in Y$ with $\beta \leq \alpha$, let $\{\phi_{\alpha,\beta} : S_\alpha \rightarrow S_\beta\}$ be a collection of homomorphisms such that

- 1) if $\gamma \leq \beta \leq \alpha$ in Y then $\phi_{\alpha,\beta}\phi_{\beta,\gamma} = \phi_{\alpha,\gamma}$.*

Theorem

Let Y be a semilattice. For each $\alpha \in Y$ let S_α be a semigroup such that $S_\alpha \cap S_\beta = \emptyset$ if $\alpha \neq \beta$. Let $\alpha, \beta \in Y$ with $\beta \leq \alpha$, let $\{\phi_{\alpha,\beta} : S_\alpha \rightarrow S_\beta\}$ be a collection of homomorphisms such that

- 1) if $\gamma \leq \beta \leq \alpha$ in Y then $\phi_{\alpha,\beta}\phi_{\beta,\gamma} = \phi_{\alpha,\gamma}$.
- 2) for each $\alpha \in Y$ in $\phi_{\alpha,\alpha}$ is the identity on S_α .

Theorem

Let Y be a semilattice. For each $\alpha \in Y$ let S_α be a semigroup such that $S_\alpha \cap S_\beta = \emptyset$ if $\alpha \neq \beta$. Let $\alpha, \beta \in Y$ with $\beta \leq \alpha$, let

$\{\phi_{\alpha,\beta} : S_\alpha \rightarrow S_\beta\}$ be a collection of homomorphisms such that

1) if $\gamma \leq \beta \leq \alpha$ in Y then $\phi_{\alpha,\beta}\phi_{\beta,\gamma} = \phi_{\alpha,\gamma}$.

2) for each $\alpha \in Y$ in $\phi_{\alpha,\alpha}$ is the identity on S_α .

Then $S = \bigcup_{\alpha \in Y} S_\alpha$ is a semigroup where for $x \in S_\alpha, y \in S_\beta$ the product in S is given by

$$xy = (x\phi_{\alpha,\alpha\beta})(y\phi_{\beta,\alpha\beta}).$$

Theorem

Let Y be a semilattice. For each $\alpha \in Y$ let S_α be a semigroup such that $S_\alpha \cap S_\beta = \emptyset$ if $\alpha \neq \beta$. Let $\alpha, \beta \in Y$ with $\beta \leq \alpha$, let

$\{\phi_{\alpha,\beta} : S_\alpha \rightarrow S_\beta\}$ be a collection of homomorphisms such that

1) if $\gamma \leq \beta \leq \alpha$ in Y then $\phi_{\alpha,\beta}\phi_{\beta,\gamma} = \phi_{\alpha,\gamma}$.

2) for each $\alpha \in Y$ in $\phi_{\alpha,\alpha}$ is the identity on S_α .

Then $S = \bigcup_{\alpha \in Y} S_\alpha$ is a semigroup where for $x \in S_\alpha, y \in S_\beta$ the product in S is given by


$$xy = (x\phi_{\alpha,\alpha\beta})(y\phi_{\beta,\alpha\beta}).$$

If replace a semigroup S_α by a group G_α then it will be called a Clifford semigroup


Example




Example




Example




Example



Example




Example



$$0_\alpha * 1_\alpha = (0_\alpha)\varphi_{\alpha,\alpha}(1_\alpha)\varphi_{\alpha,\alpha} = 0_\alpha 1_\alpha = 1_\alpha$$

$$1_\alpha * 1_\beta = (1_\alpha)\varphi_{\alpha,\alpha\beta}(1_\beta)\varphi_{\beta,\alpha\beta} = (1_\alpha)\varphi_{\alpha,\beta}(1_\beta)\varphi_{\beta,\beta} = 0_\beta 1_\beta = 1_\beta$$

Example




$$0_\alpha * 1_\alpha = (0_\alpha)\varphi_{\alpha,\alpha}(1_\alpha)\varphi_{\alpha,\alpha} = 0_\alpha 1_\alpha$$

$$1_\alpha * 1_\beta = (1_\alpha)\varphi_{\alpha,\alpha\beta}(1_\beta)\varphi_{\beta,\alpha\beta} = (1_\alpha)\varphi_{\alpha,\beta}(1_\beta)\varphi_{\beta,\beta} = 0_\beta 1_\beta = 1_\beta$$

Thus $(S = \{0_\alpha, 1_\alpha, 0_\beta, 1_\beta\}, *)$ is a semigroup.

Example



$$\varphi_{\gamma,\alpha}\varphi_{\alpha,\nu} = \varphi_{\gamma,\beta}\varphi_{\beta,\nu}$$

$$(a_\gamma)\varphi_{\gamma,\alpha}\varphi_{\alpha,\nu} = (a_\gamma)\varphi_{\gamma,\beta}\varphi_{\beta,\nu}$$

We begin by describing an important construction.

Definition

Let S be a semigroup, I, Λ be nonempty sets, and P be a $\Lambda \times I$ matrix with entries $p_{\lambda,i}$ from S . Define

$$M = M(S; I, \Lambda; P)$$

with the multiplication

$$(i, s, \lambda)(j, t, \mu) = (i, sp_{\lambda,j}t, \mu)$$

Then M is a semigroup called a Rees matrix semigroup over S . This is an important technique in semigroup theory for constructing new semigroups from old ones

Definition

Let S be a semigroup. An element $a \in S$ is called completely regular if there exists $x \in S$ such that $a = axa$, $ax = xa$.

Definition

Let S be a semigroup. An element $a \in S$ is called completely regular if there exists $x \in S$ such that $a = axa$, $ax = xa$.

Definition

Let S be a semigroup. Then S is simple if it does not have any proper ideals.

Definition

Let S be a semigroup. An element $a \in S$ is called completely regular if there exists $x \in S$ such that $a = axa$, $ax = xa$.

Definition

Let S be a semigroup. Then S is simple if it does not have any proper ideals.

Definition

Let e be an idempotent in a semigroup S . Then e is a primitive Idempotent if for each idempotent f in S such that $f \leq e$ then $f = e$.

Clifford

Clifford proved that completely regular semigroups are semilattices of completely simple semigroups and conversely.

Clifford

Clifford proved that completely regular semigroups are semilattices of completely simple semigroups and conversely.

Lallement

Using Clifford's representation of completely regular semigroups reduced the problem of their structure to structure of completely simple semigroups and certain functions among them and their translation hulls.

Theorem (Lallement theorem)

Let Y be a semilattice. For each $\alpha \in Y$ let S_α be a completely simple semigroup such that $S_\alpha \cap S_\beta = \emptyset$ if $\alpha \neq \beta$. For each pair α, β , $\alpha \geq \beta$, let $\{\Phi_{\alpha,\beta} : S_\alpha \rightarrow \Omega(S_\beta)\}$ be a function satisfying the following conditions:

- 1) $\Phi_{\alpha,\alpha} : a \rightarrow \pi_\alpha$ ($a \in S_\alpha$),
- 2) $(S_\alpha \Phi_{\alpha,\alpha\beta})(S_\beta \Phi_{\beta,\alpha\beta}) \subseteq \Pi(S_{\alpha,\beta})$,
- 3) if $\alpha\beta > \gamma$ and $a \in S_\alpha$, $b \in S_\beta$, then

$$[(a\Phi_{\alpha,\alpha\beta})(b\Phi_{\beta,\alpha\beta})]\Phi_{\alpha\beta,\alpha\beta}^{-1}\Phi_{\alpha\beta,\gamma} = (a\Phi_{\alpha,\gamma})(b\Phi_{\beta,\gamma}).$$

On $S = \bigcup_{\alpha \in Y} S_\alpha$ define a multiplication $*$ by

$$a * b = [(a\Phi_{\alpha,\alpha\beta})(b\Phi_{\beta,\alpha\beta})]\Phi_{\alpha\beta,\alpha\beta}^{-1}\Phi_{\alpha\beta,\gamma} \quad (a \in S_\alpha, b \in S_\beta).$$

Then S is a completely regular semigroup. Conversely, every completely regular semigroup can be so constructed.

The problem is reduced to

- 1) the structure of completely simple semigroups,

The problem is reduced to

- 1) the structure of completely simple semigroups,
- 2) the structure of their translational hulls,

The problem is reduced to

- 1) the structure of completely simple semigroups,
- 2) the structure of their translational hulls,
- 3) a determination of the functions satisfying the above condition.

The problem is reduced to

- 1) the structure of completely simple semigroups,
- 2) the structure of their translational hulls,
- 3) a determination of the functions satisfying the above condition.

The problem 1) has been solved for example,

The problem is reduced to

- 1) the structure of completely simple semigroups,
- 2) the structure of their translational hulls,
- 3) a determination of the functions satisfying the above condition.

The problem 1) has been solved for example,

Theorem

(David Rees 1940) A semigroup is completely simple if and only if it is isomorphic to a Rees matrix semigroup over a group.

The problem is reduced to

- 1) the structure of completely simple semigroups,
- 2) the structure of their translational hulls,
- 3) a determination of the functions satisfying the above condition.

The problem 1) has been solved for example,

Theorem

(David Rees 1940) A semigroup is completely simple if and only if it is isomorphic to a Rees matrix semigroup over a group.

There are also many research on the generalizations of Rees theorem with applications to some other classes of semigroups.

The problem 2) has been quasi-solved in view of several constructions of the translational hull of a Rees matrix semigroup.

The problem 2) has been quasi-solved in view of several constructions of the translational hull of a Rees matrix semigroup. The problem 3) has no solution whatsoever except in very special cases.

The problem 2) has been quasi-solved in view of several constructions of the translational hull of a Rees matrix semigroup. The problem 3) has no solution whatsoever except in very special cases.

Theorem

Let Y be a semilattice. For each $\alpha \in Y$ let E_α be a rectangular band such that $E_\alpha \cap E_\beta = \emptyset$ if $\alpha \neq \beta$. For each $\alpha, \beta \in Y$ with $\beta \leq \alpha$, let $\{\phi_{\alpha,\beta} : E_\alpha \rightarrow E_\beta\}$ be a collection of homomorphisms such that

- 1) if $\gamma \leq \beta \leq \alpha$ in Y then $\phi_{\alpha,\beta}\phi_{\beta,\gamma} = \phi_{\alpha,\gamma}$.*
- 2) for each $\alpha \in Y$ in $\phi_{\alpha,\alpha}$ is the identity on E_α .*

Then $S = \bigcup_{\alpha \in Y} E_\alpha$ is a semigroup where for $x \in E_\alpha, y \in E_\beta$ the product in S is given by

$$xy = (x)(\phi_{\alpha,\alpha\beta})(y\phi_{\beta,\alpha\beta}).$$

Then S is a band and conversely.

Example

$$\text{Mod}[x(yz) \approx (xy)z, x \approx x^3, (xyz)^2 \approx (xz)^2] \quad \text{Mod}[x(yz) \approx (xy)z, x \approx x^3]$$

$$\text{Mod}[x(yz) \approx (xy)z, x \approx x^3, \\ (xyz)^2 \approx (xz)^2, xyzx \approx xzyx]$$

???????

Regular band [RB]

Bands

Lemma

Let $S \in \text{Mod}[(xy)z \approx x(yz), x \approx x^3, (xyz)^2 \approx (xz)^2, xyzx \approx xzyx]$.

Then

1) $(xyzx)^2 = xy^2x$

2) $E(S) = \{x^2 \mid x \in S\}$

3) S is completely simple

Proof. 1) Take $x, y, z \in S$.

$$\begin{aligned}(xyzx)^2 &= (xyzx)(xyzx) \\&= x(yzxx)yzx \\&= x(yz)(xxyz)x \\&= xyz(xy xz)x \\&= xyzx(yzx)x \\&= x(yzx)^2x \\&= x(yx)^2x \\&= xyxyxx \\&= x^3y^2x \\&= xy^2x\end{aligned}$$

2) Obviously, by $(x^2)^2 = x^3x = x^2$

3) By the property $x = x^3$ implies that S is regular. Now we consider for each $e, f \in E(S)$ such that $f = ef = fe$, then

$$\begin{aligned} ef &= fe \\ (ef)ef &= (fe)ef \\ (eeff) &= feef \\ efefe &= fefee \\ ef^2e^2 &= fe^3 \\ efe &= f^2e \\ (efe)^2 &= (fe)^2 \\ (ee)^2 &= f^2 \\ e &= f. \end{aligned}$$

Hence S is completely simple.

Proposition (Howie)

Let e be an idempotent of a semigroup S . Then

$$G_e = \{a \in S \mid a \in eS \cap Se, e \in aS \cap Sa\}$$

Proposition (Howie)

Let e be an idempotent of a semigroup S . Then

$$G_e = \{a \in S \mid a \in eS \cap Se, e \in aS \cap Sa\}$$

Lemma

Let $S \in \text{Mod}[x(yz) \approx (xy)z, x \approx x^3, (xyz)^2 \approx (xz)^2, xyzx \approx xzyx]$. Then the set

$$G_{x^2} = \{y \in S \mid y^2 = x^2\}$$

is an abelian group.

Lemma

Let $S \in \text{Mod}[x(yz) \approx (xy)z, x \approx x^3, (xyz)^2 \approx (xz)^2, xyzx \approx xzyx]$ and $x^2 \in E(S)$. Then the set

$$L_{x^2} = \{(ax^2)^2 \mid a \in S\}$$

is a left zero semigroup and $lx^2 = l$ for every $l \in L_{x^2}$.

Lemma

Let $S \in \text{Mod}[x(yz) \approx (xy)z, x \approx x^3, (xyz)^2 \approx (xz)^2, xyzx \approx xzyx]$ and $x^2 \in E(S)$. Then the set

$$L_{x^2} = \{(ax^2)^2 \mid a \in S\}$$

is a left zero semigroup and $lx^2 = l$ for every $l \in L_{x^2}$.

Lemma

Let $S \in \text{Mod}[x(yz) \approx (xy)z, x \approx x^3, (xyz)^2 \approx (xz)^2, xyzx \approx xzyx]$ and $x^2 \in E(S)$. Then the set

$$R_{x^2} = \{(x^2a)^2 \mid a \in S\}$$

is a right zero semigroup and $x^2r = r$ for every $r \in R_{x^2}$.

Theorem

Let $S \in \text{Mod}[x(yz) \approx (xy)z, x \approx x^3, (xyz)^2 \approx (xz)^2, xyzx \approx xzyx]$.
Then

$$S \cong M[G_{x^2}; L_{x^2}, R_{x^2}; P]$$

where $G_{x^2}, L_{x^2}, R_{x^2}$ are described above and P is a sandwich Rees Matrix.

Example

$$\text{Mod}[x(yz) \approx (xy)z, x \approx x^3, (xyz)^2 \approx (xz)^2] \quad \text{Mod}[x(yz) \approx (xy)z, x \approx x^3]$$

$$\text{Mod}[x(yz) \approx (xy)z, x \approx x^3, \\ (xyz)^2 \approx (xz)^2, xyzx \approx xzyx]$$


$$\text{Mod}[x(yz) \approx (xy)z, x \approx x^3, \\ xyzx \approx xzyx]$$

Regular band[RB]


Bands

Example

$$\text{Mod}[x(yz) \approx (xy)z, x \approx x^3, (xyz)^2 \approx (xz)^2] \quad \text{Mod}[x(yz) \approx (xy)z, x \approx x^3]$$



Rectangular band[RB]



Bands

Thank You !