

On Regular Ordered Ternary Semigroups

Nareupanat Lekkoksung

Department of Mathematics, Faculty of Science,
Khon Kaen University, Thailand

3 July 2014

Outline

- ① Motivation
- ② Introduction
- ③ Main Results
- ④ References

Outline

- ① Motivation
- ② Introduction
- ③ Main Results
- ④ References

Motivation

- In 1932 [10], D. H. Lehmer investigated certain ternary algebraic system called triplexes which turn out to be ternary groups.
- In 1965 [15], Zadeh introduced the concept of fuzzy sets. And F. M. Sioson studied ideal theory in ternary semigroups in the same year.
- In 1983 [12], M. L. Santiago developed the theory of ternary semigroups.
- In 1981, 1991 and 1993 [6, 7, 8], fuzzy semigroups have been first considered by Kuroki.

Motivation

- In 2008 [3], T. K. Dutta, S. Kar and B. K. Maity studied some properties of regular ternary semigroups, completely regular ternary semigroups, intra-regular ternary semigroups and characterized them by using various ideals of ternary semigroups.
- In 2012 [5], S. Kar and P. Sarkar introduced and characterized the concept of fuzzy quasi-ideal and fuzzy bi-ideal in ternary semigroups and studied their properties in ternary semigroup.

Outline

- ① Motivation
- ② Introduction
- ③ Main Results
- ④ References

Our aim of this talk is threefold:

- 1 To characterize regular ordered ternary semigroups in terms of their fuzzy generalized bi-ideals and fuzzy quasi-ideals.
- 2 To characterize regular ordered ternary semigroups in terms of their fuzzy generalized bi-ideals and fuzzy ideals, fuzzy bi-ideals and fuzzy ideals.
- 3 To characterize regular ordered ternary semigroups in terms of fuzzy generalized bi-ideals, fuzzy left ideals and fuzzy right ideals and fuzzy bi-ideals, fuzzy left ideals and fuzzy right ideals.

Definition ([13])

A non-empty set S is called a **ternary semigroup** if there exists a ternary operation $[\cdot] : S \times S \times S \rightarrow S$ satisfying the following condition:

$$[[abc]de] = [a[bcd]e] = [ab[cde]],$$

for all $a, b, c, d, e \in S$.

For simplicity, we shall write $[abc]$ as abc .

Any semigroup can be reduced to a ternary semigroup.
However, S. Banach showed that a ternary semigroup does not necessarily reduce to a semigroup by this examples.

Example

Let $S = \{-i, 0, i\}$. Then S is a ternary semigroup under the multiplication over complex numbers while S is **not** a semigroup under complex number multiplication.

Definition ([1])

A ternary semigroup S is called an **ordered ternary semigroup** if there is a partial order \leq on S such that $a \leq b$ implies $acd \leq bcd$, $cad \leq cbd$ and $cda \leq cdb$, for all $a, b, c, d \in S$.

Example

Let \mathbb{N} be the set of all natural numbers. We define a ternary operation $[\cdot]$ by usual multiplication, that is, $[abc] = abc$ and define a partial order \leq by usual less than or equal to relation. Then $(\mathbb{N}, [\cdot], \leq)$ is an ordered ternary semigroup.

Definition ([1])

A ternary semigroup S is called an **ordered ternary semigroup** if there is a partial order \leq on S such that $a \leq b$ implies $acd \leq bcd$, $cad \leq cbd$ and $cda \leq cdb$, for all $a, b, c, d \in S$.

Example

Let \mathbb{N} be the set of all natural numbers. We define a ternary operation $[\cdot]$ by usual multiplication, that is, $[abc] = abc$ and define a partial order \leq by usual less than or equal to relation. Then $(\mathbb{N}, [\cdot], \leq)$ is an ordered ternary semigroup.

Definition ([15])

Let X be a non-empty set. A mapping $f : X \rightarrow [0, 1]$ is called a **fuzzy subset** of X .

Let $a \in S$, we define

$$A_a := \{(x, y, z) \in S \times S \times S : a \leq xyz\}$$

and let 1 be a fuzzy subset of S defined as follows

$$1(x) = 1$$

for each $x \in S$.

Definition ([15])

Let X be a non-empty set. A mapping $f : X \rightarrow [0, 1]$ is called a **fuzzy subset** of X .

Let $a \in S$, we define

$$A_a := \{(x, y, z) \in S \times S \times S : a \leq xyz\}$$

and let 1 be a fuzzy subset of S defined as follows

$$1(x) = 1$$

for each $x \in S$.

$$\begin{aligned}
 & (\forall a \in S)(f \circ g \circ h)(a) \\
 &= \begin{cases} \sup_{(u,v,w) \in A_a} \{\min\{f(u), g(v), h(w)\}\} & \text{if } A_a \neq \emptyset, \\ 0 & \text{if } A_a = \emptyset \end{cases}
 \end{aligned}$$

$f \subseteq g$ if and only if $f(x) \leq g(x)$.

$$(f \cap g)(x) = \min\{f(x), g(x)\}$$

and

$$(f \cup g)(x) = \max\{f(x), g(x)\}.$$

Definition ([14])

Let S be an ordered ternary semigroup. A fuzzy subset f of S is called a **fuzzy ternary subsemigroup of S** if for any $x, y, z \in S$,

- ① $f(xyz) \geq \min\{f(x), f(y), f(z)\}$ and
- ② $x \leq y$ implies $f(x) \geq f(y)$.

Definition ([2])

Let S be an ordered ternary groupoid. A fuzzy subset f of S is called a **fuzzy right [resp. left, lateral] ideal of S** if for any $x, y, z \in S$,

- ① $f(xyz) \geq f(x)$ [resp. $f(xyz) \geq f(z)$, $f(xyz) \geq f(y)$] and
- ② $x \leq y$ implies $f(x) \geq f(y)$.

We call f a **fuzzy ideal** of an ordered ternary semigroup S if f is a fuzzy right, left and lateral ideal of S .

Definition

Let S be an ordered ternary groupoid. A fuzzy subset f of S is called a **fuzzy quasi-ideal of S** if

- ① $(f \circ 1 \circ 1) \cap (1 \circ f \circ 1) \cap (1 \circ 1 \circ f) \subseteq f$,
- ② $(f \circ 1 \circ 1) \cap (1 \circ 1 \circ f \circ 1 \circ 1) \cap (1 \circ 1 \circ f) \subseteq f$ and
- ③ $x \leq y$ implies $f(x) \geq f(y)$ for every $x, y \in S$.

Definition ([14])

Let S be an ordered ternary semigroup. A fuzzy subset f of S is called a **fuzzy generalized bi-ideal of S** if for any $u, v, w, x, y \in S$,

- ① $f(uvwx y) \geq \min\{f(u), f(w), f(y)\}$ and
- ② $x \leq y$ implies $f(x) \geq f(y)$.

Definition ([14])

Let S be an ordered ternary semigroup. A fuzzy subset f of S is called a **fuzzy bi-ideal of S** if for any $u, v, w, x, y \in S$,

- ① f is a fuzzy ternary subsemigroup of S ,
- ② $f(uvwxy) \geq \min\{f(u), f(w), f(y)\}$ and
- ③ $x \leq y$ implies $f(x) \geq f(y)$.

An ordered ternary semigroup S is called **regular** [3] if for each $a \in S$ there exists $x \in S$ such that $a \leq axa$.

Outline

- ① Motivation
- ② Introduction
- ③ Main Results
- ④ References

Main Result 1

For an ordered ternary semigroup S , the following assertions are equivalent:

- (1) S is regular.
- (2) $f = f \circ 1 \circ f \circ 1 \circ f$ for every fuzzy generalized bi-ideal f of S .
- (3) $f = f \circ 1 \circ f \circ 1 \circ f$ for every fuzzy bi-ideal f of S .
- (4) $f = f \circ 1 \circ f \circ 1 \circ f$ for every fuzzy quasi-ideal f of S .

Main Result 2

Let S be an ordered ternary semigroup. Then the following statements are equivalent:

- (1) S is regular.
- (2) $f \circ g \circ f \circ g \circ f = f \cap g$ for any fuzzy generalized bi-ideal f and any fuzzy ideal g of S .
- (3) $f \circ g \circ f \circ g \circ f = f \cap g$ for any fuzzy bi-ideal f and any fuzzy ideal g of S .

Main Result 3

Let S be an ordered ternary semigroup. Then the following statements are equivalent:

- (1) S is regular.
- (2) $h \cap f \cap g \subseteq h \circ 1 \circ f \circ 1 \circ g$ for every fuzzy generalized bi-ideal f , every fuzzy left ideal g and every fuzzy right ideal h of S .
- (3) $h \cap f \cap g \subseteq h \circ 1 \circ f \circ 1 \circ g$ for every fuzzy bi-ideal f , every fuzzy left ideal g and every fuzzy right ideal h of S .

Outline

- ① Motivation
- ② Introduction
- ③ Main Results
- ④ References

References I

- [1] T. Changphas, *A Note on Quasi and Bi-Ideals in Ordered Ternary Semigroups*, Int. Journal of Math. Analysis, **6** (2012), no. 11, 527–532.
- [2] R. Chinram and S. Saelee, *Fuzzy Ideals and Fuzzy Filters of Ordered Ternary Semigroups*, Journal of Mathematics Research, **2** (2010), no. 1, 93–97.
- [3] T. K. Dutta, S. Kar and B. K. Maity, *On Ideal in Regular Ternary Semigroups*, Discussiones Mathematicae, General Algebra and Applications, **28** (2008), 147–159.
- [4] A. Iampan, *On Ordered Ideal Extensions of Ordered Ternary Semigroups*, Lobachevskii Journal of Mathematics, **31** (2010), no. 1, 13–17.

References II

- [5] S. Kar and P. Sarkar, *Fuzzy quasi-ideals and fuzzy bi-ideals of ternary semigroups*, *Annals of Fuzzy Mathematics and Informations*, **4** (2012), no. 2, 407–423.
- [6] N. Kuroki, *On fuzzy ideals and fuzzy bi-ideals in semigroups*, *Fuzzy Sets and Systems*, **5** (1981), 203–215.
- [7] N. Kuroki, *On fuzzy semigroups*, *Inf. Sci.*, **53** (1991), 203–236.
- [8] N. Kuroki, *Fuzzy semiprime quasi-ideals in semigroups*, *Inf. Sci.*, **75** (1993), 201–211.
- [9] D. M. Lee and S. K. Lee, *On intra-regular ordered semigroups*, *Kangweon-Kyungki Math. Jour*, **14** (2006), no. 1, 95–100.

References III

- [10] D. H. Lehmer, *A ternary analogue of Abelian groups*, Amer. Jr. of Math., **59** (1972), 329–338.
- [11] S. Lekkoksung and N. Lekkoksung, *On Intra-Regular Ordered Ternary Semigroups*, Int. Journal of Math. Analysis, **6** (2012), no. 2, 69–73.
- [12] M. L. Santiago, *Some contributions to the study of ternary semigroups and semiheaps*, (Ph. D. Thesis, 1983, University of Madras).
- [13] F. M. Sioson, *Ideal theory in ternary semigroups*, Math. Japonica., **10** (1965), 63–84.

References IV

- [14] N. Yaqoob, S. Abdullah, N. Rehman and M. Naeem,
Roughness and Fuzziness in Ordered Ternary Semigroups,
World Applied Sciences Journal, **17** (2012), no. 12,
1683–1693.
- [15] L. A. Zadeh, Fuzzy Sets, *Information and Control*, **8**
(1965), 338–353.

Thank you for your attention!