



TECHNISCHE
UNIVERSITÄT
WIEN

Vienna University of Technology

Centraliser clones on finite domains¹

Mike Behrisch

Institute of Computer Languages,
Theory and Logic Group,
Vienna University of Technology

2nd July 2014
Prague, Czech Republic

¹Supported by the Austrian Science Fund (FWF) under grant I836-N23 

Outline

- 1 A Galois theory based on commutation
- 2 Structure of the lattice of centralisers
- 3 Minimal/maximal centralisers

Why centraliser clones?

CSP, again!

$\underline{T} = \langle T; Q \rangle$ finite rel structure, of finite signature, **template**.

CSP (\underline{T}) is the **decision problem** \exists hom. $h: \underline{A} \rightarrow \underline{T}$? for
finite \underline{A} of the same signature.

Why centraliser clones?

CSP, again!

$\underline{T} = \langle T; Q \rangle$ finite rel structure, of finite signature, **template**.

$\text{CSP}(\underline{T})$ is the **decision problem** $\exists \text{ hom. } h: \underline{A} \rightarrow \underline{T}?$ for finite \underline{A} of the same signature.

Complexity

$\text{CSP}(\underline{T})$ is **always in NP**, **some** CSP's are **in P**.

Why centraliser clones?

CSP, again!

$\underline{T} = \langle T; Q \rangle$ finite rel structure, of finite signature, **template**.
 CSP (\underline{T}) is the **decision problem** $\exists \text{ hom. } h: \underline{A} \rightarrow \underline{T}?$ for
 finite \underline{A} of the same signature.

Complexity

CSP (\underline{T}) is **always in NP**, some CSP's are **in P**.

CSP dichotomy conjecture

Feder/Vardi: $|T| = 2 \Rightarrow$ **either** CSP (\underline{T}) **in P** or **NP-complete**.
Conjecture: This **extends to all finite domains**.

Why centraliser clones?

CSP, again!

$\underline{T} = \langle T; Q \rangle$ finite rel structure, of finite signature, **template**.
 CSP (\underline{T}) is the **decision problem** $\exists \text{ hom. } h: \underline{A} \rightarrow \underline{T}?$ for
 finite \underline{A} of the same signature.

Complexity

CSP (\underline{T}) is **always in NP**, **some** CSP's are **in P**.

CSP dichotomy conjecture

Feder/Vardi: $|T| = 2 \Rightarrow$ **either** CSP (\underline{T}) **in P** or **NP-complete**.
Conjecture: This **extends to all finite domains**.

Connection to centraliser clones

Feder/Madelaine/Stewart | Broniek: It **suffices** to establish the
 dichotomy conjecture for \underline{T} , where $\text{Pol } \underline{T}$ is a **centraliser clone**.

Outline

- 1 A Galois theory based on commutation
- 2 Structure of the lattice of centralisers
- 3 Minimal/maximal centralisers

Operations and their graphs

Finitary operations

- For $k \in \mathbb{N}_+$ a func $f: A^k \longrightarrow A$ is a k -ary operation on A
- $O_A^{(k)} := A^{A^k}$ set of k -ary operations on A
- $O_A := \bigcup_{k \in \mathbb{N}_+} O_A^{(k)}$ set of all finitary operations on A

Operations and their graphs

Finitary operations

- For $k \in \mathbb{N}_+$ a func $f: A^k \rightarrow A$ is a **k -ary operation on A**
- $O_A^{(k)} := A^{A^k}$ **set of k -ary operations on A**
- $O_A := \bigcup_{k \in \mathbb{N}_+} O_A^{(k)}$ **set of all finitary operations on A**

Graphs

For $f \in O_A^{(n)}$, $F \subseteq O_A$

$$\begin{aligned} f^\bullet &:= \{ (x_1, \dots, x_n, x_{n+1}) \in A^{n+1} \mid x_{n+1} = f(x_1, \dots, x_n) \} \\ &= \{ (x_1, \dots, x_n, f(x_1, \dots, x_n)) \mid (x_1, \dots, x_n) \in A^n \} \end{aligned}$$

$$F^\bullet := \{ f^\bullet \mid f \in F \}$$

Basis of the Galois connection

Commutation

For $n, m \in \mathbb{N}_+$, $f \in O_A^{(n)}$, $g \in O_A^{(m)}$,

$$f \text{ commutes with } g :\iff f \perp g :\iff f \triangleright g^\bullet$$

Basis of the Galois connection

Commutation

For $n, m \in \mathbb{N}_+$, $f \in O_A^{(n)}$, $g \in O_A^{(m)}$,

$$f \text{ commutes with } g : \Longleftrightarrow f \perp g : \Longleftrightarrow f \triangleright g^\bullet$$

Lemma

For $n, m \in \mathbb{N}_+$, $f \in O_A^{(n)}$, $g \in O_A^{(m)}$, we have

$$f \perp g \Longleftrightarrow f : \langle A; g \rangle^n \longrightarrow \langle A; g \rangle \text{ is a homomorphism.}$$

A more concrete characterisation of commutation

Lemma

For $n, m \in \mathbb{N}_+$, $f \in O_A^{(n)}$, $g \in O_A^{(m)}$, we have

$$f \perp g \iff$$

A more concrete characterisation of commutation

Lemma

For $n, m \in \mathbb{N}_+$, $f \in O_A^{(n)}$, $g \in O_A^{(m)}$, we have

$$f \perp g \iff \forall X = (x_{i,j})_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n}} \in A^{m \times n}:$$

$$\begin{array}{ccc} x_{1,1} & \cdots & x_{1,n} \\ \vdots & \ddots & \vdots \\ x_{m,1} & \cdots & x_{m,n} \end{array}$$

A more concrete characterisation of commutation

Lemma

For $n, m \in \mathbb{N}_+$, $f \in O_A^{(n)}$, $g \in O_A^{(m)}$, we have

$$f \perp g \iff \forall X = (x_{i,j})_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n}} \in A^{m \times n}:$$

$$\begin{array}{ccc} g & & g \\ \left(\begin{array}{ccc} x_{1,1} & \cdots & x_{1,n} \\ \vdots & \ddots & \vdots \\ x_{m,1} & \cdots & x_{m,n} \end{array} \right) & & \\ = & & = \\ g(x(\cdot, 1)) \cdots g(x(\cdot, n)) & & \end{array}$$

A more concrete characterisation of commutation

Lemma

For $n, m \in \mathbb{N}_+$, $f \in O_A^{(n)}$, $g \in O_A^{(m)}$, we have

$$f \perp g \iff \forall X = (x_{i,j})_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n}} \in A^{m \times n}:$$

$$\begin{array}{ccc} g & & g \\ \left(\begin{array}{ccc} x_{1,1} & \cdots & x_{1,n} \\ \vdots & \ddots & \vdots \\ x_{m,1} & \cdots & x_{m,n} \end{array} \right) & & \\ = & & = \\ g(x(\cdot, 1)) \cdots g(x(\cdot, n)) & & \\ \in g^\bullet & & \in g^\bullet \end{array}$$

A more concrete characterisation of commutation

Lemma

For $n, m \in \mathbb{N}_+$, $f \in O_A^{(n)}$, $g \in O_A^{(m)}$, we have

$$f \perp g \iff \forall X = (x_{i,j})_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n}} \in A^{m \times n}:$$

$$\begin{array}{ccc} \begin{array}{c} g \\ \left(\right. \\ x_{1,1} \quad \cdots \quad x_{1,n} \\ \vdots \quad \ddots \quad \vdots \\ x_{m,1} \quad \cdots \quad x_{m,n} \\ \left. \right) \\ = \\ f(g(x(\cdot, 1)) \cdots g(x(\cdot, n))) \\ \in g^\bullet \end{array} & = & \begin{array}{c} f(x(1, \cdot)) \\ \vdots \\ f(x(m, \cdot)) \\ \\ y \end{array} \end{array}$$

A more concrete characterisation of commutation

Lemma

For $n, m \in \mathbb{N}_+$, $f \in O_A^{(n)}$, $g \in O_A^{(m)}$, we have

$$f \perp g \iff \forall X = (x_{i,j})_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n}} \in A^{m \times n}:$$

$$\begin{array}{ccc} \begin{array}{c} g \\ \left(\begin{array}{ccc} x_{1,1} & \cdots & x_{1,n} \\ \vdots & \ddots & \vdots \\ x_{m,1} & \cdots & x_{m,n} \end{array} \right) \end{array} & = & f(x(1, \cdot)) \\ & & \vdots \\ \begin{array}{c} f \\ \left(\begin{array}{ccc} x_{1,1} & \cdots & x_{1,n} \\ \vdots & \ddots & \vdots \\ x_{m,1} & \cdots & x_{m,n} \end{array} \right) \end{array} & = & f(x(m, \cdot)) \\ & & \vdots \\ & = & y \\ & \in g^\bullet & \in g^\bullet \end{array}$$

A more concrete characterisation of commutation

Lemma

For $n, m \in \mathbb{N}_+$, $f \in O_A^{(n)}$, $g \in O_A^{(m)}$, we have

$$f \perp g \iff \forall X = (x_{i,j})_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n}} \in A^{m \times n}:$$

$$\begin{array}{ccc}
 \begin{array}{c} g \\ \left(\right. \\ f(\quad x_{1,1} \quad \cdots \quad x_{1,n} \quad) = \\ \quad \vdots \quad \ddots \quad \vdots \\ f(\quad x_{m,1} \quad \cdots \quad x_{m,n} \quad) = \\ \left. \right) \\ = \end{array} &
 \begin{array}{c} g \\ \left(\right. \\ \quad \quad \quad x_{1,n} \quad) = \\ \quad \quad \quad \vdots \\ \quad \quad \quad x_{m,n} \quad) = \\ \left. \right) \\ = \end{array} &
 \begin{array}{c} g \\ \left(\right. \\ f(x(1, \cdot)) \\ \quad \vdots \\ f(x(m, \cdot)) \\ \left. \right) \\ = \end{array} \\
 f(g(x(\cdot, 1)) \cdots g(x(\cdot, n))) & = & g(f(x(1, \cdot)), \dots, f(x(m, \cdot))) \\
 \in g^\bullet & &
 \end{array}$$

Commutativity is symmetric

Corollary

For $f, g \in O_A$ we have $f \perp g \iff g \perp f$.

Galois correspondence induced by commutation

Definition

For $F \subseteq O_A$ we put

$$F^* := \{g \in O_A \mid \forall f \in F: g \perp f\} \quad (\text{centraliser of } F)$$

$$F^{**} \quad (\text{bicentraliser of } F, \text{ bicentral closure of } F)$$

$$\mathcal{C}_A := \{F^* \mid F \subseteq O_A\} \quad (\text{lattice of centraliser clones})$$

Galois correspondence induced by commutation

Definition

For $F \subseteq O_A$ we put

$$F^* := \{g \in O_A \mid \forall f \in F: g \perp f\} \quad (\text{centraliser of } F)$$

$$F^{**} \quad (\text{bicentraliser of } F, \text{ bicentral closure of } F)$$

$$\mathcal{C}_A := \{F^* \mid F \subseteq O_A\} \quad (\text{lattice of centraliser clones})$$

Alternatively

For $F \subseteq O_A$ we have

$$F^* = \text{Pol}_A F^\bullet \quad (\text{centralisers are clones!})$$

$$= \{f \in O_A \mid f^\bullet \in \text{Inv}_A F\}$$

Comparison with clone closure

Observation

Centralisers

= usual Pol – Inv where **relations** are **confined to graphs**

Comparison with clone closure

Observation

Centralisers

= usual Pol – Inv where **relations** are **confined to graphs**

Lemma

For $F \subseteq O_A$ we have $F^* = \langle F \rangle_{O_A}^*$.

Comparison with clone closure

Observation

Centralisers

= usual Pol – Inv where **relations** are **confined to graphs**

Lemma

For $F \subseteq O_A$ we have $F^* = \langle F \rangle_{O_A}^*$.

Proof.

$$g \in F^* \iff g^\bullet \in \text{Inv}_A F = \text{Inv}_A \langle F \rangle_{O_A} \iff g \in \langle F \rangle_{O_A}^*. \quad \square$$

Comparison with clone closure

Observation

Centralisers

= usual Pol – Inv where **relations** are **confined to graphs**

Lemma

For $F \subseteq O_A$ we have $F^* = \langle F \rangle_{O_A}^*$.

Corollary

For $F \subseteq O_A$ it is $F^{**} = \langle F \rangle_{O_A}^{**}$, i.e. $\langle F \rangle_{O_A} \subseteq F^{**}$.

Comparison with clone closure

Observation

Centralisers

= usual Pol – Inv where **relations** are **confined to graphs**

Lemma

For $F \subseteq O_A$ we have $F^* = \langle F \rangle_{O_A}^*$.

Corollary

For $F \subseteq O_A$ it is $F^{**} = \langle F \rangle_{O_A}^{**}$, i.e. $\langle F \rangle_{O_A} \subseteq F^{**}$.

Proof.

$$F^* = \langle F \rangle_{O_A}^* \implies F^{**} = \langle F \rangle_{O_A}^{**} \supseteq \langle F \rangle_{O_A}.$$



Outline

- 1 A Galois theory based on commutation
- 2 Structure of the lattice of centralisers
- 3 Minimal/maximal centralisers

Finitely many centralisers

Theorem (R. Pöschel, unpublished)

$|A| = k$ *finite* $\implies \forall F \subseteq O_A \exists Q_F \subseteq R_A^{(4)}$ *4-ary rel's*:

$$F^* = \left(\langle F \rangle_{O_A}^{(k)} \right)^* \cap \text{Pol}_A Q_F.$$

Finitely many centralisers

Theorem (R. Pöschel, unpublished)

$|A| = k$ *finite* $\implies \forall F \subseteq O_A \exists Q_F \subseteq R_A^{(4)}$ *4-ary rel's*:

$$F^* = \left(\langle F \rangle_{O_A}^{(k)} \right)^* \cap \text{Pol}_A Q_F.$$

Corollary (S. Burris/R. D. Willard, 1987, different proof)

$|A| = k$ *finite* $\implies |\mathcal{C}_A|$ *finite*.

Finitely many centralisers

Theorem (R. Pöschel, unpublished)

$|A| = k$ *finite* $\implies \forall F \subseteq O_A \exists Q_F \subseteq R_A^{(4)}$ *4-ary rel's*:

$$F^* = \left(\langle F \rangle_{O_A}^{(k)} \right)^* \cap \text{Pol}_A Q_F.$$

Corollary (S. Burris/R. D. Willard, 1987, different proof)

$|A| = k$ *finite* $\implies |\mathcal{C}_A|$ *finite*.

decomp: $\mathcal{C}_A \longrightarrow \mathfrak{P}(O_A^{(k)}) \times \mathfrak{P}(\mathfrak{P}(A^4))$ is *inj.*, as
 $\Sigma \longmapsto (\Sigma^{*(k)}, Q_{\Sigma^*})$
 $(\Sigma^{*(k)})^* \cap \text{Pol}_A Q_{\Sigma^*} = (\langle \Sigma^* \rangle_{O_A}^{(k)})^* \cap \text{Pol}_A Q_{\Sigma^*} = \Sigma^{**} = \Sigma.$

Special cases

$|A| = 2$, A. B. Кузнецов, 1977

$|C_A| = 25$,

Special cases

$|A| = 2$, A. В. Кузнецов, 1977

$|C_A| = 25$,



Special cases

$|A| = 2$, A. B. Кузнецов, 1977

$|C_A| = 25$,

⊖ Кузнецов's proof unavailable,

Special cases

$|A| = 2$, A. B. Кузнецов, 1977

$|C_A| = 25$,

⊖ Кузнецов's proof unavailable,

⊕ sketched by László Szabó (1978), reproved by Miki Hermann, 2008

Special cases

$|A| = 2$, A. В. Кузнецов, 1977

$|C_A| = 25$,

⊖ Кузнецов's proof unavailable,

⊕ sketched by László Szabó (1978), reproved by Miki Hermann, 2008

$|A| = 3$, А. Ф. Данильченко, 1974–79

$|C_A| = 2986$,

Special cases

$|A| = 2$, А. В. Кузнецов, 1977

$|C_A| = 25$,

⊖ Кузнецов's proof unavailable,

⊕ sketched by László Szabó (1978), reproved by Miki Hermann, 2008

$|A| = 3$, А. Ф. Данильченко, 1974–79

$|C_A| = 2986$,

⊕ available, readable

Special cases

$|A| = 2$, А. В. Кузнецов, 1977

$|C_A| = 25$,

⊖ Кузнецов's proof unavailable,

⊕ sketched by László Szabó (1978), reproved by Miki Hermann, 2008

$|A| = 3$, А. Ф. Данильченко, 1974–79

$|C_A| = 2986$,

⊕ available, readable

⊖ does not give intuition how to find these clones in general

Special cases

$|A| = 2$, А. В. Кузнецов, 1977

$|C_A| = 25$,

⊖ Кузнецов's proof unavailable,

⊕ sketched by László Szabó (1978), reproved by Miki Hermann, 2008

$|A| = 3$, А. Ф. Данильченко, 1974–79

$|C_A| = 2986$,

⊕ available, readable

⊖ does not give intuition how to find these clones in general

$|A| = 4$

Special cases

$|A| = 2$, А. В. Кузнецов, 1977

$|C_A| = 25$,

⊖ Кузнецов's proof unavailable,

⊕ sketched by László Szabó (1978), reproved by Miki Hermann, 2008

$|A| = 3$, А. Ф. Данильченко, 1974–79

$|C_A| = 2986$,

⊕ available, readable

⊖ does not give intuition how to find these clones in general

$|A| = 4$

We don't know, but we would like to.

More structural properties/questions

C_A is finite \implies

More structural properties/questions

\mathcal{C}_A is finite \implies

- finitely many atoms/co-atoms,

More structural properties/questions

\mathcal{C}_A is finite \implies

- finitely many atoms/co-atoms,
- $\forall F \in \mathcal{C}_A \setminus \{\perp\} \exists A \in \mathcal{C}_A \text{ atom: } A \subseteq F.$
- $\forall F \in \mathcal{C}_A \setminus \{\top\} \exists C \in \mathcal{C}_A \text{ co-atom: } F \subseteq C.$

More structural properties/questions

\mathcal{C}_A is finite \implies

- finitely many atoms/co-atoms,
- $\forall F \in \mathcal{C}_A \setminus \{\perp\} \exists A \in \mathcal{C}_A \text{ atom: } A \subseteq F.$
- $\forall F \in \mathcal{C}_A \setminus \{\top\} \exists C \in \mathcal{C}_A \text{ co-atom: } F \subseteq C.$
- char of \perp, \top ? (easy)

More structural properties/questions

\mathcal{C}_A is finite \implies

- finitely many atoms/co-atoms,
- $\forall F \in \mathcal{C}_A \setminus \{\perp\} \exists A \in \mathcal{C}_A \text{ atom: } A \subseteq F.$
- $\forall F \in \mathcal{C}_A \setminus \{\top\} \exists C \in \mathcal{C}_A \text{ co-atom: } F \subseteq C.$
- char of \perp, \top ? (easy)
- char of \bigcap -irreducibles? (not easy)

More structural properties/questions

\mathcal{C}_A is finite \implies

- finitely many atoms/co-atoms,
- $\forall F \in \mathcal{C}_A \setminus \{\perp\} \exists A \in \mathcal{C}_A \text{ atom: } A \subseteq F.$
- $\forall F \in \mathcal{C}_A \setminus \{\top\} \exists C \in \mathcal{C}_A \text{ co-atom: } F \subseteq C.$
- char of \perp, \top ? (easy)
- char of \bigcap -irreducibles? (not easy)
every $F \in \mathcal{C}_A$ is \bigcap of \bigcap -irreducibles

More structural properties/questions

\mathcal{C}_A is finite \implies

- finitely many atoms/co-atoms,
- $\forall F \in \mathcal{C}_A \setminus \{\perp\} \exists A \in \mathcal{C}_A \text{ atom: } A \subseteq F.$
- $\forall F \in \mathcal{C}_A \setminus \{\top\} \exists C \in \mathcal{C}_A \text{ co-atom: } F \subseteq C.$
- char of \perp, \top ? (easy)
- char of \bigcap -irreducibles? (not easy)
every $F \in \mathcal{C}_A$ is \bigcap of \bigcap -irreducibles

Данильченко's method

Describe a **generator** for each \bigcap -irreducible $F \in \mathcal{C}_A$

\top and \perp of \mathcal{C}_A

Lemma

We have $O_A^ = J_A$.*

\top and \perp of \mathcal{C}_A

Lemma

We have $O_A^* = J_A$.

Proof.

W.l.o.g. $|A| \geq 2$.

\top and \perp of \mathcal{C}_A

Lemma

We have $O_A^* = J_A$.

Proof.

W.l.o.g. $|A| \geq 2$.

$$f \in O_A^* \implies f^\bullet \in \text{Inv}_A O_A = \text{Diag}_A \implies \exists \epsilon: f^\bullet = d_\epsilon$$

\top and \perp of \mathcal{C}_A

Lemma

We have $O_A^* = J_A$.

Proof.

W.l.o.g. $|A| \geq 2$.

$$f \in O_A^* \implies f^\bullet \in \text{Inv}_A O_A = \text{Diag}_A \implies \exists \varepsilon: f^\bullet = d_\varepsilon$$

$$f \notin J_A \implies \varepsilon = \Delta, \text{ i.e. } f^\bullet = \nabla$$

\top and \perp of \mathcal{C}_A

Lemma

We have $O_A^* = J_A$.

Proof.

W.l.o.g. $|A| \geq 2$.

$$f \in O_A^* \implies f^\bullet \in \text{Inv}_A O_A = \text{Diag}_A \implies \exists \varepsilon: f^\bullet = d_\varepsilon$$

$$f \notin J_A \implies \varepsilon = \Delta, \text{ i.e. } f^\bullet = \nabla \text{ (no graph).}$$



\top and \perp of \mathcal{C}_A

Lemma

We have $O_A^* = J_A$.

Proof.

W.l.o.g. $|A| \geq 2$.

$$f \in O_A^* \implies f^\bullet \in \text{Inv}_A O_A = \text{Diag}_A \implies \exists \varepsilon: f^\bullet = d_\varepsilon$$

$$f \notin J_A \implies \varepsilon = \Delta, \text{ i.e. } f^\bullet = \nabla \text{ (no graph).}$$



Corollary

For $F \subseteq O_A$ it is $F^* = O_A \iff F \subseteq J_A$.

Roadmap

- Co-atoms are special \bigcap -irreducibles
- Co-atoms and **atoms** are related
- Atoms are related to **minimal clones**

Outline

- 1 A Galois theory based on commutation
- 2 Structure of the lattice of centralisers
- 3 Minimal/maximal centralisers

Parametrically pre-complete sets

Lemma

For $F \subseteq O_A$ tfae:

$$F^{**} \subsetneq O_A \quad \forall G \supsetneq F: \quad G^{**} = O_A \quad (1)$$

$$J_A \subsetneq F^* \quad \forall G \supsetneq F: \quad G^* = J_A \quad (2)$$

$$J_A \subsetneq F^* \quad \forall g \in F^* \setminus J_A: \quad F = g^* \quad (3)$$

Parametrically pre-complete sets

Lemma

For $F \subseteq O_A$ tfae:

$$F^{**} \subsetneq O_A \quad \forall G \supsetneq F: \quad G^{**} = O_A \quad (1)$$

$$J_A \subsetneq F^* \quad \forall G \supsetneq F: \quad G^* = J_A \quad (2)$$

$$J_A \subsetneq F^* \quad \forall g \in F^* \setminus J_A: \quad F = g^* \quad (3)$$

Corollary

$F \subseteq O_A$ *param. pre-complete* $\implies F = f^*$ for some $f \in F^* \setminus J_A$.

Minimal functions

Proposition

$F \subseteq O_A$ *parametrically pre-complete*

$\implies \exists M$ *minimal clone*: $F = M^*$

$\implies \exists f \in O_A^{(\leq \max(3, |A|))}$ *minimal function*: $F = f^*$.

Minimal functions

Proposition

$F \subseteq O_A$ *parametrically pre-complete*

$\implies \exists M$ *minimal clone*: $F = M^*$

$\implies \exists f \in O_A^{(\leq \max(3, |A|))}$ *minimal function*: $F = f^*$.

Proof.

- $F^* \not\supseteq J_A \implies F^* \supseteq M \not\supseteq J_A$
for some minimal clone $M = \langle f \rangle_{O_A}$ generated by one of Rosenberg's minimal functions.

Minimal functions

Proposition

$F \subseteq O_A$ *parametrically pre-complete*

$\implies \exists M$ *minimal clone*: $F = M^*$

$\implies \exists f \in O_A^{(\leq \max(3, |A|))}$ *minimal function*: $F = f^*$.

Proof.

- $F^* \not\supseteq J_A \implies F^* \supseteq M \not\supseteq J_A$
for some minimal clone $M = \langle f \rangle_{O_A}$ generated by one of Rosenberg's minimal functions.
- Equivalently: $F \subseteq M^* = \langle f \rangle_{O_A}^* = f^* \subsetneq O_A$.

Minimal functions

Proposition

$F \subseteq O_A$ *parametrically pre-complete*

$\implies \exists M$ *minimal clone*: $F = M^*$

$\implies \exists f \in O_A^{(\leq \max(3, |A|))}$ *minimal function*: $F = f^*$.

Proof.

- $F^* \not\supseteq J_A \implies F^* \supseteq M \not\supseteq J_A$
for some minimal clone $M = \langle f \rangle_{O_A}$ generated by one of Rosenberg's minimal functions.
- Equivalently: $F \subseteq M^* = \langle f \rangle_{O_A}^* = f^* \subsetneq O_A$.
- Since $M^* \supseteq F$ and $(M^*)^{**} = M^* \neq O_A$,
 $\implies F = M^* = f^*$. □

Unary case

Unary minimal functions from Rosenberg's theorem

$f \neq \text{id}_A$ and ($f \circ f = f$ or $f^p = \text{id}_A$ for some prime p)

Unary case

Unary minimal functions from Rosenberg's theorem

$f \neq \text{id}_A$ and ($f \circ f = f$ or $f^p = \text{id}_A$ for some prime p)

Recall

All of these unary functions generate **indeed** minimal clones.

Unary case

Unary minimal functions from Rosenberg's theorem

$f \neq \text{id}_A$ and ($f \circ f = f$ or $f^p = \text{id}_A$ for some prime p)

Recall

All of these unary functions generate **indeed** minimal clones.

Question

Which of these yield **maximal centralisers**?

Almost all

Theorem (W. Harnau, 1974)

f^* is a co-atom in $\mathcal{V}_A := \left\{ g^* \mid g \in O_A^{(1)} \right\}$ if and only if

- ① $f \neq \text{id}_A$ and $f \circ f = f$, or
- ② $f^p = \text{id}_A$ for some prime p and
 f does *not* have *exactly one fixed point*

Almost all

Theorem (W. Harnau, 1974)

f^* is a co-atom in $\mathcal{V}_A := \left\{ g^* \mid g \in O_A^{(1)} \right\}$ if and only if

- ① $f \neq \text{id}_A$ and $f \circ f = f$, or
- ② $f^p = \text{id}_A$ for some prime p and
 f does *not* have *exactly one fixed point*

Proposition

$$f \in O_A^{(1)} \implies f^{**} \subseteq \left\langle O_A^{(1)} \right\rangle_{O_A}$$

Almost all

Theorem (W. Harnau, 1974)

f^* is a co-atom in $\mathcal{V}_A := \left\{ g^* \mid g \in O_A^{(1)} \right\}$ if and only if

- ① $f \neq \text{id}_A$ and $f \circ f = f$, or
- ② $f^p = \text{id}_A$ for some prime p and f does *not* have *exactly one fixed point*

Proposition

$$f \in O_A^{(1)} \implies f^{**} \subseteq \left\langle O_A^{(1)} \right\rangle_{O_A}$$

Corollary

*All functions from Harnau's theorem yield **maximal centralisers** (**minimal bicentralisers**).*

Future work

- consider other classes of minimal functions
- exploit characterisation of maximal clones
- better algorithms for checking commutation condition
- automatic computation of bicentral closure up to a certain arity
- better upper bounds for the minimal arity needed to
** -generate centraliser clones