

Weighted Clones and Valued CSPs

Stanislav Živný

Algebras & Clones fest
Prague, Czech republic, 07/2014



EPSRC

Engineering and Physical Sciences
Research Council

*The London
Mathematical
Society*



THE
ROYAL
SOCIETY

Motivation

CSPs \Leftrightarrow Clones

[*Jeavons et al. JACM'97, Jeavons TCS'98, Bulatov et al. SICOMP'05*]


Motivation

CSPs \Leftrightarrow Clones


[Jeavons et al. JACM'97, Jeavons TCS'98, Bulatov et al. SICOMP'05]

3-Colouring

G




K_3




k -Clique

K_k



G



Constraint Satisfaction Problems

$$\text{CSP}(\mathcal{A}, \mathcal{B})$$



Given two classes \mathcal{A} and \mathcal{B} of relational structures
and $\mathbf{A} \in \mathcal{A}$, $\mathbf{B} \in \mathcal{B}$, decide whether $\mathbf{A} \rightarrow \mathbf{B}$.

Non-Uniform $\text{CSP}(\mathcal{A}, \mathcal{B})$: $\mathcal{A} = \text{all}$, $\mathcal{B} = \{\Gamma\}$

$$\text{CSP}(\Gamma) = \text{CSP}(-, \{\Gamma\}), \text{ finite relational structure } \Gamma$$


Non-Uniform $\text{CSP}(\mathcal{A}, \mathcal{B})$: $\mathcal{A} = \text{all}$, $\mathcal{B} = \{\Gamma\}$

$$\text{CSP}(\Gamma) = \text{CSP}(-, \{\Gamma\}), \text{ finite relational structure } \Gamma$$



Non-Uniform CSP(\mathcal{A}, \mathcal{B}): $\mathcal{A} = \text{all}$, $\mathcal{B} = \{\Gamma\}$

$\text{CSP}(\Gamma) = \text{CSP}(-, \{\Gamma\})$, finite relational structure Γ



CSPs \Leftrightarrow Clones

[Jeavons et al. JACM'97, Jeavons TCS'98, Bulatov et al. SICOMP'05]

Motivation

CSPs \Leftrightarrow Clones

[Jeavons et al. JACM'97, Jeavons TCS'98, Bulatov et al. SICOMP'05]

Complexity of CSP(Γ)

1. $\text{CSP}(\Gamma') \equiv_p \text{CSP}(\Gamma)$
where $\Gamma' = \text{RelClone}(\Gamma)$ is the closure of Γ under $\exists, \wedge, =$
2. relational clones $\text{RelClone}(\Gamma)$
 \approx clones $\text{Pol}(\Gamma)$
3. properties of $\text{Pol}(\Gamma)$

Complexity of $\text{CSP}(\Gamma)$

1. $\text{CSP}(\Gamma') \equiv_p \text{CSP}(\Gamma)$
where $\Gamma' = \text{RelClone}(\Gamma)$ is the closure of Γ under $\exists, \wedge, =$
2. relational clones $\text{RelClone}(\Gamma)$
 \approx clones $\text{Pol}(\Gamma)$
3. properties of $\text{Pol}(\Gamma)$

Example: $\text{Pol}(\Gamma)$ contains a Mal'cev operation $\Rightarrow \text{CSP}(\Gamma)$ tractable

[Bulatov & Dalmau SICOMP'06]

CSPs \Leftrightarrow Clones

[Jeavons et al. JACM'97, Jeavons TCS'98, Bulatov et al. SICOMP'05]

Valued CSPs \Leftrightarrow Weighted Clones

[Cohen, Cooper, Creed, Jeavons, Ž. SICOMP'13]

Valued CSPs \Leftrightarrow Weighted Clones

[Cohen, Cooper, Creed, Jeavons, Ž. SICOMP'13]

Valued CSPs

- ▶ fixed finite set D
- ▶ $\overline{\mathbb{Q}} = \mathbb{Q} \cup \{\infty\}$

Valued CSPs

- ▶ fixed finite set D
- ▶ $\overline{\mathbb{Q}} = \mathbb{Q} \cup \{\infty\}$

Let Γ be a finite set of functions $\phi : D^m \rightarrow \overline{\mathbb{Q}}$. An instance of VCSP(Γ) is an optimisation problem of the form

$$\min_{x_1, \dots, x_n \in D} \phi_1(x_{1,1}, \dots, x_{1,m_1}) + \dots + \phi_q(x_{q,1}, \dots, x_{q,m_q})$$

where $x_{i,j} \in \{x_1, \dots, x_n\}$ and all functions $\phi_i \in \Gamma$.

Valued CSPs

- ▶ fixed finite set D
- ▶ $\overline{\mathbb{Q}} = \mathbb{Q} \cup \{\infty\}$

weighted relations

Let Γ be a finite set of functions $\phi : D^m \rightarrow \overline{\mathbb{Q}}$. An instance of $\text{VCSP}(\Gamma)$ is an optimisation problem of the form

$$\min_{x_1, \dots, x_n \in D} \phi_1(x_{1,1}, \dots, x_{1,m_1}) + \dots + \phi_q(x_{q,1}, \dots, x_{q,m_q})$$

where $x_{i,j} \in \{x_1, \dots, x_n\}$ and all functions $\phi_i \in \Gamma$.

Valued CSPs

- ▶ fixed finite set D
- ▶ $\overline{\mathbb{Q}} = \mathbb{Q} \cup \{\infty\}$

Let Γ be a finite set of functions $\phi : D^m \rightarrow \overline{\mathbb{Q}}$. An instance of VCSP(Γ) is an optimisation problem of the form

$$\min_{x_1, \dots, x_n \in D} \phi_1(x_{1,1}, \dots, x_{1,m_1}) + \dots + \phi_q(x_{q,1}, \dots, x_{q,m_q})$$

where $x_{i,j} \in \{x_1, \dots, x_n\}$ and all functions $\phi_i \in \Gamma$.

Valued CSPs

- ▶ fixed finite set D
- ▶ $\overline{\mathbb{Q}} = \mathbb{Q} \cup \{\infty\}$

possibly different m for different $\phi \in \Gamma$

Let Γ be a finite set of functions $\phi : D^m \rightarrow \overline{\mathbb{Q}}$. An instance of $\text{VCSP}(\Gamma)$ is an optimisation problem of the form

$$\min_{x_1, \dots, x_n \in D} \phi_1(x_{1,1}, \dots, x_{1,m_1}) + \dots + \phi_q(x_{q,1}, \dots, x_{q,m_q})$$

where $x_{i,j} \in \{x_1, \dots, x_n\}$ and all functions $\phi_i \in \Gamma$.

Valued CSPs

- ▶ fixed finite set D
- ▶ $\overline{\mathbb{Q}} = \mathbb{Q} \cup \{\infty\}$

Let Γ be a finite set of functions $\phi : D^m \rightarrow \overline{\mathbb{Q}}$. An instance of VCSP(Γ) is an optimisation problem of the form

$$\min_{x_1, \dots, x_n \in D} \phi_1(x_{1,1}, \dots, x_{1,m_1}) + \dots + \phi_q(x_{q,1}, \dots, x_{q,m_q})$$

where $x_{i,j} \in \{x_1, \dots, x_n\}$ and all functions $\phi_i \in \Gamma$.

Valued CSPs

- ▶ fixed finite set D
- ▶ $\overline{\mathbb{Q}} = \mathbb{Q} \cup \{\infty\}$

Let Γ be a finite set of functions $\phi : D^m \rightarrow \overline{\mathbb{Q}}$. An instance of $\text{VCSP}(\Gamma)$ is an optimisation problem of the form

$$\min_{x_1, \dots, x_n \in D} \phi_1(x_{1,1}, \dots, x_{1,m_1}) + \dots + \phi_q(x_{q,1}, \dots, x_{q,m_q})$$

where $x_{i,j} \in \{x_1, \dots, x_n\}$ and all functions $\phi_i \in \Gamma$.

Complexity of $\text{VCSP}(\Gamma)$ for all possible Γ !

Valued CSPs

Let Γ be a finite set of functions $\phi : D^m \rightarrow \overline{\mathbb{Q}}$. An instance of $\text{VCSP}(\Gamma)$ is an optimisation problem of the form

$$\min_{x_1, \dots, x_n \in D} \phi_1(x_{1,1}, \dots, x_{1,m_1}) + \dots + \phi_q(x_{q,1}, \dots, x_{q,m_q})$$

where $x_{i,j} \in \{x_1, \dots, x_n\}$ and all functions $\phi_i \in \Gamma$.

Monotone NAE 3-SAT

$$\min_{x_1, \dots, x_n \in \{T, F\}} \sum_{(x_i, x_j, x_k) \in C} \phi_{\text{nae}}(x_i, x_j, x_k)$$

x	y	z	$\phi_{\text{nae}}(x, y, z)$
T	T	T	∞
F	F	F	∞
*	*	*	0

Valued CSPs

Let Γ be a finite set of functions $\phi : D^m \rightarrow \overline{\mathbb{Q}}$. An instance of $\text{VCSP}(\Gamma)$ is an optimisation problem of the form

$$\min_{x_1, \dots, x_n \in D} \phi_1(x_{1,1}, \dots, x_{1,m_1}) + \dots + \phi_q(x_{q,1}, \dots, x_{q,m_q})$$

where $x_{i,j} \in \{x_1, \dots, x_n\}$ and all functions $\phi_i \in \Gamma$.

(s, t) -Min-Cut

$$\begin{array}{ll} \min_{x_1, \dots, x_n \in \{a, b\}} & \sum_{(x_i, x_j) \in E(G)} \phi_{ij}(x_i, x_j) \\ \text{s.t.} & x_s = a, x_t = b \end{array}$$

x	y	$\phi_{ij}(x, y)$
a	a	0
a	b	1
b	a	1
b	b	0

Valued CSPs

Let Γ be a finite set of functions $\phi : D^m \rightarrow \overline{\mathbb{Q}}$. An instance of $\text{VCSP}(\Gamma)$ is an optimisation problem of the form

$$\min_{x_1, \dots, x_n \in D} \phi_1(x_{1,1}, \dots, x_{1,m_1}) + \dots + \phi_q(x_{q,1}, \dots, x_{q,m_q})$$

where $x_{i,j} \in \{x_1, \dots, x_n\}$ and all functions $\phi_i \in \Gamma$.

Max-Cut

$$\min_{x_1, \dots, x_n \in \{a, b\}} \sum_{(x_i, x_j) \in E(G)} \phi_{\neq}(x_i, x_j)$$

x	y	$\phi_{\neq}(x, y)$
a	a	1
a	b	0
b	a	0
b	b	1

Valued CSPs

Let Γ be a finite set of functions $\phi : D^m \rightarrow \overline{\mathbb{Q}}$. An instance of $\text{VCSP}(\Gamma)$ is an optimisation problem of the form

$$\min_{x_1, \dots, x_n \in D} \phi_1(x_{1,1}, \dots, x_{1,m_1}) + \dots + \phi_q(x_{q,1}, \dots, x_{q,m_q})$$

where $x_{i,j} \in \{x_1, \dots, x_n\}$ and all functions $\phi_i \in \Gamma$.

Vertex Cover

$$\min_{x_1, \dots, x_n \in \{0,1\}} \left(\sum_{(x_i, x_j) \in E(G)} \phi_{\text{vc}}(x_i, x_j) + \sum_{x_i \in V(G)} \eta(x_i) \right)$$

x	$\eta(x)$
0	0
1	1

x	y	$\phi_{\text{vc}}(x, y)$
0	0	∞
*	*	0

Valued CSPs

Let Γ be a finite set of functions $\phi : D^m \rightarrow \overline{\mathbb{Q}}$. An instance of $\text{VCSP}(\Gamma)$ is an optimisation problem of the form

$$\min_{x_1, \dots, x_n \in D} \phi_1(x_{1,1}, \dots, x_{1,m_1}) + \dots + \phi_q(x_{q,1}, \dots, x_{q,m_q})$$

where $x_{i,j} \in \{x_1, \dots, x_n\}$ and all functions $\phi_i \in \Gamma$.

Codomain of functions from Γ :

Valued CSPs

Let Γ be a finite set of functions $\phi : D^m \rightarrow \overline{\mathbb{Q}}$. An instance of VCSP(Γ) is an optimisation problem of the form

$$\min_{x_1, \dots, x_n \in D} \phi_1(x_{1,1}, \dots, x_{1,m_1}) + \dots + \phi_q(x_{q,1}, \dots, x_{q,m_q})$$

where $x_{i,j} \in \{x_1, \dots, x_n\}$ and all functions $\phi_i \in \Gamma$.

Codomain of functions from Γ :

- ▶ $\{0, \infty\}$: CSPs (feasibility)

Valued CSPs

Let Γ be a finite set of functions $\phi : D^m \rightarrow \overline{\mathbb{Q}}$. An instance of VCSP(Γ) is an optimisation problem of the form

$$\min_{x_1, \dots, x_n \in D} \phi_1(x_{1,1}, \dots, x_{1,m_1}) + \dots + \phi_q(x_{q,1}, \dots, x_{q,m_q})$$

where $x_{i,j} \in \{x_1, \dots, x_n\}$ and all functions $\phi_i \in \Gamma$.

Codomain of functions from Γ :

- ▶ $\{0, \infty\}$: CSPs (feasibility)
- ▶ $\{0, 1\}$: Max-CSPs (max #satisfied constraints)

Valued CSPs

Let Γ be a finite set of functions $\phi : D^m \rightarrow \overline{\mathbb{Q}}$. An instance of $\text{VCSP}(\Gamma)$ is an optimisation problem of the form

$$\min_{x_1, \dots, x_n \in D} \phi_1(x_{1,1}, \dots, x_{1,m_1}) + \dots + \phi_q(x_{q,1}, \dots, x_{q,m_q})$$

where $x_{i,j} \in \{x_1, \dots, x_n\}$ and all functions $\phi_i \in \Gamma$.

Codomain of functions from Γ :

- ▶ $\{0, \infty\}$: CSPs (feasibility)
- ▶ $\{0, 1\}$: Max-CSPs (max #satisfied constraints)
- ▶ \mathbb{Q} : Finite-Valued CSPs (optimisation)

Valued CSPs

Let Γ be a finite set of functions $\phi : D^m \rightarrow \overline{\mathbb{Q}}$. An instance of VCSP(Γ) is an optimisation problem of the form

$$\min_{x_1, \dots, x_n \in D} \phi_1(x_{1,1}, \dots, x_{1,m_1}) + \dots + \phi_q(x_{q,1}, \dots, x_{q,m_q})$$

where $x_{i,j} \in \{x_1, \dots, x_n\}$ and all functions $\phi_i \in \Gamma$.

Codomain of functions from Γ :

- ▶ $\{0, \infty\}$: CSPs (feasibility)
- ▶ $\{0, 1\}$: Max-CSPs (max #satisfied constraints)
- ▶ \mathbb{Q} : Finite-Valued CSPs (optimisation)
- ▶ $\overline{\mathbb{Q}}$: (General-)Valued CSPs (feasibility & optimisation)

Valued CSPs

Let Γ be a finite set of functions $\phi : D^m \rightarrow \overline{\mathbb{Q}}$. An instance of $\text{VCSP}(\Gamma)$ is an optimisation problem of the form

$$\min_{x_1, \dots, x_n \in D} \phi_1(x_{1,1}, \dots, x_{1,m_1}) + \dots + \phi_q(x_{q,1}, \dots, x_{q,m_q})$$

where $x_{i,j} \in \{x_1, \dots, x_n\}$ and all functions $\phi_i \in \Gamma$.

Codomain of functions from Γ :

- ▶ $\{0, \infty\}$: CSPs (feasibility)
- ▶ $\{0, 1\}$: Max-CSPs (max #satisfied constraints)
- ▶ \mathbb{Q} : Finite-Valued CSPs (optimisation)
- ▶ $\overline{\mathbb{Q}}$: (General-)Valued CSPs (feasibility & optimisation)

Valued CSPs \Leftrightarrow Weighted Clones

[Cohen, Cooper, Creed, Jeavons, Ž. SICOMP'13]

This Talk

Valued CSPs \Leftrightarrow Weighted Clones

[Cohen, Cooper, Creed, Jeavons, Ž. SICOMP'13]

Outline

operations \rightarrow weightings

clones \rightarrow weighted clones

polymorphisms \rightarrow weighted polymorphisms

Operations and Clones

A k -ary **operation** on D is a mapping $f : D^k \rightarrow D$.

A **clone** C on D is a set of operations on D closed under superposition and containing all projections.

Weightings

A k -ary **weighting** of a clone C is a function $\omega : C^{(k)} \rightarrow \mathbb{Q}$ such that $\omega(f) < 0$ only if f is a projection and

$$\sum_{f \in C^{(k)}} \omega(f) = 0.$$

Weightings

A k -ary **weighting** of a clone C is a function $\omega : C^{(k)} \rightarrow \mathbb{Q}$ such that $\omega(f) < 0$ only if f is a projection and

$$\sum_{f \in C^{(k)}} \omega(f) = 0.$$

- ▶ $\omega(f) = c\omega'(f)$ for every $f \in C^{(k)}$ (scaling by $c \in \mathbb{Q}_{\geq 0}$)

Weightings

A k -ary **weighting** of a clone C is a function $\omega : C^{(k)} \rightarrow \mathbb{Q}$ such that $\omega(f) < 0$ only if f is a projection and

$$\sum_{f \in C^{(k)}} \omega(f) = 0.$$

- ▶ $\omega(f) = c\omega'(f)$ for every $f \in C^{(k)}$ (scaling by $c \in \mathbb{Q}_{\geq 0}$)
- ▶ $\omega(f) = \omega_1(f) + \omega_2(f)$ (addition)

Weightings

A k -ary **weighting** of a clone C is a function $\omega : C^{(k)} \rightarrow \mathbb{Q}$ such that $\omega(f) < 0$ only if f is a projection and

$$\sum_{f \in C^{(k)}} \omega(f) = 0.$$

- ▶ $\omega(f) = c\omega'(f)$ for every $f \in C^{(k)}$ (scaling by $c \in \mathbb{Q}_{\geq 0}$)
- ▶ $\omega(f) = \omega_1(f) + \omega_2(f)$ (addition)
- ▶ $\omega[g_1, \dots, g_k](f') = \sum_{\substack{f \in C^{(k)} \\ f[g_1, \dots, g_k] = f'}} \omega(f)$ (superposition)

Superposition Example

- ▶ D totally ordered

Superposition Example

- ▶ D totally ordered
- ▶ binary min, max $\in C$

Superposition Example

- ▶ D totally ordered
- ▶ binary $\min, \max \in C$
- ▶ 4-ary weighting ω

$$\omega(f) = \begin{cases} -1 & \text{if } f \in \{e_1^{(4)}, e_2^{(4)}, e_3^{(4)}, e_4^{(4)}\} \\ +1 & \text{if } f \in \{\max_{12}^{(4)}, \min_{12}^{(4)}, \max_{34}^{(4)}, \min_{34}^{(4)}\} \\ 0 & \text{o/w} \end{cases}$$

Superposition Example

- ▶ D totally ordered
- ▶ binary $\min, \max \in C$
- ▶ 4-ary weighting ω

$$\omega(f) = \begin{cases} -1 & \text{if } f \in \{e_1^{(4)}, e_2^{(4)}, e_3^{(4)}, e_4^{(4)}\} \\ +1 & \text{if } f \in \{\max_{12}^{(4)}, \min_{12}^{(4)}, \max_{34}^{(4)}, \min_{34}^{(4)}\} \\ 0 & \text{o/w} \end{cases}$$

$$\langle g_1, g_2, g_3, g_4 \rangle = \langle e_1^{(3)}, e_2^{(3)}, e_3^{(3)}, \max_{12}^{(3)} \rangle$$

Superposition Example

- ▶ D totally ordered
- ▶ binary $\min, \max \in C$
- ▶ 4-ary weighting ω

$$\omega(f) = \begin{cases} -1 & \text{if } f \in \{e_1^{(4)}, e_2^{(4)}, e_3^{(4)}, e_4^{(4)}\} \\ +1 & \text{if } f \in \{\max_{12}^{(4)}, \min_{12}^{(4)}, \max_{34}^{(4)}, \min_{34}^{(4)}\} \\ 0 & \text{o/w} \end{cases}$$

$$\langle g_1, g_2, g_3, g_4 \rangle = \langle e_1^{(3)}, e_2^{(3)}, e_3^{(3)}, \max_{12}^{(3)} \rangle$$

$$e_1^{(4)}[g_1, g_2, g_3, g_4] = g_1 = e_1^{(3)}$$

Superposition Example

- ▶ D totally ordered
- ▶ binary $\min, \max \in C$
- ▶ 4-ary weighting ω

$$\omega(f) = \begin{cases} -1 & \text{if } f \in \{e_1^{(4)}, e_2^{(4)}, e_3^{(4)}, e_4^{(4)}\} \\ +1 & \text{if } f \in \{\max_{12}^{(4)}, \min_{12}^{(4)}, \max_{34}^{(4)}, \min_{34}^{(4)}\} \\ 0 & \text{o/w} \end{cases}$$

$$\langle g_1, g_2, g_3, g_4 \rangle = \langle e_1^{(3)}, e_2^{(3)}, e_3^{(3)}, \max_{12}^{(3)} \rangle$$

$$e_2^{(4)}[g_1, g_2, g_3, g_4] = g_2 = e_2^{(3)}$$

Superposition Example

- ▶ D totally ordered
- ▶ binary $\min, \max \in C$
- ▶ 4-ary weighting ω

$$\omega(f) = \begin{cases} -1 & \text{if } f \in \{e_1^{(4)}, e_2^{(4)}, e_3^{(4)}, e_4^{(4)}\} \\ +1 & \text{if } f \in \{\max_{12}^{(4)}, \min_{12}^{(4)}, \max_{34}^{(4)}, \min_{34}^{(4)}\} \\ 0 & \text{o/w} \end{cases}$$

$$\langle g_1, g_2, g_3, g_4 \rangle = \langle e_1^{(3)}, e_2^{(3)}, e_3^{(3)}, \max_{12}^{(3)} \rangle$$

$$e_3^{(4)}[g_1, g_2, g_3, g_4] = g_3 = e_3^{(3)}$$

Superposition Example

- ▶ D totally ordered
- ▶ binary $\min, \max \in C$
- ▶ 4-ary weighting ω

$$\omega(f) = \begin{cases} -1 & \text{if } f \in \{e_1^{(4)}, e_2^{(4)}, e_3^{(4)}, e_4^{(4)}\} \\ +1 & \text{if } f \in \{\max_{12}^{(4)}, \min_{12}^{(4)}, \max_{34}^{(4)}, \min_{34}^{(4)}\} \\ 0 & \text{o/w} \end{cases}$$

$$\langle g_1, g_2, g_3, g_4 \rangle = \langle e_1^{(3)}, e_2^{(3)}, e_3^{(3)}, \max_{12}^{(3)} \rangle$$

$$e_4^{(4)}[g_1, g_2, g_3, g_4] = g_4 = \max_{12}^{(3)}$$

Superposition Example

- ▶ D totally ordered
- ▶ binary $\min, \max \in C$
- ▶ 4-ary weighting ω

$$\omega(f) = \begin{cases} -1 & \text{if } f \in \{e_1^{(4)}, e_2^{(4)}, e_3^{(4)}, e_4^{(4)}\} \\ +1 & \text{if } f \in \{\text{max}_{12}^{(4)}, \min_{12}^{(4)}, \max_{34}^{(4)}, \min_{34}^{(4)}\} \\ 0 & \text{o/w} \end{cases}$$

$$\langle g_1, g_2, g_3, g_4 \rangle = \langle e_1^{(3)}, e_2^{(3)}, e_3^{(3)}, \max_{12}^{(3)} \rangle$$

$$\text{max}_{12}^{(4)}[g_1, g_2, g_3, g_4] = \max_{12}^{(3)}$$

Superposition Example

- ▶ D totally ordered
- ▶ binary $\min, \max \in C$
- ▶ 4-ary weighting ω

$$\omega(f) = \begin{cases} -1 & \text{if } f \in \{e_1^{(4)}, e_2^{(4)}, e_3^{(4)}, e_4^{(4)}\} \\ +1 & \text{if } f \in \{\max_{12}^{(4)}, \textcolor{red}{\min}_{12}^{(4)}, \max_{34}^{(4)}, \min_{34}^{(4)}\} \\ 0 & \text{o/w} \end{cases}$$

$$\langle g_1, g_2, g_3, g_4 \rangle = \langle e_1^{(3)}, e_2^{(3)}, e_3^{(3)}, \max_{12}^{(3)} \rangle$$

$$\textcolor{red}{\min}_{12}^{(4)}[g_1, g_2, g_3, g_4] = \min_{12}^{(3)}$$

Superposition Example

- ▶ D totally ordered
- ▶ binary $\min, \max \in C$
- ▶ 4-ary weighting ω

$$\omega(f) = \begin{cases} -1 & \text{if } f \in \{e_1^{(4)}, e_2^{(4)}, e_3^{(4)}, e_4^{(4)}\} \\ +1 & \text{if } f \in \{\max_{12}^{(4)}, \min_{12}^{(4)}, \textcolor{red}{\max}_{34}^{(4)}, \min_{34}^{(4)}\} \\ 0 & \text{o/w} \end{cases}$$

$$\langle g_1, g_2, g_3, g_4 \rangle = \langle e_1^{(3)}, e_2^{(3)}, e_3^{(3)}, \max_{12}^{(3)} \rangle$$

$$\textcolor{red}{\max}_{34}^{(4)}[g_1, g_2, g_3, g_4] = \max_{123}^{(3)}$$

Superposition Example

- ▶ D totally ordered
- ▶ binary $\min, \max \in C$
- ▶ 4-ary weighting ω

$$\omega(f) = \begin{cases} -1 & \text{if } f \in \{e_1^{(4)}, e_2^{(4)}, e_3^{(4)}, e_4^{(4)}\} \\ +1 & \text{if } f \in \{\max_{12}^{(4)}, \min_{12}^{(4)}, \max_{34}^{(4)}, \min_{34}^{(4)}\} \\ 0 & \text{o/w} \end{cases}$$

$$\langle g_1, g_2, g_3, g_4 \rangle = \langle e_1^{(3)}, e_2^{(3)}, e_3^{(3)}, \max_{12}^{(3)} \rangle$$

$$\min_{34}^{(4)}[g_1, g_2, g_3, g_4] = \min^{(3)}(\max_{12}^{(3)}, e_3^{(3)})$$

Superposition Example

- ▶ D totally ordered
- ▶ binary $\min, \max \in C$
- ▶ 4-ary weighting ω

$$\omega(f) = \begin{cases} -1 & \text{if } f \in \{e_1^{(4)}, e_2^{(4)}, e_3^{(4)}, e_4^{(4)}\} \\ +1 & \text{if } f \in \{\max_{12}^{(4)}, \min_{12}^{(4)}, \max_{34}^{(4)}, \min_{34}^{(4)}\} \\ 0 & \text{o/w} \end{cases}$$

$$\langle g_1, g_2, g_3, g_4 \rangle = \langle e_1^{(3)}, e_2^{(3)}, e_3^{(3)}, \max_{12}^{(3)} \rangle$$

$$\omega' = \omega[g_1, g_2, g_3, g_4]$$

Superposition Example

- ▶ D totally ordered
- ▶ binary $\min, \max \in C$
- ▶ 4-ary weighting ω

$$\omega(f) = \begin{cases} -1 & \text{if } f \in \{e_1^{(4)}, e_2^{(4)}, e_3^{(4)}, e_4^{(4)}\} \\ +1 & \text{if } f \in \{\max_{12}^{(4)}, \min_{12}^{(4)}, \max_{34}^{(4)}, \min_{34}^{(4)}\} \\ 0 & \text{o/w} \end{cases}$$

$$\langle g_1, g_2, g_3, g_4 \rangle = \langle e_1^{(3)}, e_2^{(3)}, e_3^{(3)}, \max_{12}^{(3)} \rangle$$

$$\omega' = \omega[g_1, g_2, g_3, g_4]$$

$$\omega'(f) = \begin{cases} -1 & \text{if } f \in \{e_1^{(3)}, e_2^{(3)}, e_3^{(3)}\} \\ +1 & \text{if } f \in \{\max_{123}^{(3)}, \min_{12}^{(3)}, \min^{(3)}(\max_{12}^{(3)}, e_3^{(3)})\} \\ 0 & \text{o/w} \end{cases}$$

Superposition as Matrix Multiplication

$$\omega = (f_1, \dots, f_{|C^{(k)}|})$$

Superposition as Matrix Multiplication

$$\omega = (f_1, \dots, f_{|C^{(k)}|}) \quad g_i \in C^{(\ell)} \text{ for } 1 \leq i \leq k$$

Superposition as Matrix Multiplication

$$\begin{aligned}\omega &= (f_1, \dots, f_{|C^{(k)}|}) \\ \omega[g_1, \dots, g_k] &= (h_1, \dots, h_{|C^{(\ell)}|})\end{aligned}$$

$$g_i \in C^{(\ell)} \text{ for } 1 \leq i \leq k$$

Superposition as Matrix Multiplication

$$\omega = (f_1, \dots, f_{|C^{(k)}|})$$
$$\omega[g_1, \dots, g_k] = (h_1, \dots, h_{|C^{(\ell)}|})$$

$$g_i \in C^{(\ell)} \text{ for } 1 \leq i \leq k$$

$$\left(\begin{array}{c} \text{G} \end{array} \right)_{|C^{(k)}| \times |C^{(\ell)}|}$$

Superposition as Matrix Multiplication

$$\omega = (f_1, \dots, f_{|C^{(k)}|})$$
$$\omega[g_1, \dots, g_k] = (h_1, \dots, h_{|C^{(\ell)}|})$$

$$g_i \in C^{(\ell)} \text{ for } 1 \leq i \leq k$$

$$f \begin{pmatrix} h \\ 1 \\ \mathbf{G} \end{pmatrix}_{|C^{(k)}| \times |C^{(\ell)}|}$$

$$f[g_1, \dots, g_k] = h$$

Superposition as Matrix Multiplication

$$\omega = (f_1, \dots, f_{|C^{(k)}|})$$
$$\omega[g_1, \dots, g_k] = (h_1, \dots, h_{|C^{(\ell)}|})$$

$$g_i \in C^{(\ell)} \text{ for } 1 \leq i \leq k$$

$$\left(\begin{array}{c} \text{G} \end{array} \right)_{|C^{(k)}| \times |C^{(\ell)}|}$$

Superposition as Matrix Multiplication

$$\begin{aligned}\omega &= (f_1, \dots, f_{|C^{(k)}|}) \\ \omega[g_1, \dots, g_k] &= (h_1, \dots, h_{|C^{(\ell)}|})\end{aligned}$$

$$g_i \in C^{(\ell)} \text{ for } 1 \leq i \leq k$$

$$\omega[g_1, \dots, g_k] = \omega \cdot \left(\begin{array}{c} \text{G} \\ |C^{(k)}| \times |C^{(\ell)}| \end{array} \right)$$

Superposition Example, cont'd

- ▶ D totally ordered
- ▶ binary $\min, \max \in C$
- ▶ 4-ary weighting ω

$$\omega(f) = \begin{cases} -1 & \text{if } f \in \{e_1^{(4)}, e_2^{(4)}, e_3^{(4)}, e_4^{(4)}\} \\ +1 & \text{if } f \in \{\max_{12}^{(4)}, \min_{12}^{(4)}, \max_{34}^{(4)}, \min_{34}^{(4)}\} \\ 0 & \text{o/w} \end{cases}$$

$$\langle g_1, g_2, g_3, g_4 \rangle = \langle e_1^{(4)}, \max_{23}^{(4)}, \min_{23}^{(4)}, e_4^{(4)} \rangle$$

Superposition Example, cont'd

- ▶ D totally ordered
- ▶ binary $\min, \max \in C$
- ▶ 4-ary weighting ω

$$\omega(f) = \begin{cases} -1 & \text{if } f \in \{e_1^{(4)}, e_2^{(4)}, e_3^{(4)}, e_4^{(4)}\} \\ +1 & \text{if } f \in \{\max_{12}^{(4)}, \min_{12}^{(4)}, \max_{34}^{(4)}, \min_{34}^{(4)}\} \\ 0 & \text{o/w} \end{cases}$$

$$\langle g_1, g_2, g_3, g_4 \rangle = \langle e_1^{(4)}, \max_{23}^{(4)}, \min_{23}^{(4)}, e_4^{(4)} \rangle$$

$$\omega' = \omega[g_1, g_2, g_3, g_4]$$

Superposition Example, cont'd

- ▶ D totally ordered
- ▶ binary $\min, \max \in C$
- ▶ 4-ary weighting ω

$$\omega(f) = \begin{cases} -1 & \text{if } f \in \{e_1^{(4)}, e_2^{(4)}, e_3^{(4)}, e_4^{(4)}\} \\ +1 & \text{if } f \in \{\max_{12}^{(4)}, \min_{12}^{(4)}, \max_{34}^{(4)}, \min_{34}^{(4)}\} \\ 0 & \text{o/w} \end{cases}$$

$$\langle g_1, g_2, g_3, g_4 \rangle = \langle e_1^{(4)}, \max_{23}^{(4)}, \min_{23}^{(4)}, e_4^{(4)} \rangle$$

$$\omega' = \omega[g_1, g_2, g_3, g_4]$$

$$\omega'(f) = \begin{cases} -1 & \text{if } f \in \{e_1^{(3)}, \max_{23}^{(4)}, \min_{23}^{(4)}, e_4^{(4)}\} \\ +1 & \text{if } f \in \left\{ \begin{array}{l} \max_{123}^{(4)}, \min_{123}^{(4)}(e_1^{(4)}, \max_{23}^{(4)}), \\ \max_{234}^{(4)}(\min_{23}^{(4)}, e_4^{(4)}), \min_{234}^{(4)} \end{array} \right\} \\ 0 & \text{o/w} \end{cases}$$

Superposition Example, cont'd

- ▶ D totally ordered
- ▶ binary $\min, \max \in C$
- ▶ 4-ary weighting ω

$$\omega(f) = \begin{cases} -1 & \text{if } f \in \{e_1^{(4)}, e_2^{(4)}, e_3^{(4)}, e_4^{(4)}\} \\ +1 & \text{if } f \in \{\max_{12}^{(4)}, \min_{12}^{(4)}, \max_{34}^{(4)}, \min_{34}^{(4)}\} \\ 0 & \text{o/w} \end{cases}$$

$$\langle g_1, g_2, g_3, g_4 \rangle = \langle e_1^{(4)}, \max_{23}^{(4)}, \min_{23}^{(4)}, e_4^{(4)} \rangle$$

$$\omega' = \omega[g_1, g_2, g_3, g_4]$$

$$\omega'(f) = \begin{cases} -1 & \text{if } f \in \{e_1^{(3)}, \max_{23}^{(4)}, \min_{23}^{(4)}, e_4^{(4)}\} \\ +1 & \text{if } f \in \left\{ \begin{array}{l} \max_{123}^{(4)}, \min_{123}^{(4)}(e_1^{(4)}, \max_{23}^{(4)}), \\ \max_{234}^{(4)}(\min_{23}^{(4)}, e_4^{(4)}), \min_{234}^{(4)} \end{array} \right\} \\ 0 & \text{o/w} \end{cases}$$

Weightings

A k -ary **weighting** of a clone C is a function $\omega : C^{(k)} \rightarrow \mathbb{Q}$ such that $\omega(f) < 0$ only if f is a projection and

$$\sum_{f \in C^{(k)}} \omega(f) = 0.$$

- ▶ $\omega(f) = c\omega'(f)$ for every $f \in C^{(k)}$ (scaling by $c \in \mathbb{Q}_{\geq 0}$)
- ▶ $\omega(f) = \omega_1(f) + \omega_2(f)$ (addition)
- ▶ $\omega[g_1, \dots, g_k](f') = \sum_{\substack{f \in C^{(k)} \\ f[g_1, \dots, g_k] = f'}} \omega(f)$ (**proper** superposition)

Weightings

A k -ary **weighting** of a clone C is a function $\omega : C^{(k)} \rightarrow \mathbb{Q}$ such that $\omega(f) < 0$ only if f is a projection and

$$\sum_{f \in C^{(k)}} \omega(f) = 0.$$

Lemma: Improper superpositions can be eliminated!

- ▶ $\omega(f) = c\omega'(f)$ for every $f \in C^{(k)}$ (scaling by $c \in \mathbb{Q}_{\geq 0}$)
- ▶ $\omega(f) = \omega_1(f) + \omega_2(f)$ (addition)
- ▶ $\omega[g_1, \dots, g_k](f') = \sum_{\substack{f \in C^{(k)} \\ f[g_1, \dots, g_k] = f'}} \omega(f)$ (**proper** superposition)

Weighted Clones

A **weighted clone** W is a non-empty set of weightings of the support clone C which is closed under non-negative scaling, addition of weightings of equal arity, and proper superpositions with operations from C .

Weighted Clones

A **weighted clone** W is a non-empty set of weightings of the support clone C which is closed under non-negative scaling, addition of weightings of equal arity, and proper superpositions with operations from C .

- ▶ W_C : all possible weightings of C

Weighted Clones

A **weighted clone** W is a non-empty set of weightings of the support clone C which is closed under non-negative scaling, addition of weightings of equal arity, and proper superpositions with operations from C .

- ▶ W_C : all possible weightings of C
- ▶ W_C^0 : 0-valued weightings of C
(i.e., for each arity $k \geq 1$, $\omega_k(f) = 0$ for all $f \in C^{(k)}$)

Weighted Clones

A **weighted clone** W is a non-empty set of weightings of the support clone C which is closed under non-negative scaling, addition of weightings of equal arity, and proper superpositions with operations from C .

- ▶ W_C : all possible weightings of C
- ▶ W_C^0 : 0-valued weightings of C
(i.e., for each arity $k \geq 1$, $\omega_k(f) = 0$ for all $f \in C^{(k)}$)

Structure of Weighted Clones!

Valued CSPs \Leftrightarrow Weighted Clones

[Cohen, Cooper, Creed, Jeavons, Ž. SICOMP'13]

This Talk

Valued CSPs \Leftrightarrow Weighted Clones

[Cohen, Cooper, Creed, Jeavons, Ž. SICOMP'13]

Polymorphisms

Let $\phi : D^m \rightarrow \overline{\mathbb{Q}}$ and let $\text{Feas}(\phi) = \{\mathbf{x} \in D^m \mid \phi(\mathbf{x}) \text{ is finite}\}$.
An operation $f : D^k \rightarrow D$ is a **polymorphism** of ϕ if, for any $\mathbf{x}_1, \dots, \mathbf{x}_k \in \text{Feas}(\phi)$ we have $f(\mathbf{x}_1, \dots, \mathbf{x}_k) \in \text{Feas}(\phi)$.

Polymorphisms

Let $\phi : D^m \rightarrow \overline{\mathbb{Q}}$ and let $\text{Feas}(\phi) = \{\mathbf{x} \in D^m \mid \phi(\mathbf{x}) \text{ is finite}\}$.
An operation $f : D^k \rightarrow D$ is a **polymorphism** of ϕ if, for any $\mathbf{x}_1, \dots, \mathbf{x}_k \in \text{Feas}(\phi)$ we have $f(\mathbf{x}_1, \dots, \mathbf{x}_k) \in \text{Feas}(\phi)$.

$$D = \{0, 1\}, m = 3$$

Polymorphisms

Let $\phi : D^m \rightarrow \overline{\mathbb{Q}}$ and let $\text{Feas}(\phi) = \{\mathbf{x} \in D^m \mid \phi(\mathbf{x}) \text{ is finite}\}$.
An operation $f : D^k \rightarrow D$ is a **polymorphism** of ϕ if, for any $\mathbf{x}_1, \dots, \mathbf{x}_k \in \text{Feas}(\phi)$ we have $f(\mathbf{x}_1, \dots, \mathbf{x}_k) \in \text{Feas}(\phi)$.

$D = \{0, 1\}$, $m = 3$

x	y	z	$\phi(x, y, z)$
0	0	0	∞
0	0	1	5
0	1	0	∞
0	1	1	2
1	0	0	∞
1	0	1	1
1	1	0	0
1	1	1	∞

Polymorphisms

Let $\phi : D^m \rightarrow \overline{\mathbb{Q}}$ and let $\text{Feas}(\phi) = \{\mathbf{x} \in D^m \mid \phi(\mathbf{x}) \text{ is finite}\}$.
An operation $f : D^k \rightarrow D$ is a **polymorphism** of ϕ if, for any $\mathbf{x}_1, \dots, \mathbf{x}_k \in \text{Feas}(\phi)$ we have $f(\mathbf{x}_1, \dots, \mathbf{x}_k) \in \text{Feas}(\phi)$.

$D = \{0, 1\}$, $m = 3$, $k = 2$, $f = \min$

x	y	z	$\phi(x, y, z)$
0	0	0	∞
0	0	1	5
0	1	0	∞
0	1	1	2
1	0	0	∞
1	0	1	1
1	1	0	0
1	1	1	∞

Polymorphisms

Let $\phi : D^m \rightarrow \overline{\mathbb{Q}}$ and let $\text{Feas}(\phi) = \{\mathbf{x} \in D^m \mid \phi(\mathbf{x}) \text{ is finite}\}$.
An operation $f : D^k \rightarrow D$ is a **polymorphism** of ϕ if, for any $\mathbf{x}_1, \dots, \mathbf{x}_k \in \text{Feas}(\phi)$ we have $f(\mathbf{x}_1, \dots, \mathbf{x}_k) \in \text{Feas}(\phi)$.

$D = \{0, 1\}$, $m = 3$, $k = 2$, $f = \min$

x	y	z	$\phi(x, y, z)$
0	0	0	∞
0	0	1	5
0	1	0	∞
0	1	1	2
1	0	0	∞
1	0	1	1
1	1	0	0
1	1	1	∞

$$\mathbf{x}_1 = (0, 1, 1)$$

Polymorphisms

Let $\phi : D^m \rightarrow \overline{\mathbb{Q}}$ and let $\text{Feas}(\phi) = \{\mathbf{x} \in D^m \mid \phi(\mathbf{x}) \text{ is finite}\}$.
An operation $f : D^k \rightarrow D$ is a **polymorphism** of ϕ if, for any $\mathbf{x}_1, \dots, \mathbf{x}_k \in \text{Feas}(\phi)$ we have $f(\mathbf{x}_1, \dots, \mathbf{x}_k) \in \text{Feas}(\phi)$.

$D = \{0, 1\}$, $m = 3$, $k = 2$, $f = \min$

x	y	z	$\phi(x, y, z)$
0	0	0	∞
0	0	1	5
0	1	0	∞
0	1	1	2
1	0	0	∞
1	0	1	1
1	1	0	0
1	1	1	∞

$$\mathbf{x}_1 = (0, 1, 1) \in \text{Feas}(\phi)$$

Polymorphisms

Let $\phi : D^m \rightarrow \overline{\mathbb{Q}}$ and let $\text{Feas}(\phi) = \{\mathbf{x} \in D^m \mid \phi(\mathbf{x}) \text{ is finite}\}$.
An operation $f : D^k \rightarrow D$ is a **polymorphism** of ϕ if, for any $\mathbf{x}_1, \dots, \mathbf{x}_k \in \text{Feas}(\phi)$ we have $f(\mathbf{x}_1, \dots, \mathbf{x}_k) \in \text{Feas}(\phi)$.

$D = \{0, 1\}$, $m = 3$, $k = 2$, $f = \min$

x	y	z	$\phi(x, y, z)$
0	0	0	∞
0	0	1	5
0	1	0	∞
0	1	1	2
1	0	0	∞
1	0	1	1
1	1	0	0
1	1	1	∞

$$\begin{aligned}\mathbf{x}_1 &= (0, 1, 1) \in \text{Feas}(\phi) \\ \mathbf{x}_2 &= (1, 0, 1)\end{aligned}$$

Polymorphisms

Let $\phi : D^m \rightarrow \overline{\mathbb{Q}}$ and let $\text{Feas}(\phi) = \{\mathbf{x} \in D^m \mid \phi(\mathbf{x}) \text{ is finite}\}$.
An operation $f : D^k \rightarrow D$ is a **polymorphism** of ϕ if, for any $\mathbf{x}_1, \dots, \mathbf{x}_k \in \text{Feas}(\phi)$ we have $f(\mathbf{x}_1, \dots, \mathbf{x}_k) \in \text{Feas}(\phi)$.

$D = \{0, 1\}$, $m = 3$, $k = 2$, $f = \min$

x	y	z	$\phi(x, y, z)$
0	0	0	∞
0	0	1	5
0	1	0	∞
0	1	1	2
1	0	0	∞
1	0	1	1
1	1	0	0
1	1	1	∞

$$\mathbf{x}_1 = (0, 1, 1) \in \text{Feas}(\phi)$$

$$\mathbf{x}_2 = (1, 0, 1) \in \text{Feas}(\phi)$$

Polymorphisms

Let $\phi : D^m \rightarrow \overline{\mathbb{Q}}$ and let $\text{Feas}(\phi) = \{\mathbf{x} \in D^m \mid \phi(\mathbf{x}) \text{ is finite}\}$.
An operation $f : D^k \rightarrow D$ is a **polymorphism** of ϕ if, for any $\mathbf{x}_1, \dots, \mathbf{x}_k \in \text{Feas}(\phi)$ we have $f(\mathbf{x}_1, \dots, \mathbf{x}_k) \in \text{Feas}(\phi)$.

$D = \{0, 1\}$, $m = 3$, $k = 2$, $f = \min$

x	y	z	$\phi(x, y, z)$
0	0	0	∞
0	0	1	5
0	1	0	∞
0	1	1	2
1	0	0	∞
1	0	1	1
1	1	0	0
1	1	1	∞

$$\begin{array}{rcl} \mathbf{x}_1 & = & (0, 1, 1) \in \text{Feas}(\phi) \\ \mathbf{x}_2 & = & (1, 0, 1) \in \text{Feas}(\phi) \\ \hline f(\mathbf{x}_1, \mathbf{x}_2) & = & (0, 0, 1) \end{array}$$

Polymorphisms

Let $\phi : D^m \rightarrow \overline{\mathbb{Q}}$ and let $\text{Feas}(\phi) = \{\mathbf{x} \in D^m \mid \phi(\mathbf{x}) \text{ is finite}\}$.
An operation $f : D^k \rightarrow D$ is a **polymorphism** of ϕ if, for any $\mathbf{x}_1, \dots, \mathbf{x}_k \in \text{Feas}(\phi)$ we have $f(\mathbf{x}_1, \dots, \mathbf{x}_k) \in \text{Feas}(\phi)$.

$D = \{0, 1\}$, $m = 3$, $k = 2$, $f = \min$

x	y	z	$\phi(x, y, z)$
0	0	0	∞
0	0	1	5
0	1	0	∞
0	1	1	2
1	0	0	∞
1	0	1	1
1	1	0	0
1	1	1	∞

$$\begin{array}{rcll} \mathbf{x}_1 & = & (0, 1, 1) & \in \text{Feas}(\phi) \\ \mathbf{x}_2 & = & (1, 0, 1) & \in \text{Feas}(\phi) \\ \hline f(\mathbf{x}_1, \mathbf{x}_2) & = & (0, 0, 1) & \in \text{Feas}(\phi) \end{array}$$

Polymorphisms

Let $\phi : D^m \rightarrow \overline{\mathbb{Q}}$ and let $\text{Feas}(\phi) = \{\mathbf{x} \in D^m \mid \phi(\mathbf{x}) \text{ is finite}\}$.
An operation $f : D^k \rightarrow D$ is a **polymorphism** of ϕ if, for any $\mathbf{x}_1, \dots, \mathbf{x}_k \in \text{Feas}(\phi)$ we have $f(\mathbf{x}_1, \dots, \mathbf{x}_k) \in \text{Feas}(\phi)$.

$D = \{0, 1\}$, $m = 3$, $k = 2$, $f = \min$

x	y	z	$\phi(x, y, z)$
0	0	0	∞
0	0	1	5
0	1	0	∞
0	1	1	2
1	0	0	∞
1	0	1	1
1	1	0	0
1	1	1	∞

$$\begin{array}{rclcl} \mathbf{x}_1 & = & (1, 0, 1) & \in \text{Feas}(\phi) \\ \mathbf{x}_2 & = & (1, 1, 0) & \in \text{Feas}(\phi) \\ \hline f(\mathbf{x}_1, \mathbf{x}_2) & = & (1, 0, 0) & \notin \text{Feas}(\phi) \end{array}$$

f is **not** a polymorphism of ϕ

Weighted Polymorphisms

Let $\phi : D^m \rightarrow \overline{\mathbb{Q}}$ be a function

Weighted Polymorphisms

Let $\phi : D^m \rightarrow \overline{\mathbb{Q}}$ be a function and let $C \subseteq \text{Pol}(\phi)$ be a clone of operations.

Weighted Polymorphisms

Let $\phi : D^m \rightarrow \overline{\mathbb{Q}}$ be a function and let $C \subseteq \text{Pol}(\phi)$ be a clone of operations. A k -ary weighting $\omega : C^{(k)} \rightarrow \mathbb{Q}$
($\sum_{f \in C^{(k)}} \omega(f) = 0$, $\omega(f) < 0$ only if f a projection)

Weighted Polymorphisms

Let $\phi : D^m \rightarrow \overline{\mathbb{Q}}$ be a function and let $C \subseteq \text{Pol}(\phi)$ be a clone of operations. A k -ary weighting $\omega : C^{(k)} \rightarrow \mathbb{Q}$ ($\sum_{f \in C^{(k)}} \omega(f) = 0, \omega(f) < 0$ only if f a projection) is a k -ary **weighted polymorphism** of ϕ if

Weighted Polymorphisms

Let $\phi : D^m \rightarrow \overline{\mathbb{Q}}$ be a function and let $C \subseteq \text{Pol}(\phi)$ be a clone of operations. A k -ary weighting $\omega : C^{(k)} \rightarrow \mathbb{Q}$ ($\sum_{f \in C^{(k)}} \omega(f) = 0$, $\omega(f) < 0$ only if f a projection) is a k -ary **weighted polymorphism** of ϕ if for any $\mathbf{x}_1, \dots, \mathbf{x}_k \in \text{Feas}(\phi)$

$$\sum_{f \in C^{(k)}} \omega(f) \phi(f(\mathbf{x}_1, \dots, \mathbf{x}_k)) \leq 0.$$

Weighted Polymorphisms

Let $\phi : D^m \rightarrow \overline{\mathbb{Q}}$ be a function and let $C \subseteq \text{Pol}(\phi)$ be a clone of operations. A k -ary weighting $\omega : C^{(k)} \rightarrow \mathbb{Q}$ ($\sum_{f \in C^{(k)}} \omega(f) = 0$, $\omega(f) < 0$ only if f a projection) is a k -ary **weighted polymorphism** of ϕ if for any $\mathbf{x}_1, \dots, \mathbf{x}_k \in \text{Feas}(\phi)$

$$\sum_{f \in C^{(k)}} \omega(f) \phi(f(\mathbf{x}_1, \dots, \mathbf{x}_k)) \leq 0.$$

Equivalently, a probability distribution μ over $C^{(k)} \subseteq \text{Pol}^{(k)}(\phi)$

$$\mathbb{E}_{f \sim \mu} [\phi(f(\mathbf{x}_1, \dots, \mathbf{x}_k))] \leq \text{avg}\{\phi(\mathbf{x}_1), \dots, \phi(\mathbf{x}_k)\}.$$

Weighted Polymorphisms

Let $\phi : D^m \rightarrow \overline{\mathbb{Q}}$ be a function and let $C \subseteq \text{Pol}(\phi)$ be a clone of operations. A k -ary weighting $\omega : C^{(k)} \rightarrow \mathbb{Q}$ ($\sum_{f \in C^{(k)}} \omega(f) = 0$, $\omega(f) < 0$ only if f a projection) is a k -ary **weighted polymorphism** of ϕ if for any $\mathbf{x}_1, \dots, \mathbf{x}_k \in \text{Feas}(\phi)$

$$\sum_{f \in C^{(k)}} \omega(f) \phi(f(\mathbf{x}_1, \dots, \mathbf{x}_k)) \leq 0.$$

Equivalently, a probability distribution μ over $C^{(k)} \subseteq \text{Pol}^{(k)}(\phi)$

$$\mathbb{E}_{f \sim \mu} [\phi(f(\mathbf{x}_1, \dots, \mathbf{x}_k))] \leq \text{avg}\{\phi(\mathbf{x}_1), \dots, \phi(\mathbf{x}_k)\}.$$

$\phi : D^m \rightarrow \overline{\mathbb{Q}}$ is **submodular** if for all $\mathbf{x}_1, \mathbf{x}_2 \in D^m$,

$$\phi(\min(\mathbf{x}_1, \mathbf{x}_2)) + \phi(\max(\mathbf{x}_1, \mathbf{x}_2)) - \phi(\mathbf{x}_1) - \phi(\mathbf{x}_2) \leq 0$$

Weighted Polymorphisms

Let $\phi : D^m \rightarrow \overline{\mathbb{Q}}$ be a function and let $C \subseteq \text{Pol}(\phi)$ be a clone of operations. A k -ary weighting $\omega : C^{(k)} \rightarrow \mathbb{Q}$ ($\sum_{f \in C^{(k)}} \omega(f) = 0, \omega(f) < 0$ only if f a projection) is a k -ary **weighted polymorphism** of ϕ if for any $\mathbf{x}_1, \dots, \mathbf{x}_k \in \text{Feas}(\phi)$

$$\sum_{f \in C^{(k)}} \omega(f) \phi(f(\mathbf{x}_1, \dots, \mathbf{x}_k)) \leq 0.$$

Equivalently, a probability distribution μ over $C^{(k)} \subseteq \text{Pol}^{(k)}(\phi)$

$$\mathbb{E}_{f \sim \mu} [\phi(f(\mathbf{x}_1, \dots, \mathbf{x}_k))] \leq \text{avg}\{\phi(\mathbf{x}_1), \dots, \phi(\mathbf{x}_k)\}.$$

$\phi : D^m \rightarrow \overline{\mathbb{Q}}$ is **submodular** if for all $\mathbf{x}_1, \mathbf{x}_2 \in D^m$,

$$\phi(\min(\mathbf{x}_1, \mathbf{x}_2)) + \phi(\max(\mathbf{x}_1, \mathbf{x}_2)) - \phi(\mathbf{x}_1) - \phi(\mathbf{x}_2) \leq 0$$

$$\omega_{\text{sub}}(\min) = \omega_{\text{sub}}(\max) = +1 \text{ and } \omega_{\text{sub}}(e_1^{(2)}) = \omega_{\text{sub}}(e_2^{(2)}) = -1$$

Weighted Polymorphisms

Let $\phi : D^m \rightarrow \overline{\mathbb{Q}}$ be a function and let $C \subseteq \text{Pol}(\phi)$ be a clone of operations. A k -ary weighting $\omega : C^{(k)} \rightarrow \mathbb{Q}$ ($\sum_{f \in C^{(k)}} \omega(f) = 0$, $\omega(f) < 0$ only if f a projection) is a k -ary **weighted polymorphism** of ϕ if for any $\mathbf{x}_1, \dots, \mathbf{x}_k \in \text{Feas}(\phi)$

$$\sum_{f \in C^{(k)}} \omega(f) \phi(f(\mathbf{x}_1, \dots, \mathbf{x}_k)) \leq 0.$$

Equivalently, a probability distribution μ over $C^{(k)} \subseteq \text{Pol}^{(k)}(\phi)$

$$\mathbb{E}_{f \sim \mu} [\phi(f(\mathbf{x}_1, \dots, \mathbf{x}_k))] \leq \text{avg}\{\phi(\mathbf{x}_1), \dots, \phi(\mathbf{x}_k)\}.$$

$\phi : D^m \rightarrow \overline{\mathbb{Q}}$ is **submodular** if for all $\mathbf{x}_1, \mathbf{x}_2 \in D^m$,

$$\phi(\min(\mathbf{x}_1, \mathbf{x}_2)) + \phi(\max(\mathbf{x}_1, \mathbf{x}_2)) - \phi(\mathbf{x}_1) - \phi(\mathbf{x}_2) \leq 0$$

$$\omega_{\text{sub}}(\min) = \omega_{\text{sub}}(\max) = +1 \text{ and } \omega_{\text{sub}}(e_1^{(2)}) = \omega_{\text{sub}}(e_2^{(2)}) = -1$$

$$\mu_{\text{sub}}(\min) = \mu_{\text{sub}}(\max) = \frac{1}{2}$$

Valued CSPs \Leftrightarrow Weighted Clones

[Cohen, Cooper, Creed, Jeavons, Ž. SICOMP'13]

This Talk

Valued CSPs \Leftrightarrow Weighted Clones

[Cohen, Cooper, Creed, Jeavons, Ž. SICOMP'13]

Galois Connection

$\text{wRelClone}(\Gamma)$ closure of Γ under scaling, addition, min

Galois Connection

$wRelClone(\Gamma)$ closure of Γ under scaling, addition, min
 $wClone(W)$ smallest weighted clone containing W

Galois Connection

$wRelClone(\Gamma)$ closure of Γ under scaling, addition, min
 $wClone(W)$ **smallest weighted clone** containing W

Galois Connection

$wRelClone(\Gamma)$ closure of Γ under scaling, addition, min
 $wClone(W)$ smallest weighted clone containing W

Galois Connection

$wRelClone(\Gamma)$	closure of Γ under scaling, addition, min
$wClone(W)$	smallest weighted clone containing W
$wPol(\Gamma)$	weighted polymorphisms of Γ

Galois Connection

$\text{wRelClone}(\Gamma)$	closure of Γ under scaling, addition, min
$\text{wClone}(W)$	smallest weighted clone containing W
$\text{wPol}(\Gamma)$	weighted polymorphisms of Γ
$\text{Imp}(W)$	functions ϕ with $\omega \in \text{wPol}(\phi)$ for all $\omega \in W$

Galois Connection


$\text{wRelClone}(\Gamma)$	closure of Γ under scaling, addition, min
$\text{wClone}(W)$	smallest weighted clone containing W
$\text{wPol}(\Gamma)$	weighted polymorphisms of Γ
$\text{Imp}(W)$	functions ϕ with $\omega \in \text{wPol}(\phi)$ for all $\omega \in W$

Theorem [Cohen, Cooper, Creed, Jeavons, Ž. SICOMP'13]


For any finite Γ , $\text{Imp}(\text{wPol}(\Gamma)) = \text{wRelClone}(\Gamma)$.

For any finite W , $\text{wPol}(\text{Imp}(W)) = \text{wClone}(W)$.

Galois Connection in Picture 1



Galois Connection in Picture 2



Complexity of VCSP(Γ)

1. $\text{VCSP}(\Gamma') \equiv_p \text{VCSP}(\Gamma)$,
 $\Gamma' = \text{wRelClone}(\Gamma)$ closure of Γ under $+$, scaling, min
2. weighted relational clones $\text{wRelClone}(\Gamma)$
 \approx weighted clones $\text{wPol}(\Gamma)$
3. properties of $\text{wPol}(\Gamma)$

Complexity of VCSP(Γ)

1. $\text{VCSP}(\Gamma') \equiv_p \text{VCSP}(\Gamma)$,
 $\Gamma' = \text{wRelClone}(\Gamma)$ closure of Γ under $+$, scaling, min
2. weighted relational clones $\text{wRelClone}(\Gamma)$
 \approx weighted clones $\text{wPol}(\Gamma)$
3. properties of $\text{wPol}(\Gamma)$

Example: $\text{wPol}(\Gamma)$ contains $\omega_{\text{sub}} \Rightarrow \text{VCSP}(\Gamma)$ tractable

[Iwata et al. JACM'01, Schrijver JCTB'00]

Tractability

A weighted clone W with support clone C is **tractable** if $\text{VCSP}(\text{Imp}(W))$ belongs to PTIME.

Tractability

A weighted clone W with support clone C is **tractable** if $\text{VCSP}(\text{Imp}(W))$ belongs to PTIME.

Easy to show:

- ▶ If $C = J_D$ (projections on D) then either $W = W_C$ or $W = W_C^0$, both are NP-hard.
- ▶ If $W = W_C^0$ for some C then W is NP-hard.

[Cohen, Cooper, Creed, Jeavons, Ž. SICOMP'13]

Tractability

A weighted clone W with support clone C is **tractable** if $\text{VCSP}(\text{Imp}(W))$ belongs to PTIME.

Easy to show:

- ▶ If $C = J_D$ (projections on D) then either $W = W_C$ or $W = W_C^0$, both are NP-hard.
- ▶ If $W = W_C^0$ for some C then W is NP-hard.

[Cohen, Cooper, Creed, Jeavons, Ž. SICOMP'13]

Consequently, unless W is NP-hard W contains a **nontrivial** weighting ω , i.e., ω assigns a positive weight to a non-projection.

Idempotency

Wlog we can restrict to weighted clones W which are:

- ▶ **surjective**

for every unary $\omega \in W$, $\omega(f) > 0 \Rightarrow f$ bijection [Thapper & Ž. '14]

Idempotency

Wlog we can restrict to weighted clones W which are:

- ▶ **surjective**

for every unary $\omega \in W$, $\omega(f) > 0 \Rightarrow f$ bijection [Thapper & Ž. '14]

- ▶ **idempotent**

for every $\omega \in W$, $\omega(f) > 0 \Rightarrow f$ idempotent [Ochremiak '14]
(thus, the only $\omega(f) > 0$ in unary $\omega \in W$ are projections)

Necessary Conditions for Tractability

Theorem [*Creed & Ž. CP'11/SICOMP'13*]

Any weighted clone W containing a nontrivial weighting contains a weighting that assigns positive weight to either:

1. A set of unary operations that are not projections; or
2. A set of binary idempotent operations that are not projections; or
3. A set of ternary operations that are majority operations, minority operations, Pixley operations or semiprojections; or
4. A set of k -ary semiprojections (for some $k > 3$).

Necessary Conditions for Tractability

Theorem [*Creed & Ž. CP'11/SICOMP'13*]

Any weighted clone W containing a nontrivial weighting contains a weighting that assigns positive weight to either:

1. A set of unary operations that are not projections; or
2. A set of binary idempotent operations that are not projections; or
3. A set of ternary operations that are majority operations, minority operations, Pixley operations or semiprojections; or
4. A set of k -ary semiprojections (for some $k > 3$).

Proof: follows the proof of Rosenberg's classification.

Necessary Conditions for Tractability

Theorem [Thapper & Ž. '14]

Any weighted clone W containing a nontrivial weighting contains a weighting that assigns positive weight to either:

1. A set of unary operations that are not projections; or
2. A set of binary idempotent operations that are not projections; or
3. A set of ternary operations that are either:
 - 3.1 a set of majority operations; or
 - 3.2 a set of minority operations; or
 - 3.3 a set of majority operations with total weight 2 and a set of minority operations with total weight 1; or
4. A set of k -ary semiprojections (for some $k \geq 3$).

Necessary Conditions for Tractability

Theorem [Thapper & Ž. '14]

Any weighted clone W containing a nontrivial weighting contains a weighting that assigns positive weight to either:

1. A set of unary operations that are not projections; or
2. A set of binary idempotent operations that are not projections; or
3. A set of ternary operations that are either:
 - 3.1 a set of majority operations; or
 - 3.2 a set of minority operations; or
 - 3.3 a set of majority operations with total weight 2 and a set of minority operations with total weight 1; or
4. A set of k -ary semiprojections (for some $k \geq 3$).

(1) Pixley/semi out, (2) interplay of majorities and minorities.

Necessary Conditions for Tractability

Theorem [Thapper & Ž. '14]

Any weighted clone W containing a nontrivial weighting contains a weighting that assigns positive weight to either:

1. A set of unary operations that are not projections; or
2. A set of binary idempotent operations that are not projections; or
3. A set of ternary operations that are either:
 - 3.1 a set of majority operations; or
 - 3.2 a set of minority operations; or
 - 3.3 a set of majority operations with total weight 2 and a set of minority operations with total weight 1; or
4. A set of k -ary semiprojections (for some $k \geq 3$).

Necessary Conditions for Tractability

Theorem [Thapper & Ž. '14]

Any weighted clone W containing a nontrivial weighting contains a weighting that assigns positive weight to either:

1. A set of unary operations that are not projections; or
2. A set of binary idempotent operations that are not projections; or
3. A set of ternary operations that are either:
 - 3.1 a set of majority operations; or
 - 3.2 a set of minority operations; or
 - 3.3 a set of majority operations with total weight 2 and a set of minority operations with total weight 1; or
4. A set of k -ary semiprojections (for some $k \geq 3$).

Proof: Gordan's Theorem (duality of linear programming).

Necessary Conditions for Tractability

Theorem [Thapper & Ž. '14]

Any weighted clone W containing a nontrivial weighting contains a weighting that assigns positive weight to either:

1. A set of unary operations that are not projections; or
2. A set of binary idempotent operations that are not projections; or
3. A set of ternary operations that are either:
 - 3.1 a set of majority operations; or
 - 3.2 a set of minority operations; or
 - 3.3 a set of majority operations with total weight 2 and a set of minority operations with total weight 1; or
4. A set of k -ary semiprojections (for some $k \geq 3$).

Necessary Conditions for Tractability

Theorem [Thapper & Ž. '14]

Any weighted clone W containing a nontrivial weighting contains a weighting that assigns positive weight to either:

1. A set of unary operations that are not projections; or
2. A set of binary idempotent operations that are not projections; or
3. A set of ternary operations that are either:
 - 3.1 a set of majority operations; or
 - 3.2 a set of minority operations; or
 - 3.3 a set of majority operations with total weight 2 and a set of minority operations with total weight 1; or
4. A set of k -ary semiprojections (for some $k \geq 3$).

Easy: not possible/sufficient for tractability!

Sufficient Conditions for Tractability

Theorem [Thapper & Ž. '14]

Any weighted clone W containing a nontrivial weighting contains a weighting that assigns positive weight to either:

1. A set of unary operations that are not projections; or
2. A set of binary idempotent operations that are not projections; or
3. A set of ternary operations that are either:
 - 3.1 a set of majority operations; or
 - 3.2 a set of minority operations; or
 - 3.3 a set of majority operations with total weight 2 and a set of minority operations with total weight 1; or
4. A set of k -ary semiprojections (for some $k \geq 3$).

Sufficient Conditions for Tractability

Theorem [Thapper & Ž. '14]

Any weighted clone W containing a nontrivial weighting contains a weighting that assigns positive weight to either:

1. A set of unary operations that are not projections; or
2. A set of binary idempotent operations that are not projections; or
3. A set of ternary operations that are either:
 - 3.1 a set of majority operations; or
 - 3.2 a set of minority operations; or
 - 3.3 a set of majority operations with total weight 2 and a set of minority operations with total weight 1; or
4. A set of k -ary semiprojections (for some $k \geq 3$).

Easy: sufficient for tractability as $\text{Imp}(W)$ contains only relations!

Sufficient Conditions for Tractability

Only near-unanimity operations/only edge operations!

Theorem [Thapper & Z. '14]

Any weighted clone W containing a nontrivial weighting contains a weighting that assigns positive weight to either:

1. A set of unary operations that are not projections; or
2. A set of binary idempotent operations that are not projections; or
3. A set of ternary operations that are either:
 - 3.1 a set of majority operations; or
 - 3.2 a set of minority operations; or
 - 3.3 a set of majority operations with total weight 2 and a set of minority operations with total weight 1; or
4. A set of k -ary semiprojections (for some $k \geq 3$).

Easy: sufficient for tractability as $\text{Imp}(W)$ contains only relations!

Sufficient Conditions for Tractability

Theorem [Thapper & Ž. '14]

Any weighted clone W containing a nontrivial weighting contains a weighting that assigns positive weight to either:

1. A set of unary operations that are not projections; or
2. A set of binary idempotent operations that are not projections; or
3. A set of ternary operations that are either:
 - 3.1 a set of majority operations; or
 - 3.2 a set of minority operations; or
 - 3.3 a set of majority operations with total weight 2 and a set of minority operations with total weight 1; or
4. A set of k -ary semiprojections (for some $k \geq 3$).

Sufficient Conditions for Tractability

Theorem [Thapper & Ž. '14]

Any weighted clone W containing a nontrivial weighting contains a weighting that assigns positive weight to either:

1. A set of unary operations that are not projections; or
2. A set of binary idempotent operations that are not projections; or
3. A set of ternary operations that are either:
 - 3.1 a set of majority operations; or
 - 3.2 a set of minority operations; or
 - 3.3 a set of majority operations with total weight 2 and a set of minority operations with total weight 1; or
4. A set of k -ary semiprojections (for some $k \geq 3$).

special case (2 maj & 1 minor) tractable [Kolmogorov & Ž. JACM'13]

Necessary Conditions for Tractability

Theorem [Thapper & Ž. '14]

Any weighted clone W containing a nontrivial weighting contains a weighting that assigns positive weight to either:

1. A set of unary operations that are not projections; or
2. A set of binary idempotent operations that are not projections; or
3. A set of ternary operations that are either:
 - 3.1 a set of majority operations; or
 - 3.2 a set of minority operations; or
 - 3.3 a set of majority operations with total weight 2 and a set of minority operations with total weight 1; or
4. A set of k -ary semiprojections (for some $k \geq 3$).

Holds true for any support clone $C \neq J_D$!

Necessary Conditions for Finite-Valued Tractability

Theorem [Thapper & Ž. '14]

Any weighted clone W containing a nontrivial weighting contains a weighting that assigns positive weight to either:

1. A set of unary operations that are not projections; or
2. A set of binary idempotent operations that are not projections; or
3. A set of ternary operations that are either:
 - 3.1 a set of majority operations; or
 - 3.2 a set of minority operations; or
 - 3.3 a set of majority operations with total weight 2 and a set of minority operations with total weight 1; or
4. A set of k -ary semiprojections (for some $k \geq 3$).

Easy: if the support clone $C = \mathcal{O}_D$ the only possible case

Complexity of Finite-Valued CSPs

Theorem [Thapper & Ž. STOC'13]

Let W be an idempotent weighted clone with the support clone $C = \mathcal{O}_D$ for some finite D .

1. Either W contains a binary weighting that assigns positive weight to **commutative** operations, in which case W is tractable;
2. or W is NP-hard.

Complexity of Finite-Valued CSPs

Theorem [Thapper & Ž. STOC'13]

Let W be an idempotent weighted clone with the support clone $C = \mathcal{O}_D$ for some finite D .

1. Either W contains a binary weighting that assigns positive weight to **commutative** operations, in which case W is tractable;
2. or W is NP-hard.

$C = \mathcal{O}_D$ means that functions in $\text{Imp}(W)$ are \mathbb{Q} -valued

Necessary Conditions for Conservative Tractability

Theorem [Thapper & Ž. '14]

Any weighted clone W containing a nontrivial weighting contains a weighting that assigns positive weight to either:

1. A set of unary operations that are not projections; or
2. A set of binary idempotent operations that are not projections; or
3. A set of ternary operations that are either:
 - 3.1 a set of majority operations; or
 - 3.2 a set of minority operations; or
 - 3.3 a set of majority operations with total weight 2 and a set of minority operations with total weight 1; or
4. A set of k -ary semiprojections (for some $k \geq 3$).

Easy: if all unary functions $\subseteq \text{Imp}(W)$ the only possible cases

Complexity of Conservative Valued CSPs

Theorem [Kolmogorov & Ž. JACM'13]

Let W be weighted clone on D such that $\text{Imp}(W)$ contains all unary functions (equivalently, all $\{0, 1\}$ -valued unary fns).

1. Either Γ admits a conservative binary multimorphism and a conservative ternary multimorphism and there is a family M of 2-element subsets of D , such that:
 - ▶ for every $\{a, b\} \in M$, $\omega|_{\{a,b\}}$ is a symmetric tournament pair and
 - ▶ for every $\{a, b\} \notin M$, $\omega'|_{\{a,b\}}$ is an MJNin which case W is tractable;
2. or W is NP-hard.

Complexity of Conservative Valued CSPs

Theorem [Kolmogorov & Ž. JACM'13]

Let W be weighted clone on D such that $\text{Imp}(W)$ contains all unary functions (equivalently, all $\{0, 1\}$ -valued unary fns).

1. Either Γ admits a conservative binary multimorphism and a conservative ternary multimorphism and there is a family M of 2-element subsets of D , such that:
 - ▶ for every $\{a, b\} \in M$, $\omega|_{\{a,b\}}$ is a symmetric tournament pair and
 - ▶ for every $\{a, b\} \notin M$, $\omega'|_{\{a,b\}}$ is an MJNin which case W is tractable;
2. or W is NP-hard.

multimorphism: $\omega(f) \in \mathbb{N}$ and $\omega(e_i^{(k)}) = -1$

STP: binary mm with f dual of g and both conservative commutative

MJN: ternary mm with 2 majority and 1 minority operations

Weighted Clones on $|D| = 2$


“Necessary Theorem” gives 9 nontrivial weightings,
which are atoms in the lattice of weighted clones.

8 are tractable, 1 is NP-hard.

[Creed & Ž. CP'11/SICOMP'13]


Weighted Clones on $|D| = 2$ with $C = \mathcal{O}_D$

From the 9 before 4 are atoms, 2 generate everything.




Weighted Clones on $|D| = 2$ with $C = \mathcal{O}_D$, cont'd

Uncountably many weighted clones above W_{\max} (and W_{\min}).



Future Work, Open Problems, Where to Learn More?

The complexity of valued constraint satisfaction



[Jeavons, Krokhin, Ž. '14]

Thank You



<http://zivny.cz/>