

IX.3 Hardy spaces on the unit disc

Definition. Let $\Omega \subset \mathbb{C}$ be an open set and $u : \Omega \rightarrow [-\infty, +\infty)$ be a function. The function u is said to be **subharmonic**, if it is upper semicontinuous and, moreover, whenever $a \in \Omega$ and $R > 0$ are such that $\overline{U(a, R)} \subset \Omega$, it holds

$$u(a) \leq \frac{1}{2\pi} \int_{-\pi}^{\pi} u(a + Re^{it}) dt$$

and the integral on the right-hand side is not equal to $-\infty$.

Remark: Similarly one can define **superharmonic** functions (they are lower semicontinuous, have values in $(-\infty, +\infty]$, satisfy the opposite inequality and the respective integrals are not $+\infty$). Then a function is harmonic if and only if it is simultaneously subharmonic and superharmonic.

Theorem 13. Let $\Omega \subset \mathbb{C}$ be a domain and let f be a holomorphic function on Ω which is not the constant zero function. Then the functions $\log |f|$, $\log^+ |f|$ and $|f|^p$ ($p \in (0, +\infty)$) are subharmonic on Ω .

Remark: In the above theorem we set $\log 0 = -\infty$ and $\log^+ t = (\log t)^+ = \max\{\log t, 0\}$ for $t \in [0, \infty)$.

Theorem 14. Let $\Omega \subset \mathbb{C}$ be an open set and u be a subharmonic function on Ω . Let $a \in \Omega$ and $R > 0$ be such that $\overline{U(a, R)} \subset \Omega$. Let h be a function continuous on $\overline{U(a, R)}$ and harmonic on $U(a, R)$. If $u \leq h$ on the circle $|z - a| = R$, then $u \leq h$ on $U(a, R)$.

Notation:

- $\mathbb{D} = U(0, 1) = \{z \in \mathbb{C} : |z| < 1\}$

For $f \in H(\mathbb{D})$ and $r \in [0, 1)$ set:

- $M_0(f, r) = \exp \left(\frac{1}{2\pi} \int_{-\pi}^{\pi} \log^+ |f(re^{i\theta})| d\theta \right)$
- $M_p(f, r) = \left(\frac{1}{2\pi} \int_{-\pi}^{\pi} |f(re^{i\theta})|^p d\theta \right)^{1/p}$ ($0 < p < \infty$)
- $M_\infty(f, r) = \sup_{\theta \in [-\pi, \pi)} |f(re^{i\theta})|$

Theorem 15. Let $f \in H(\mathbb{D})$.

- The function $r \mapsto M_p(f, r)$ is non-decreasing on $[0, 1)$ for any $p \in [0, \infty]$.
- The function $p \mapsto M_p(f, r)$ is non-decreasing on $(0, \infty]$ for any $r \in (0, 1)$.
- $M_0(f, r)^p \leq 1 + M_p(f, r)^p$ for any $p \in (0, \infty)$ and $r \in (0, 1)$.

Definition.

- For $f \in H(\mathbb{D})$ and $p \in [0, \infty]$ set

$$\|f\|_p = \sup_{r \in [0, 1)} M_p(f, r) = \lim_{r \rightarrow 1-} M_p(f, r).$$

- For $p \in (0, \infty]$ set

$$H^p = \{f \in H(\mathbb{D}) : \|f\|_p < \infty\}.$$

- Further, set

$$N = \{f \in H(\mathbb{D}) : \|f\|_0 < \infty\}.$$

Remark. $H^p \subset H^s \subset N$ whenever $0 < s < p \leq \infty$.

Lemma 16. Let $f \in N$. Then there are $g, h \in H(\mathbb{D})$ such that $f = g \cdot h$, $\|g\|_\infty \leq 1$, h has no roots in $\mathbb{D}$, $h \in N$ and $\|h\|_p = \|f\|_p$ for each $p \in [0, \infty]$.

Lemma 17. Let $f \in H^p$.

- If $p \geq 1$, then $M_\infty(f, r) \leq \frac{1}{1-r} \|f\|_1 \leq \frac{1}{1-r} \|f\|_p$ for $r \in (0, 1)$.
- If $p \in (0, 1)$, then $M_\infty(f, r) \leq \frac{3}{(1-r)^{1+\frac{1}{p}}} \|f\|_p$ for $r \in (0, 1)$.

Theorem 18.

- $(H^p, \|\cdot\|_p)$ is a Banach space for any $p \in [1, \infty]$.
- If $p \in (0, 1)$, then H^p is a complete metric linear space (hence an F -space) with the metric defined by the formula $\rho_p(f, g) = \|f - g\|_p^p$.

Theorem 19. Let $f \in H(\mathbb{D})$ satisfy

$$f(z) = \sum_{n=0}^{\infty} a_n z^n, \quad z \in \mathbb{D}.$$

Then

$$\|f\|_2^2 = \sum_{n=0}^{\infty} |a_n|^2.$$

If moreover $f \in H^2$, then the following assertions hold:

- (1) The limit $f^*(e^{it}) = \lim_{r \rightarrow 1-} f(re^{it})$ exists for almost all $t \in [0, 2\pi)$.
- (2) $f^* \in L^2(\mathbb{T})$
- (3) For $n \in \mathbb{Z}$ define $\varphi_n(e^{it}) = e^{int}$, $t \in [-\pi, \pi)$. Then $(\varphi_n)_{n \in \mathbb{Z}}$ is an orthonormal basis of $L^2(\mathbb{T})$ and the expansion of f^* with respect to this basis is

$$f^* = \sum_{n=0}^{\infty} a_n \varphi_n.$$

- (4) $\lim_{r \rightarrow 1-} \frac{1}{2\pi} \int_{-\pi}^{\pi} |f^*(e^{it}) - f(re^{it})|^2 dt = 0$.
- (5) $f = P[f^*]$
- (6) Let γ be the positively oriented unit circle. Then

$$f(z) = \frac{1}{2\pi i} \int_{\gamma} \frac{f^*(w)}{w - z} dw, \quad z \in \mathbb{D}.$$

Corollary. H^2 is a Hilbert space and the mapping $f \mapsto f^*$ is a linear isometry of H^2 onto the closed linear subspace of $L^2(\mathbb{T})$ generated by the functions φ_n , $n \geq 0$. (This subspace is formed by those $g \in L^2(\mathbb{T})$, whose coefficients at φ_n , $n < 0$, in the expansion with respect to the orthonormal basis φ_n , $n \in \mathbb{Z}$ vanish, i.e., by those g which satisfy $\hat{g}(n) = 0$ for $n < 0$.)

Theorem 20. Let $p \in [1, \infty]$ and $f \in H^p$. Then the limit $f^*(e^{it}) = \lim_{r \rightarrow 1-} f(re^{it})$ exists for almost all $t \in [0, 2\pi)$ and, moreover, the following assertions hold.

- (1) $f^* \in L^p(\mathbb{T})$
- (2) $\|f^*\|_p = \|f\|_p$
- (3) $f = P[f^*]$
- (4) Let γ be the positively oriented unit circle. Then

$$f(z) = \frac{1}{2\pi i} \int_{\gamma} \frac{f^*(w)}{w - z} dw, \quad z \in \mathbb{D}.$$

- (5) If $p < \infty$, then $\lim_{r \rightarrow 1-} \frac{1}{2\pi} \int_{-\pi}^{\pi} |f^*(e^{it}) - f(re^{it})|^p dt = 0$.

Theorem 21. Let $f \in H(\mathbb{D})$ and $p \in [1, \infty]$. Then $f \in H^p$ if and only if there exists $g \in L^p(\mathbb{T})$ such that $\hat{g}(n) = 0$ for $n < 0$ and $f = P[g]$.