

$$\int \frac{x^4 + 7x^3 + 16x^2 + 14x + 7}{(x+3)(x+2)^2} dx = \int \underbrace{x + \frac{2x+7}{(x+3)(x+2)^2}}_I dx$$

deleni:

$$\begin{array}{r} (x^4 + 7x^3 + 16x^2 + 14x + 7) : (x^3 + 7x^2 + 16x + 12) = x \\ -(x^4 + 7x^3 + 16x^2 + 12x) \\ \hline 2x + 7 \end{array}$$

$$\int \frac{2x+7}{(x+3)(x+2)^2} dx = \int \frac{A}{x+3} + \frac{B}{x+2} + \frac{C}{(x+2)^2} dx =$$

$$2x+7 = A(x+2)^2 + B(x+3)(x+2) + C(x+3)$$

$$x = -2 \quad -4+7 = C \quad C = 3$$

$$x = -3 \quad -6+7 = A \quad A = 1$$

$$x = 0 \quad 7 = 4 + B \cdot 6 + 3 \cdot 3 \quad -6 = 6B \quad B = -1$$

$$= \int \frac{1}{x+3} - \frac{1}{x+2} + \frac{3}{(x+2)^2} dx = \log|x+3| - \log|x+2| + 3 \frac{-1}{x+2}$$

boltomady

$$I \subseteq \frac{x^2}{2} + \log|x+3| - \log|x+2| - \frac{3}{x+2}$$

$$x \in (-\infty, -3), (-3, -2) \\ (-2, \infty)$$

$$\int \frac{x^3 - 2x^2 + 7x + 6}{(x^2 + x + 4)(x^2 - 3x + 2)} dx = \int \frac{A}{x-2} + \frac{B}{x-1} + \frac{Cx+D}{x^2+x+4} dx$$

\downarrow
 $(x-2)(x-1)$

$x \neq 1, x \neq 2$

$$x^3 - 2x^2 + 7x + 6 = A(x-1)(x^2+x+4) + B(x-2)(x^2+x+4) + (Cx+D)(x-2)(x-1)$$

$$= Ax^3 + 3Ax - 4A + Bx^3 - Bx^2 + 2Bx - 8B + Cx^3 - 3Cx^2 + 2Cx + Dx^2 - 3Dx + 2D$$

$$x^3: A + B + C = 1$$

$$x^2: -2 = -B - 3C + D$$

$$x: 7 = 3A + 2B + 2C - 3D$$

$$1: 6 = -4A - 8B + 2D$$

$$\begin{pmatrix} A & B & C & D & | & \\ 1 & 1 & 1 & 0 & | & 1 \\ 0 & -1 & -3 & 1 & | & -2 \\ 3 & 2 & 2 & -3 & | & 7 \\ -2 & -4 & 0 & 1 & | & 3 \end{pmatrix} \sim$$

$$\begin{pmatrix} 1 & 1 & 1 & 0 & | & 1 \\ 0 & -1 & -3 & 1 & | & -2 \\ 0 & -1 & -1 & -3 & | & 4 \\ 0 & -2 & 2 & 1 & | & 5 \end{pmatrix} \sim \begin{pmatrix} 1 & 1 & 1 & 0 & | & 1 \\ 0 & 1 & 3 & -1 & | & 2 \\ 0 & 0 & 2 & -4 & | & 8 \\ 0 & 0 & 4 & 7 & | & -3 \end{pmatrix} \sim \begin{pmatrix} 1 & 1 & 1 & 0 & | & 1 \\ 0 & 1 & 3 & -1 & | & 2 \\ 0 & 0 & 1 & -2 & | & 3 \\ 0 & 0 & 0 & 15 & | & -15 \end{pmatrix}$$

$$D = -1 \quad C = 1 \quad B = -2 \quad A = 2$$

Integrally

$$\int \frac{2}{x-2} dx = 2 \log|x-2|$$

$$\int \frac{-2}{x-1} dx = -2 \log|x-1|$$

$$\int \frac{x-1}{x^2+x+4} dx = \frac{1}{2} \int \frac{2x-2}{x^2+x+4} dx = \frac{1}{2} \int \frac{2x+1}{x^2+x+4} dx + \frac{1}{2} \int \frac{-3}{x^2+x+4} dx$$

$$\frac{1}{2} \log(x^2+x+4)$$

$$\downarrow$$

$$-\frac{3}{2} \int \frac{1}{(x+\frac{1}{2})^2 + \frac{15}{4}} dx$$

$$\rightarrow -\frac{3}{2} \int \frac{4}{15} \cdot \frac{1}{\left(\frac{x+\frac{1}{2}}{\frac{\sqrt{5}}{2}}\right)^2 + 1} dx = -\frac{2}{5} \int \frac{1}{\left(\frac{2x+1}{\sqrt{5}}\right)^2 + 1} dx$$

$$= -\frac{2}{5} \cdot \frac{\sqrt{5}}{2} \arctan\left(\frac{2x+1}{\sqrt{5}}\right)$$

cellem

$$\int \stackrel{c}{=} 2 \log|x-2| - 2 \log|x-1| - \frac{\sqrt{5}}{5} \arctan\left(\frac{2x+1}{\sqrt{5}}\right)$$

$$x \in \underline{\underline{(-\infty, 1), (1, 2), (2, \infty)}}$$