

KERNEL REGRESSION

$(y_1, x_1), \dots, (y_n, x_n)$ i.i.d

I. "GLOBAL" POLYNOMIALS

$$m(x) = E[Y_i | X_i = x] = \beta_0 + \beta_1 x + \beta_2 x^2 + \dots + \beta_p x^p$$

II. Regressogram

- breaks: $a_0 < a_1 < a_2 \dots < a_L$

$$\hat{m}_n^{(R)}(x) = \text{For } x \in (a_l, a_{l+1}]$$

↑
estm. of $m(x)$

$$\hat{m}_n^{(R)}(x) = \frac{\sum_{i: x_i \in (a_l, a_{l+1}]} y_i}{\#\{i: x_i \in (a_l, a_{l+1}]\}}$$

↑
estm. of $m(x)$

III. Nadaraya - Watson's estimator

- ~~piecewise~~ locally constant fitting

- for a kernel K : $\hat{m}_n^{(NW)}(x) = \sum_{i=1}^n w_{ni}(x) y_i$

$$\text{where } w_{ni}(x) = \frac{K\left(\frac{x_i - x}{h_n}\right)}{\sum_{i=1}^n K\left(\frac{x_i - x}{h_n}\right)}, \quad K\text{-kernel, } h_n\text{-bandwidth}$$

$$\text{if } K(x) = \frac{1}{2} \mathbb{1}\{|x| < 1\}$$

$$\hat{m}_n^{(NW)}(x) = \frac{\sum_{i: x_i \in (x-h_n, x+h_n)} y_i}{\#\{i: x_i \in (x-h_n, x+h_n)\}}$$

MEAN
AVERAGING

IV. Local - linear estimator

- $\hat{m}_n^{(LL)}(x) = \hat{\beta}_0(x)$, where

$$(\hat{\beta}_0(x), \hat{\beta}_1(x)) = \underset{\beta_0, \beta_1}{\operatorname{argmin}} \sum_{i=1}^n (y_i - \beta_0 - \beta_1(x_i - x))^2 K\left(\frac{x_i - x}{h_n}\right)$$

WEIGHTED REGRESSION

$$\text{if } K(x) = \frac{1}{2} \mathbb{1}\{|x| < 1\}$$

$$(\hat{\beta}_0(x), \hat{\beta}_1(x)) = \underset{\beta_0, \beta_1}{\operatorname{argmin}} \sum_{i: x_i \in (x-h_n, x+h_n)} (y_i - \beta_0 - \beta_1(x_i - x))^2$$

VI. CHOICE OF h_n

VI (A) - $h_n^{(CV)}$ = argmin $CV(h_n)$,
 $h_n > 0$

where $CV(h_n) = \frac{1}{n} \sum_{i=1}^n [Y_i - \hat{m}_{h_n}^{(-i)}(x_i)]^2$,
 ESTIM OF $m(x)$ based on $(\frac{y_1}{x_1}, \dots, (\frac{y_{i-1}}{x_{i-1}}, \frac{y_{i+1}}{x_{i+1}}, \dots, (\frac{y_n}{x_n}))$

VI (B) - NEAREST-NEIGHBOUR BW CHOICE

- Let $K(x) > 0 \Leftrightarrow x \in (-1, 1)$
- then Choose h_n so that $\#\{i: |x_i - x| \leq h_n\} \geq 2$

$\leadsto h_n^{(NN)}(x)$ is the 2-th smallest value from $\{|x_1 - x|, \dots, |x_n - x|\}$

CHOICE OF h - Transform X_i to R_i , use CV on $(\frac{y_1}{R_1}, \dots, (\frac{y_n}{R_n}))$. 2

VII. LOWESS FUNCTION

- "robustified" kernel regression

TRICKY KERNEL

- 0 step: $\hat{m}^{(0)}(x)$ is calculated as $\hat{m}_{h_n}^{(LL)}(x)$ with $h_n = \lfloor n^{\frac{1}{3}} \rfloor$

FOR $k = 1, \dots, \text{ITER}$

$$r_i^{(k)} = Y_i - \hat{m}^{(k-1)}(x_i)$$

$$\hat{\sigma}_i^{(k)} = B \left(\frac{r_i^{(k)}}{\text{med}\{|r_1^{(k)}|, \dots, |r_n^{(k)}|\}} \right), \quad i = 1, \dots, n$$

$$\hat{m}^{(k)}(x) = \hat{\beta}_0^{(k)}(x), \quad \text{where}$$

$$(\hat{\beta}_0^{(k)}(x), \hat{\beta}_1^{(k)}(x)) = \text{argmin}_{(b_0, b_1)} \sum_{i=1}^n (Y_i - b_0 - b_1(x_i - x))^2 \cdot K\left(\frac{x_i - x}{h_n}\right) \hat{\sigma}_i^{(k)}$$

VII. CONDITIONAL VARIANCE ESTIMATION $T^2(\cdot) = \text{var}(Y|X)$

1. STEP: ESTIMATE $\mu(X) = E[Y|X=X]$
AND CALCULATE RESIDUALS $\hat{\epsilon}_i = Y_i - \hat{\mu}(X_i)$

2. STEP, ESTIMATE $\text{var}(Y|X)$ BY FITTING KERNEL
REGRESSION TO $\begin{pmatrix} \hat{\epsilon}_1^2 \\ X_1 \end{pmatrix} \dots \begin{pmatrix} \hat{\epsilon}_n^2 \\ X_n \end{pmatrix}$