

Lattice - Introduction

Tomáš Koblár

23. March 2012

Lattice

Lattice -
Introduction

Tomáš Koblár

Lattice and its
basis

Minkowski's
Theorem

Computational
problems




Figure: Lattice of NaCl

Definition of Lattice

Lattice -
Introduction

Tomáš Koblár

Lattice and its
basis

Minkowski's
Theorem

Computational
problems

Definition

A lattice is set points

$$L(v_1 \dots v_n) = \{v_1 a_1 + \dots + v_n a_n \mid a_i \in \mathbb{Z} \forall i \in \{1, \dots, n\}\},$$

where v_1, \dots, v_n are linearly independent vectors in \mathbb{R}^m . These vectors are called a basis of L .

Definition of Lattice

Lattice -
Introduction

Tomáš Koblitz

Lattice and its
basis

Minkowski's
Theorem

Computational
problems

Definition

A lattice is set points

$$L(v_1 \dots v_n) = \{v_1 a_1 + \dots + v_n a_n \mid a_i \in \mathbb{Z} \forall i \in \{1, \dots, n\}\},$$

where v_1, \dots, v_n are linearly independent vectors in \mathbb{R}^m . These vectors are called a basis of L .

If we define B as the $m \times n$ matrix with columns v_1, \dots, v_n , then we can write

$$L(v_1 \dots v_n) = L(B) = \{By \mid y \in \mathbb{Z}^n\}$$

Rank of lattice is n and dimension is m . If $m = n$ then the lattice is called full-rank.

Basis

Lattice -
Introduction

Tomáš Koblár

Lattice and its
basis

Minkowski's
Theorem

Computational
problems

Basis of lattice is not unique:




Figure: Two bases of the same lattices in \mathbb{R}^2

But not every n -tuple of vectors in \mathbb{R}^n generate same lattice:




Figure: Two bases of different lattices in \mathbb{R}^2

Equivalency of lattices

Lattice -
Introduction

Tomáš Koblitz

Lattice and its
basis

Minkowski's
Theorem

Computational
problems

A matrix $A \in \mathbb{Z}^{n \times n}$ is called unimodular if $\det(A) = \pm 1$.

Remark

If matrix U is unimodular then U^{-1} is unimodular and $U^{-1} \in \mathbb{Z}^{n \times n}$

Equivalency of lattices

Lattice -
Introduction

Tomáš Koblár

Lattice and its
basis

Minkowski's
Theorem

Computational
problems

A matrix $A \in \mathbb{Z}^{n \times n}$ is called unimodular if $\det(A) = \pm 1$.

Remark

If matrix U is unimodular then U^{-1} is unimodular and $U^{-1} \in \mathbb{Z}^{n \times n}$

Lemma

Two lattices $L(B_1), L(B_2)$ are the same if and only if there exists unimodular matrix U such that $B_1 = B_2 U$.

Proof of Equivalency

Lattice -
Introduction

Tomáš Koblár

Lattice and its
basis

Minkowski's
Theorem

Computational
problems

Since $L(B_1) = L(B_2)$, there exists matrix $U \in \mathbb{Z}^{n \times n}$ that $B_1 = B_2 U$. Similarly, there exist matrix $V \in \mathbb{Z}^{n \times n}$ such that $B_2 = B_1 V$.

Hence $B_2 = B_1 V = B_2 UV$, and we get $B_2^T B_2 = V^T U^T B_2^T B_2 UV$.

Taking determinants:

$$\det(B_2^T B_2) = \det(VU)^2 \det(B_2^T B_2)$$

Hence $\det(U)\det(V) = \pm 1$ and since both matrices have integer $\det(U) = \pm 1$, as required. \square

Assuming $B_1 = B_2 U$ for unimodular matrix U implies directly $L(B_1) \subseteq L(B_2)$. From remark we obtain that U^{-1} is unimodular too, so $B_1 U^{-1} = B_2$ implies $L(B_1) \supseteq L(B_2)$. We get $L(B_1) = L(B_2)$. \square

Fundamental parallelepiped

Lattice -
Introduction

Tomáš Koblitz

Lattice and its
basis

Minkowski's
Theorem

Computational
problems

Definition

A fundamental parallelepiped for lattice $L(B)$ is defined as

$$P(B) = \{Bx \mid x \in [0, 1)^n\}.$$

We define determinant of lattice $\det L(B) = |\det(B)|$.

Fundamental parallelepiped

Lattice -
Introduction

Tomáš Koblitz

Lattice and its
basis

Minkowski's
Theorem

Computational
problems

Definition

A fundamental parallelepiped for lattice $L(B)$ is defined as

$$P(B) = \{Bx \mid x \in [0, 1)^n\}.$$

We define determinant of lattice $\det L(B) = |\det(B)|$.

Remark

The value of determinant is the volume of the fundamental parallelepiped.

From previous lemma we easily get that two lattices $L(B_1), L(B_2)$ are the same if and only if $\det L(B_1) = \det L(B_2)$.

Successive minimum

Lattice -
Introduction

Tomáš Koblár

Lattice and its
basis

Minkowski's
Theorem

Computational
problems

Another parameter of lattice is the length of the shortest nonzero vector in the lattice. We denote this length by $\lambda_1(L)$. By length we mean Euclidean norm defined as

$$\|x\| = \sqrt{\sum_{i=1}^n x_i^2}.$$

More generally:

Definition

For lattice L of rank n and for $k \leq n$ k -th successive minimum $\lambda_k(L)$ is defined as the smallest radius of closed ball around zero contains k linearly independent vectors in lattice L .

Blichfeldt's Theorem

Lattice -
Introduction

Tomáš Koblár

Lattice and its
basis

Minkowski's
Theorem

Computational
problems

Theorem

For lattice L and set $S \subseteq \mathbb{R}^n$ with $\text{vol}(S) > \det L$ there exist nonequal points $z_1, z_2 \in S$ such that $z_1 - z_2 \in L$.

Blichfeldt's Theorem

Lattice -
Introduction

Tomáš Koblár

Lattice and its
basis

Minkowski's
Theorem

Computational
problems

Theorem

For lattice L and set $S \subseteq \mathbb{R}^n$ with $\text{vol}(S) > \det L$ there exist nonequal points $z_1, z_2 \in S$ such that $z_1 - z_2 \in L$.

Proof: Let B is some basis of L . The sets $x + P(B)$ for $x \in L$ form a partition of \mathbb{R}^n .

For every $x \in L$ we define $S_x = S \cap (x + P(B))$.

Now we define $S_x^* = S_x - x$. Then $S_x^* \subseteq P(B)$ and because

$$\sum_{x \in L} \text{vol}(S_x^*) = \sum_{x \in L} \text{vol}(S_x) = \text{vol}(S) \geq \det L = \text{vol}(P(B))$$

there are x, y in L , $x \neq y$, that $S_x^* \cap S_y^* \neq \emptyset$. Let z be a point $S_x^* \cap S_y^*$. Then $z + x \in S_x \subseteq S$ and $z + y \in S_y \subseteq S$ and $0 \neq (x + z) - (y + z) = x - y \in L$. \square

Minkowski's Theorem I.

Set $S \in \mathbb{R}^n$ is centrally-symmetric if $x \in S \Rightarrow -x \in S$

Set $S \in \mathbb{R}^n$ is convex if for any $x, y \in S$ and $\lambda \in [0, 1]$

$$\lambda x + (1 - \lambda)y \in S$$

Theorem

Let L be lattice and S central symmetric convex set in \mathbb{R}^n . If $\text{vol}(S) > 2^n \cdot \det L$ then set S contains a nonzero lattice point.

Minkowski's Theorem I.

Lattice -
Introduction

Tomáš Koblár

Lattice and its
basis

Minkowski's
Theorem

Computational
problems

Set $S \in \mathbb{R}^n$ is centrally-symmetric if $x \in S \Rightarrow -x \in S$

Set $S \in \mathbb{R}^n$ is convex if for any $x, y \in S$ and $\lambda \in [0, 1]$

$$\lambda x + (1 - \lambda)y \in S$$

Theorem

Let L be lattice and S central symmetric convex set in \mathbb{R}^n . If $\text{vol}(S) > 2^n \cdot \det L$ then set S contains a nonzero lattice point.

Proof

Define $S^ = \frac{1}{2}S = \{x \mid 2x \in S\}$. Then $\text{vol}(S^*) > \det L$.*

By Blichfeldt's theorem there exist $z_1, z_2 \in S^$ such that $0 \neq z_1 - z_2 \in L$.*

By properties of S $\frac{2z_1 - 2z_2}{2} = z_1 - z_2 \in S$, as required. \square

Minkowski's Theorem II.

Lattice -
Introduction

Tomáš Koblár

Lattice and its
basis

Minkowski's
Theorem

Computational
problems

Remark

Let $B(0, r)$ denote open ball of radius r in \mathbb{R}^n then

$$\text{vol}(B(0, r)) \geq \left(\frac{2r}{\sqrt{n}} \right)^n$$

Theorem

Let L be lattice of rank n then $\lambda_1(L) \leq \sqrt{n}(\det(L))^{1/n}$

Proof of Minkowski's theorem

Lattice -
Introduction

Tomáš Koblár

Lattice and its
basis

Minkowski's
Theorem

Computational
problems

Proof

By definition of λ_1 , $B(0, \lambda_1(L))$ contains no nonzero lattice point. From previous theorem and remark we get

$$2^n \det L \geq \text{vol}(B(0, \lambda_1(L))) \geq \left(\frac{2\lambda_1(L)}{\sqrt{n}} \right)^n.$$

By rearranging

$$\sqrt{n}(\det(L))^{1/n} \geq \lambda_1(L). \quad \square$$

Easy Computational problems

Lattice -
Introduction

Tomáš Koblár

Lattice and its
basis

Minkowski's
Theorem

Computational
problems

Membership: Given a lattice basis $B \in \mathbb{Z}^{m \times n}$ and a point $v \in \mathbb{Z}^m$, decide if $v \in L(B)$.

This problem is equivalent to question if system of m linear equation $Bx = v$ has solution in \mathbb{Z}^n . Solving by Gauss elimination.

Easy Computational problems

Lattice -
Introduction

Tomáš Koblár

Lattice and its
basis

Minkowski's
Theorem

Computational
problems

Membership: Given a lattice basis $B \in \mathbb{Z}^{m \times n}$ and a point $v \in \mathbb{Z}^m$, decide if $v \in L(B)$.

This problem is equivalent to question if system of m linear equation $Bx = v$ has solution in \mathbb{Z}^n . Solving by Gauss elimination.

Equivalence: Given $B_1, B_2 \in \mathbb{Z}^{m \times n}$, decide if $L(B_1) = L(B_2)$.

If $m=n$ we can use lemma about equivalency and compute and compare determinants of B_1 and B_2 . Otherwise we can use membership problem for all columns of B_1 and matrix (basis) B_2 and then for all columns of B_2 and matrix B_1 .

The Shortest Vector Problem - SVP

Lattice -
Introduction

Tomáš Koblár

Lattice and its
basis

Minkowski's
Theorem

Computational
problems

Search SVP: Given a basis B of lattice L find a vector $v \in L(B)$ such that $\|v\| = \lambda_1(L(B))$.

Optimization SVP: Given a basis B find $\lambda_1(L(B))$.

Decisional SVP: Given a basis B and rational r , determine if $\lambda_1(L(B)) \leq r$ or not.

Aproxiamtion variants of SVP

Lattice -
Introduction

Tomáš Koblitz

Lattice and its
basis

Minkowski's
Theorem

Computational
problems

For $\gamma \geq 1$ we define following problems:

Search SVP: Given a basis B of lattice L find a nonzero vector $v \in L(B)$ such that $\|v\| \leq \gamma \cdot \lambda_1(L(B))$.

Optimization SVP $_{\gamma}$: Given a basis B find such d that $d \leq \lambda_1(L(B)) \leq d \cdot \gamma$.

Promise SVP $_{\gamma}$: Given a basis B and rational r , determine if $\lambda_1(L(B)) \leq r$ or $\lambda_1(L(B)) < r \cdot \gamma$.

Aproxiamtion variants of SVP

Lattice -
Introduction

Tomáš Koblitz

Lattice and its
basis

Minkowski's
Theorem

Computational
problems

For $\gamma \geq 1$ we define following problems:

Search SVP: Given a basis B of lattice L find a nonzero vector $v \in L(B)$ such that $\|v\| \leq \gamma \cdot \lambda_1(L(B))$.

Optimization SVP $_{\gamma}$: Given a basis B find such d that $d \leq \lambda_1(L(B)) \leq d \cdot \gamma$.

Promise SVP $_{\gamma}$: Given a basis B and rational r , determine if $\lambda_1(L(B)) \leq r$ or $\lambda_1(L(B)) < r \cdot \gamma$.

Open question: Is the search variant harder than optimization variant ?

The Shortest Independent Vector Problem - SIVP

Lattice -
Introduction

Tomáš Koblár

Lattice and its
basis

Minkowski's
Theorem

Computational
problems

For $\gamma \geq 1$:

SIVP $_{\gamma}$: Given a basis B find n linearly independent vectors in $L(B)$ of length $\leq \gamma \cdot \lambda_n(L(B))$.

The Closest vector problem - CVP

Lattice -
Introduction

Tomáš Koblár

Lattice and its
basis

Minkowski's
Theorem

Computational
problems

For parameter $\gamma \geq 1$ we define:

Search CVP_γ : Given a lattice basis B and a vector $t \in \mathbb{Z}^n$, find $v \in L(B)$ such that $\|v - t\| \leq \gamma \cdot \text{dist}(t, L(B))$.

Optimization CVP_γ : Given a lattice basis B and a vector $t \in \mathbb{Z}^n$, find d such that $d \leq \text{dist}(t, L(B)) \leq d \cdot \gamma$.

Promise CVP_γ : Given a lattice basis B and a vector $t \in \mathbb{Z}^n$ and rational r , determine if $\text{dist}(t, L(B)) \leq r$ or $\text{dist}(t, L(B)) < r \cdot \gamma$.

Facts about complexity

Lattice -
Introduction

Tomáš Koblitz

Lattice and its
basis

Minkowski's
Theorem

Computational
problems

General SVP and CVP are NP-hard problem.

Approximating problems are probably not NP-hard but the best algorithm runs in exponential time even more there is no better quantum algorithm, which means that cryptosystems based on approximating SVP or CVP are great promise for post-quantum cryptography.

Promise CVP_γ and SVP_γ are known as GapCVP_γ respective GapSVP_γ . Both these problems are in $NP \cap coNP$ for $\gamma \approx c\sqrt{n}$ where c is some constant.

Cryptosystems based on lattices

Lattice -
Introduction

Tomáš Koblitz

Lattice and its
basis

Minkowski's
Theorem

Computational
problems

- GGH encryption / signature scheme
- NTRUEncrypt and NTRUSign

End of presentation

Lattice -
Introduction

Tomáš Koblitz

Lattice and its
basis

Minkowski's
Theorem

Computational
problems

Thank you for your attention