

Rubik cube: Introduction

Tomas Koblirle

Department of Algebra
Faculty of Mathematics and Physics
Charles University in Prague

November 23, 2014

Rubik cube: The first approach

Names of faces of the Rubik cube: **U**p,**D**own,**F**ront,**B**ack,**R**ight and **L**eft.

Rubik cube: The first approach

Names of faces of the Rubik cube: **U**p,**D**own,**F**ront,**B**ack,**R**ight and **L**eft.

The rotations of the U face is denoted as U .

Definition

Let X be a set and G be a group. We say that G **acts on the set** X on the right provided by following conditions hold:

- 1 Each g belonging to G gives rise to a function

$$\phi_g : X \rightarrow X$$

- 2 The identity of G defines the identity function on X .
- 3 For $g, h \in G$ then $\phi_{gh}(x) = \phi_h(\phi_g(x))$.

The relative position of the centers of the faces of the Rubik cube is not changed by any move.

The relative position of the centers of the faces of the Rubik cube is not changed by any move.

The Rubik cube group is

$$\langle U, D, F, B, R, L \rangle \subset S_{54}$$


The second approach

Divide and conquer:

The centers are fixed. There remain edge and corner subcubes.

Orientation of vertices

Figure: Orientation of vertices the Rubik Cube



Definition

Let G, H are groups and ϕ a homomorphism

$$\phi : H \rightarrow \text{Aut}(G).$$

We define a multiplication on the set $G \times H$ by

$$(x_1, y_1) \cdot (x_2, y_2) = (x_1 \cdot \phi(y_1)(x_2), y_1 \cdot y_2).$$


We get a group called semi-direct product, denoted $G \rtimes H$.

Group of Rubik corner subcubes

$$C_3^8 \rtimes S_8$$

Orientation of edges

Figure: Orientation of edges of the Rubik Cube



Group of Rubik edge subcubes

$$C_2^{12} \rtimes S_{12}$$

The direct product of two group S, T is the group

$$S \times T = \{(s, t) | s \in S, t \in T\}$$

with intuitive operations.

Moves change the orientation

Corner subcubes: $v : H \rightarrow C_3^8$.

$$v(F) = (2, 0, 1, 0, 1, 0, 0, 2)$$

$$v(U) = (0, 0, 0, 0, 0, 0, 0, 0)$$

$$v(R) = (1, 2, 2, 1, 0, 0, 0, 0)$$

Moves change the orientation

Edge subcubes: $v : H \rightarrow C_2^{12}$.

$$v(F) = (1, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0)$$

$$v(U) = (1, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0)$$

$$v(R) = (0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0)$$

First Fundamental Theorem

A position of the Rubik cube is determined by

- permutation of center subcubes
- permutation of edge subcubes
- permutation of corner subcubes
- flips of corner markings
- flips of edge markings

The Rubik cube group is

$$\langle U, D, F, B, R, L \rangle \subset (C_3^8 \rtimes S_8) \times (C_2^{12} \rtimes S_{12})$$

Thank you for your attention!

Any Questions? Or can I offer some interesting facts...

Interesting facts I



Ernő Rubik, born 13th July 1944 in Budapest, Hungarian inventor, architect and professor of architecture.

Inventions: Rubik cube, Rubik's tower, Rubik's magic,...

Really a Magic cube:

How many moves we have to do to solve the Rubik cube ?

Really a Magic cube:

How many moves we have to do to solve the Rubik cube ?

There is over 43×10^{18} possible positions, 43 quintillion.

Interesting facts III

Some numbers:

5,55s solved a single cube by Mats Valk

9,03s solved one-handed by Feliks Zemdeg

23.19s blindfolded(including memorization) by Marcin Kowalczyk

41 of 41 multiple blindfolded by Marcin Kowalczyk