

Structures with small orbit growth

Bertalan Bodor,
joint work with Manuel Bodirsky

TU Dresden

Špindlerův Mlýn, 6th September 2018

Main result

Theorem

The following classes of (countable) structures are equal.

Main result

Theorem

The following classes of (countable) structures are equal.

- 1 The class of all structures \mathfrak{A} for which there exist constants c and $d < 1$ so that the number of injective n -orbits of $\text{Aut}(\mathfrak{A})$ is at most cn^{dn} . ($\mathcal{K}_{\text{exp}+}$)

Theorem

The following classes of (countable) structures are equal.

- ① *The class of all structures \mathfrak{A} for which there exist constants c and $d < 1$ so that the number of injective n -orbits of $\text{Aut}(\mathfrak{A})$ is at most cn^{dn} . ($\mathcal{K}_{\text{exp}+}$)*
- ② *The class of finite coverings of reducts of unary ω -categorical structures. ($F(R(\mathcal{U}))$)*

Definition

\mathfrak{B} is a **reduct** of \mathfrak{A} if they have the same domain set; and all constants, functions and relations of \mathfrak{B} are first-order definable in \mathfrak{A} .

Reducts

Definition

\mathfrak{B} is a **reduct** of \mathfrak{A} if they have the same domain set; and all constants, functions and relations of \mathfrak{B} are first-order definable in \mathfrak{A} .

Two reducts are considered the same iff they are **interdefinable**, that is they are reducts of one another.

Reducts

Definition

\mathfrak{B} is a **reduct** of \mathfrak{A} if they have the same domain set; and all constants, functions and relations of \mathfrak{B} are first-order definable in \mathfrak{A} .

Two reducts are considered the same iff they are **interdefinable**, that is they are reducts of one another.

Notation

If \mathcal{C} is a class of structures, then $R(\mathcal{C})$ denotes the class of reducts of structures in \mathcal{C} .

Reducts and automorphism groups

Facts

Reducts and automorphism groups

Facts

- If \mathfrak{B} is a reduct of \mathfrak{A} , then $\text{Aut}(\mathfrak{B}) \supset \text{Aut}(\mathfrak{A})$.

Reducts and automorphism groups

Facts

- If \mathfrak{B} is a reduct of \mathfrak{A} , then $\text{Aut}(\mathfrak{B}) \supset \text{Aut}(\mathfrak{A})$.
- $\text{Aut}(\mathfrak{A})$: **closed** in $\text{Sym}(\mathfrak{A})$.

Reducts and automorphism groups

Facts

- If \mathfrak{B} is a reduct of \mathfrak{A} , then $\text{Aut}(\mathfrak{B}) \supset \text{Aut}(\mathfrak{A})$.
- $\text{Aut}(\mathfrak{A})$: **closed** in $\text{Sym}(\mathfrak{A})$.

Definition

\mathfrak{A} is **ω -categorical** iff for all n $\text{Aut}(\mathfrak{A})$ has finitely many n -orbits.

Reducts and automorphism groups

Facts

- If \mathfrak{B} is a reduct of \mathfrak{A} , then $\text{Aut}(\mathfrak{B}) \supset \text{Aut}(\mathfrak{A})$.
- $\text{Aut}(\mathfrak{A})$: **closed** in $\text{Sym}(\mathfrak{A})$.

Definition

\mathfrak{A} is **ω -categorical** iff for all n $\text{Aut}(\mathfrak{A})$ has finitely many n -orbits.

Theorem (Ryll-Nardzewski, Engeler, Svenonius)

For ω -categorical structures $\mathfrak{B} \mapsto \text{Aut}(\mathfrak{B})$ is a bijection between

Reducts and automorphism groups

Facts

- If \mathfrak{B} is a reduct of \mathfrak{A} , then $\text{Aut}(\mathfrak{B}) \supset \text{Aut}(\mathfrak{A})$.
- $\text{Aut}(\mathfrak{A})$: **closed** in $\text{Sym}(\mathfrak{A})$.

Definition

\mathfrak{A} is **ω -categorical** iff for all n $\text{Aut}(\mathfrak{A})$ has finitely many n -orbits.

Theorem (Ryll-Nardzewski, Engeler, Svenonius)

For ω -categorical structures $\mathfrak{B} \mapsto \text{Aut}(\mathfrak{B})$ is a bijection between

- the **reducts** of \mathfrak{A} (up to interdefinability) and

Reducts and automorphism groups

Facts

- If \mathfrak{B} is a reduct of \mathfrak{A} , then $\text{Aut}(\mathfrak{B}) \supset \text{Aut}(\mathfrak{A})$.
- $\text{Aut}(\mathfrak{A})$: **closed** in $\text{Sym}(\mathfrak{A})$.

Definition

\mathfrak{A} is **ω -categorical** iff for all n $\text{Aut}(\mathfrak{A})$ has finitely many n -orbits.

Theorem (Ryll-Nardzewski, Engeler, Svenonius)

For ω -categorical structures $\mathfrak{B} \mapsto \text{Aut}(\mathfrak{B})$ is a bijection between

- the **reducts** of \mathfrak{A} (up to interdefinability) and
- the **closed supergroups** of $\text{Aut}(\mathfrak{A})$.

Unary structures

Notation

$\mathcal{U} :=$ class of unary ω -categorical structures.

Unary structures

Notation

$\mathcal{U} :=$ class of unary ω -categorical structures.

What is in \mathcal{U} ?

Unary structures

Notation

$\mathcal{U} :=$ class of unary ω -categorical structures.

What is in \mathcal{U} ?

- Finite union of pure sets.

Unary structures

Notation

$\mathcal{U} :=$ class of unary ω -categorical structures.

What is in \mathcal{U} ?

- Finite union of pure sets.
- In terms of automorphism groups: $\prod_{i=1}^n \text{Sym}(X_i)$.

Reducts of unary structures

What is in $R(\mathcal{U})$?

Reducts of unary structures


What is in $R(\mathcal{U})$?

- We can join orbits (we are still in \mathcal{U}).


Reducts of unary structures

What is in $R(\mathcal{U})$?

- We can join orbits (we are still in \mathcal{U}).
- We can add **flips** of orbits. Example: 2 infinite cliques.



K_ω




K_ω

Reducts of unary structures

What is in $R(\mathcal{U})$?

- We can join orbits (we are still in \mathcal{U}).
- We can add **flips** of orbits. Example: 2 infinite cliques.




Reducts of unary structures

What is in $R(\mathcal{U})$?

- We can join orbits (we are still in \mathcal{U}).
- We can add **flips** of orbits. Example: 2 infinite cliques.

That is all. (...)



Definition

\mathfrak{A} is a finite covering of \mathfrak{B} if there is a $\pi : \mathfrak{A} \rightarrow \mathfrak{B}$ such that

Definition

\mathfrak{A} is a finite covering of \mathfrak{B} if there is a $\pi : \mathfrak{A} \rightarrow \mathfrak{B}$ such that

- 1 π is surjective,

Definition

\mathfrak{A} is a finite covering of \mathfrak{B} if there is a $\pi : \mathfrak{A} \rightarrow \mathfrak{B}$ such that

- ① π is surjective,
- ② $\pi^{-1}(w)$ is finite for all $w \in \mathfrak{B}$,

Definition

\mathfrak{A} is a finite covering of \mathfrak{B} if there is a $\pi : \mathfrak{A} \rightarrow \mathfrak{B}$ such that

- ① π is surjective,
- ② $\pi^{-1}(w)$ is finite for all $w \in \mathfrak{B}$,
- ③ $\text{Aut}(\mathfrak{A})$ preserves the partition $\{\pi^{-1}(w) : w \in \mathfrak{B}\}$,

Definition

\mathfrak{A} is a finite covering of \mathfrak{B} if there is a $\pi : \mathfrak{A} \rightarrow \mathfrak{B}$ such that

- ① π is surjective,
- ② $\pi^{-1}(w)$ is finite for all $w \in \mathfrak{B}$,
- ③ $\text{Aut}(\mathfrak{A})$ preserves the partition $\{\pi^{-1}(w) : w \in \mathfrak{B}\}$,
- ④ The action of $\text{Aut}(\mathfrak{A})$ on \mathfrak{B} is $\text{Aut}(\mathfrak{B})$.


Finite coverings

Examples

Finite coverings

Examples


- $\mathfrak{B} \times F$ is a finite covering of \mathfrak{B} ,



Finite coverings

Examples


- $\mathfrak{B} \times F$ is a finite covering of \mathfrak{B} ,
- $\mathfrak{B} \times F$, but each element is labelled in F (trivial covering).



Finite coverings

Examples

- $\mathfrak{B} \times F$ is a finite covering of \mathfrak{B} ,
- $\mathfrak{B} \times F$, but each element is labelled in F (trivial covering).



Notation

$F(\mathcal{C})$ is the class of finite coverings of structures in \mathcal{C} .

Reducts of finite coverings

Structures in $F(\mathcal{U})$: can be described (...).

Reducts of finite coverings

Structures in $F(\mathcal{U})$: can be described (...).
What about reducts?

Reducts of finite coverings

Structures in $F(\mathcal{U})$: can be described (...).
What about reducts?

Observations

- $R(\mathcal{U}) \subset R(F(\mathcal{U}))$.

Reducts of finite coverings

Structures in $F(\mathcal{U})$: can be described (...).
What about reducts?

Observations

- $R(\mathcal{U}) \subset R(F(\mathcal{U}))$.
- $F(R(\mathcal{U})) \subset R(F(\mathcal{U}))$.

Reducts of finite coverings

Structures in $F(\mathcal{U})$: can be described (...).
What about reducts?

Observations

- $R(\mathcal{U}) \subset R(F(\mathcal{U}))$.
- $F(R(\mathcal{U})) \subset R(F(\mathcal{U}))$.

Is there anything else?

Notation

$\mathcal{O}_n(\mathfrak{A})$ is the number of orbits of the action of $\text{Aut}(\mathfrak{A})$ on the set of **injective** n -tuples of \mathfrak{A} .

Notation

$\mathcal{O}_n(\mathfrak{A})$ is the number of orbits of the action of $\text{Aut}(\mathfrak{A})$ on the set of **injective** n -tuples of \mathfrak{A} .

Orbit growth in \mathcal{U} : $\mathcal{O}_n \approx c^n$.

Orbit growth

Notation

$\mathcal{O}_n(\mathfrak{A})$ is the number of orbits of the action of $\text{Aut}(\mathfrak{A})$ on the set of **injective** n -tuples of \mathfrak{A} .

Orbit growth in \mathcal{U} : $\mathcal{O}_n \approx c^n$.

Orbit growth in $F(\mathcal{U})$: $\mathcal{O}_n \approx$

Notation

$\mathcal{O}_n(\mathfrak{A})$ is the number of orbits of the action of $\text{Aut}(\mathfrak{A})$ on the set of **injective** n -tuples of \mathfrak{A} .

Orbit growth in \mathcal{U} : $\mathcal{O}_n \approx c^n$.

Orbit growth in $F(\mathcal{U})$: $\mathcal{O}_n \approx$

Orbit growth in $F(\mathbb{N})$: $\mathcal{O}_n \approx \#(\text{partitions of limited size}) \approx n^{dn} : d < 1$.

Orbit growth

Notation

$\mathcal{O}_n(\mathfrak{A})$ is the number of orbits of the action of $\text{Aut}(\mathfrak{A})$ on the set of **injective** n -tuples of \mathfrak{A} .

Orbit growth in \mathcal{U} : $\mathcal{O}_n \approx c^n$.

Orbit growth in $F(\mathcal{U})$: $\mathcal{O}_n \approx$

Orbit growth in $F(\mathbb{N})$: $\mathcal{O}_n \approx \#(\text{partitions of limited size}) \approx n^{dn} : d < 1$.

Why? The orbit of an injective n -tuple is determined by which of its entries are π -equivalent.

Orbit growth

Notation

$\mathcal{O}_n(\mathfrak{A})$ is the number of orbits of the action of $\text{Aut}(\mathfrak{A})$ on the set of **injective** n -tuples of \mathfrak{A} .

Orbit growth in \mathcal{U} : $\mathcal{O}_n \approx c^n$.

Orbit growth in $F(\mathcal{U})$: $\mathcal{O}_n \approx c^n n^{dn} : d < 1 \approx n^{dn} : d < 1$.

Orbit growth in $F(\mathbb{N})$: $\mathcal{O}_n \approx \#(\text{partitions of limited size}) \approx n^{dn} : d < 1$.

Why? The orbit of an injective n -tuple is determined by which if its entries are π -equivalent.

Orbit growth

Notation

$\mathcal{O}_n(\mathfrak{A})$ is the number of orbits of the action of $\text{Aut}(\mathfrak{A})$ on the set of **injective** n -tuples of \mathfrak{A} .

Orbit growth in \mathcal{U} : $\mathcal{O}_n \approx c^n$.

Orbit growth in $F(\mathcal{U})$: $\mathcal{O}_n \approx c^n n^{dn} : d < 1 \approx n^{dn} : d < 1$.

Orbit growth in $F(\mathbb{N})$: $\mathcal{O}_n \approx \#(\text{partitions of limited size}) \approx n^{dn} : d < 1$.

Why? The orbit of an injective n -tuple is determined by which if its entries are π -equivalent.

Definition

- $\mathcal{K}_{\text{exp}} := \{\mathfrak{A} : \exists c (\mathcal{O}_n(\mathfrak{A}) \leq c^n)\},$
- $\mathcal{K}_{\text{exp}+} := \{\mathfrak{A} : \exists c, d < 1 (\mathcal{O}_n(\mathfrak{A}) \leq cn^{dn})\}.$

Theorem

- ① $\mathcal{K}_{\text{exp}} = R(\mathcal{U})$,
- ② $\mathcal{K}_{\text{exp}+} = F(R(\mathcal{U}))$.

Orbit growth

Theorem

- ① $\mathcal{K}_{\text{exp}} = R(\mathcal{U})$,
- ② $\mathcal{K}_{\text{exp}+} = F(R(\mathcal{U}))$.

Remarks

- “ \supset ”: easy,

Orbit growth

Theorem

- ① $\mathcal{K}_{\text{exp}} = R(\mathcal{U})$,
- ② $\mathcal{K}_{\text{exp}+} = F(R(\mathcal{U}))$.

Remarks

- “ \supset ”: easy,
- $(2) \rightarrow (1)$: “easy”.

Orbit growth

Theorem

- ① $\mathcal{K}_{\text{exp}} = R(\mathcal{U})$,
- ② $\mathcal{K}_{\text{exp}+} = F(R(\mathcal{U}))$.

Remarks

- “ \supset ”: easy,
- $(2) \rightarrow (1)$: “easy”.

Corollary

The class $F(R(\mathcal{U}))$ is closed under taking reducts, and thus $F(R(\mathcal{U})) = R(F(\mathcal{U}))$.

Orbit growth

Theorem

- ① $\mathcal{K}_{\text{exp}} = R(\mathcal{U})$,
- ② $\mathcal{K}_{\text{exp}+} = F(R(\mathcal{U}))$.

Remarks

- “ \supset ”: easy,
- $(2) \rightarrow (1)$: “easy”.

Corollary

The class $F(R(\mathcal{U}))$ is closed under taking reducts, and thus $F(R(\mathcal{U})) = R(F(\mathcal{U}))$.

Corollary

$\mathcal{K}_{\text{exp}+} = F(R(\mathcal{U}))$ is the smallest nontrivial class of countable structures which is closed under R , F , and C (adding finitely many constants).

Orbit growth

Ideas for the proof of $\mathcal{K}_{exp} = R(\mathcal{U})$

Primitive case:

Theorem (Macpherson, 1984)

If $G \leq \text{Sym}(X)$, G primitive, but not highly transitive, then $\mathcal{O}_n(G) \geq \frac{n!}{p(n)}$ for some polynomial p .

Orbit growth

Ideas for the proof of $\mathcal{K}_{\text{exp}} = R(\mathcal{U})$

Primitive case:

Theorem (Macpherson, 1984)

If $G \leq \text{Sym}(X)$, G primitive, but not highly transitive, then $\mathcal{O}_n(G) \geq \frac{n!}{p(n)}$ for some polynomial p .

Corollary

We are done in the primitive case. That is if $\mathfrak{A} \in \mathcal{K}_{\text{exp}}$ (or in $\mathcal{K}_{\text{exp}+}$), and $\text{Aut}(\mathfrak{A})$ is primitive, then $\text{Aut}(\mathfrak{A}) = \text{Sym}(\mathfrak{A})$.

Orbit growth

Ideas for the proof of $\mathcal{K}_{exp} = R(\mathcal{U})$

More general case:

Setting:

- $G \leq \text{Sym}(X)$ closed,

Orbit growth

Ideas for the proof of $\mathcal{K}_{exp} = R(\mathcal{U})$

More general case:

Setting:

- $G \leq \text{Sym}(X)$ closed,
- G is not primitive,

Orbit growth

Ideas for the proof of $\mathcal{K}_{exp} = R(\mathcal{U})$

More general case:

Setting:

- $G \leq \text{Sym}(X)$ closed,
- G is not primitive,
- G has no finite orbits,

Orbit growth

Ideas for the proof of $\mathcal{K}_{exp} = R(\mathcal{U})$

More general case:

Setting:

- $G \leq \text{Sym}(X)$ closed,
- G is not primitive,
- G has no finite orbits,
- $\mathcal{O}_n(G) \leq 2^n$.

Orbit growth

Ideas for the proof of $\mathcal{K}_{exp} = R(\mathcal{U})$

More general case:

Setting:

- $G \leq \text{Sym}(X)$ closed,
- G is not primitive,
- G has no finite orbits,
- $\mathcal{O}_n(G) \leq 2^n$.

Steps:

- There is one nontrivial partition of G with finitely many classes: X_1, X_2 .

We need $G \geq \text{Sym}(X_1) \times \text{Sym}(X_2)$.

Orbit growth

Ideas for the proof of $\mathcal{K}_{exp} = R(\mathcal{U})$

More general case:

Setting:

- $G \leq \text{Sym}(X)$ closed,
- G is not primitive,
- G has no finite orbits,
- $\mathcal{O}_n(G) \leq 2^n$.

Steps:

- There is one nontrivial partition of G with finitely many classes: X_1, X_2 .
We need $G \geq \text{Sym}(X_1) \times \text{Sym}(X_2)$.
- We show that $G|_{X_i}$ is primitive and closed $\rightarrow G|_{X_i} = \text{Sym}(X_i)$.

Orbit growth

Ideas for the proof of $\mathcal{K}_{exp} = R(\mathcal{U})$

More general case:

Setting:

- $G \leq \text{Sym}(X)$ closed,
- G is not primitive,
- G has no finite orbits,
- $\mathcal{O}_n(G) \leq 2^n$.

Steps:

- There is one nontrivial partition of G with finitely many classes: X_1, X_2 .
We need $G \geq \text{Sym}(X_1) \times \text{Sym}(X_2)$.
- We show that $G|_{X_i}$ is primitive and closed $\rightarrow G|_{X_i} = \text{Sym}(X_i)$.
- Then if $G \not\geq \text{Sym}(X_1) \times \text{Sym}(X_2)$, then there is bijection $e : X_1 \rightarrow X_2$ such that $\{(x, e(x))\}$ is an invariant partition.

Not possible because of orbit growth.

Thomas' conjecture

Conjecture (Thomas, 1991)

Every homogeneous structure over a finite relational language has finitely many reducts.

Thomas' conjecture

Conjecture (Thomas, 1991)

Every homogeneous structure over a finite relational language has finitely many reducts.

Solved for several individual structures, but not much is known in general.

Thomas' conjecture

Conjecture (Thomas, 1991)

Every homogeneous structure over a finite relational language has finitely many reducts.

Solved for several individual structures, but not much is known in general.

Theorem

Every structure in $\mathcal{K}_{\text{exp}+}$ has finitely many reducts.

Thomas' conjecture

Conjecture (Thomas, 1991)

Every homogeneous structure over a finite relational language has finitely many reducts.

Solved for several individual structures, but not much is known in general.

Theorem

Every structure in $\mathcal{K}_{\text{exp}+}$ has finitely many reducts.

We need:

- $\mathcal{K}_{\text{exp}+} = F(R(\mathcal{U}))$,
- A quasi-characterization of structures in $F(R(\mathcal{U}))$.

Definition

$\text{CSP}(\mathfrak{A})$ is the following computational problem:

Definition

CSP(\mathfrak{A}) is the following computational problem:

- Input: primitive positive formula φ in \mathfrak{A} ($\varphi \equiv \exists\exists\ldots\exists(\wedge(\text{atomic}))$)

Definition

CSP(\mathfrak{A}) is the following computational problem:

- Input: primitive positive formula φ in \mathfrak{A} ($\varphi \equiv \exists\exists\dots\exists(\wedge(\text{atomic}))$)
- Question: Is φ true?

Definition

$\text{CSP}(\mathfrak{A})$ is the following computational problem:

- Input: primitive positive formula φ in \mathfrak{A} ($\varphi \equiv \exists\exists\dots\exists(\wedge(\text{atomic}))$)
- Question: Is φ true?

Dichotomy conjecture: $\text{CSP}(\mathfrak{A})$ is either in **P** or **NP**-complete.

Future work: CSPs in \mathcal{K}_{exp+}

Definition

$\text{CSP}(\mathfrak{A})$ is the following computational problem:

- Input: primitive positive formula φ in \mathfrak{A} ($\varphi \equiv \exists\exists\ldots\exists(\bigwedge(\text{atomic}))$)
- Question: Is φ true?

Dichotomy conjecture: $\text{CSP}(\mathfrak{A})$ is either in **P** or **NP**-complete.

Theorem (Bulatov, 2017; Zhuk 2017,)

The dichotomy conjecture is true if \mathfrak{A} is finite.

Future work: CSPs in \mathcal{K}_{exp+}

Definition

$\text{CSP}(\mathfrak{A})$ is the following computational problem:

- Input: primitive positive formula φ in \mathfrak{A} ($\varphi \equiv \exists \exists \dots \exists (\wedge (\text{atomic}))$)
- Question: Is φ true?

Dichotomy conjecture: $\text{CSP}(\mathfrak{A})$ is either in **P** or **NP**-complete.

Theorem (Bulatov, 2017; Zhuk 2017,)

The dichotomy conjecture is true if \mathfrak{A} is finite.

Theorem (Bodirsky, Mottet, 2016)

The dichotomy conjecture is true for structures in $R(\mathcal{U})$. (Using the finite dichotomy conjecture.)

Future work: CSPs in \mathcal{K}_{exp+}

What about CSPs in finite coverings?

Future work: CSPs in \mathcal{K}_{exp+}

What about CSPs in finite coverings?

Some partial results:

Future work: CSPs in \mathcal{K}_{exp+}

What about CSPs in finite coverings?

Some partial results:

- Dichotomy solved for finite coverings of the pure set assuming that the relation “ $\pi(a) = \pi(b)$ ” is p.p. definable,

Future work: CSPs in \mathcal{K}_{exp+}

What about CSPs in finite coverings?

Some partial results:

- Dichotomy solved for finite coverings of the pure set assuming that the relation “ $\pi(a) = \pi(b)$ ” is p.p. definable,
- Also solved for infinitely many edges.

Future work: CSPs in \mathcal{K}_{exp+}

What about CSPs in finite coverings?

Some partial results:

- Dichotomy solved for finite coverings of the pure set assuming that the relation “ $\pi(a) = \pi(b)$ ” is p.p. definable,
- Also solved for infinitely many edges.

Conjecture

$\mathfrak{A} \in \mathcal{K}_{exp+}$.

If $\text{Pol}(\mathfrak{A})$ contains a (canonical) pseudo-cyclic term, then $\text{CSP}(\mathfrak{A})$ is in **P**, otherwise it is **NP**-complete.

Future work

Some other questions:

- Describe all structures in $\mathcal{K}_{exp+} = F(R(U))$.

Some other questions:

- Describe all structures in $\mathcal{K}_{exp+} = F(R(U))$.
- Decidability problems in \mathcal{K}_{exp+} .

Thank you for your attention!



European Research Council

Established by the European Commission

This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 681988).